

Non-classical states of atomic ensembles: fundamentals and applications in quantum metrology

Luca Pezzè and Augusto Smerzi

*QSTAR, INO-CNR and LENS,
Largo Enrico Fermi 2, 50125 Firenze,
Italy**

Markus K. Oberthaler

*Kirchhoff-Institut für Physik,
Universität Heidelberg,
Im Neuenheimer Feld 227,
69120 Heidelberg,
Germany*

Roman Schmied and Philipp Treutlein

*Department of Physics,
University of Basel,
Klingelbergstrasse 82, 4056 Basel,
Switzerland*

(Dated: September 7, 2016)

Quantum technologies exploit entanglement to revolutionize computing, measurement, and communication. This has stimulated the research in different areas of physics to engineer and manipulate fragile many-particle entangled states. Progress has been particularly rapid for atoms. Thanks to the large and tunable nonlinearities and the well developed techniques for trapping, controlling and counting, many groundbreaking experiments have demonstrated the generation of entangled states of trapped ions, cold and ultracold gases of neutral atoms. Moreover, atoms can couple strongly to external forces and light fields, which makes them ideal for ultra-precise sensing and time keeping. All these factors call for generating non-classical atomic states designed for phase estimation in atomic clocks and atom interferometers, exploiting many-body entanglement to increase the sensitivity of precision measurements. The goal of this article is to review and illustrate successful theoretical proposals and experiments with atomic ensembles that have demonstrated many-particle entanglement and quantum-enhanced metrology.

PACS numbers: 03.67.Bg, 03.67.Mn, 03.75.Dg, 03.75.Gg, 06.20.Dk, 42.50.Dv

CONTENTS

I. Introduction	2	5. The maximum likelihood estimator	8
A. Entanglement useful for quantum metrology	4	6. Method of moments	9
B. Generation of metrologically useful entanglement in atomic ensembles	4	7. Bayesian estimation	9
C. Outline of this review	4	C. Entanglement and phase sensitivity	9
II. Fundamentals	4	1. Multiparticle entanglement	10
A. Collective spin systems	5	2. Sensitivity bound for separable states: the standard quantum limit	10
B. Phase estimation	6	3. Coherent spin states	10
1. Cramér-Rao bound and Fisher information	6	4. Useful entanglement for quantum metrology	11
2. Lower bound to the Fisher information: average moments	7	5. Metrological spin squeezing	12
3. Upper bound to the Fisher information: the quantum Fisher information	7	6. Dicke states	13
4. Phase sensitivity and statistical distance	7	7. NOON states	14
		8. Further notions of spin squeezing and their relation to entanglement	14
		9. Einstein-Podolsky-Rosen entanglement and Bell correlations	16
		D. Tomography of spin states	17
		1. Spin-noise tomography	17
		2. Visualizing spin states	17
		III. Entanglement via atomic collisions: the Bosonic Josephson Junction	18

* We highly appreciate any feedback and comments on this review. Please send these to QmetrologyRMP@gmail.com

A. Metrologically useful entanglement in the ground state of the bosonic Josephson junction	20
1. Ground state for positive nonlinearities	20
2. Ground state for negative nonlinearities	22
3. Adiabatic splitting	22
4. Experimental spin squeezing in the ground state of the BJJ	22
B. One-axis twisting dynamics	24
1. Spin squeezing and particle entanglement	25
2. Quantum interferometry	26
3. Bell correlations	26
C. Twist-and-turn dynamics	27
1. Entanglement of non-spin-squeezed states	28
D. Entanglement of two spatially-separated spinor Bose-Einstein condensates	28
IV. Entanglement via atomic collisions: spin-mixing dynamics	29
A. Spinor Bose-Einstein condensates	29
1. Spin-changing collisions	29
2. Quantum spin-mixing dynamics in the low-depletion limit	30
B. Population correlations and twin-Fock state	31
1. Number squeezing	31
2. Quantum interferometry with a twin-Fock state	31
3. Particle entanglement	32
C. Quadrature squeezing and squeezed-vacuum state	32
1. Quadrature squeezing	32
2. Interferometry with squeezed vacuum	32
3. Nonlinear SU(1,1) interferometry	33
4. Continuous variable and EPR entanglement	33
D. Other protocols to create correlated atomic pairs	34
V. Entanglement creation via atom-light interaction	35
A. Quantum state preparation using nondestructive measurements	35
1. Quantum nondemolition measurements in free space	35
2. Cavity-based quantum nondemolition measurements in the dispersive regime	37
3. Heralded atomic entanglement created by single photon detection	38
4. Cavity-based quantum nondemolition measurements in the normal mode splitting regime	39
5. Quantum state preparation via quantum Zeno dynamics	40
B. Light-mediated coherent interaction between distant atoms	41
C. Quantum state transfer from non-classical light to atoms	41
VI. Entangled states of trapped ions	42
A. Generation of metrologically useful entangled states of many ions	42
B. Quantum metrology with trapped ions	44
1. Quantum metrology with two ions	44
2. Quantum metrology with GHZ states of several ions	44
3. Spin squeezing and useful entanglement of many ions	45
VII. Working entanglement-enhanced interferometers and prospects for applications	45
A. The influence of noise and decoherence	46
1. General results	46
2. Uncorrelated decoherence	46
3. Correlated phase noise, differential interferometry, and decoherence-free subspaces	47
4. Error correction	48
5. Particle losses	48

6. Finite detection efficiency	48
B. Atomic clocks	49
1. Entanglement-assisted atomic clocks	49
2. Proposals for entanglement-assisted lattice clocks	49
C. Optical magnetometers	50
1. Entanglement-assisted optical magnetometers	51
D. Scanning-probe magnetometers using Bose-Einstein condensates	51
E. Nonlocal phase encoding	52
VIII. Outlook	52
Acknowledgments	53
References	53

I. INTRODUCTION

The most precise measurement of a physical parameter (such as the strength of a field, a force, or time) is very often obtained by mapping it to a phase that can be determined using interferometric techniques. Phase estimation is thus a unifying framework for precision measurements. It follows the general scheme outlined in Fig. 1: a probe state $\hat{\rho}_0$ of N particles is prepared, acquires a phase shift θ , and is finally detected. This conceptually simple scheme is common to all interferometric sensors: from gravitational wave detectors to atomic clocks, gyroscopes, gravimeters, just to name a few. The goal is to estimate θ with the smallest possible uncertainty $\Delta\theta$ given finite resources such as time and number of particles. The noise that determines $\Delta\theta$ can be of a technical (classical) or fundamental (quantum) nature (Braunstein and Caves, 1994; Helstrom, 1976; Holevo, 1982). Current two-mode atomic sensors are nearly—or already—limited by the so-called standard quantum limit, $\Delta\theta_{\text{SQL}} = 1/\sqrt{N}$, inherent in probes using a finite number of uncorrelated particles (Giovannetti *et al.*, 2006). This quantum noise has a statistical nature, following from the central limit theorem. Classical correlations between the probe particles do not allow to overcome this bound (Pezzè and Smerzi, 2009). Yet, the standard quantum limit is not fundamental (Caves, 1981; Wineland *et al.*, 1994; Yurke





FIG. 1 Building blocks of parameter estimation through phase estimation. Phase estimation follows four steps: *i*) preparation of the probe state $\hat{\rho}_0$; *ii*) encoding of a phase shift θ that depends on the parameter of interest; *iii*) readout (μ indicates a generic measurement result); *iv*) deriving a phase estimate $\Theta(\mu)$ from the measurement results. The uncertainty $\Delta\theta$ of the phase estimation depends crucially on all of these operations.



TRAPPED IONS

- [1] Sackett *et al.*, 2000
- [2] Meyer *et al.*, 2001
- [3] Leibfried *et al.*, 2003
- [4] Leibfried *et al.*, 2004
- [5] Leibfried *et al.*, 2005
- [15] Monz *et al.*, 2011
- [29] Bohnet *et al.*, 2015

BOSE-EINSTEIN CONDENSATES

- [6] Estève *et al.*, 2008
- [10] Gross *et al.*, 2010
- [11] Riedel *et al.*, 2010
- [16] Lücke *et al.*, 2011
- [17] Hamley *et al.*, 2012
- [19] Berrada *et al.*, 2013
- [20] Ockeloen *et al.*, 2013
- [22] Strobel *et al.*, 2014
- [24] Muessel *et al.*, 2014
- [25] Muessel *et al.*, 2015
- [30] Kruse *et al.*, 2016

COLD THERMAL ATOMS

- [7] Appel *et al.*, 2009
- [8] Leroux *et al.*, 2010a
- [9] Schleier-Smith *et al.*, 2010
- [12] Leroux *et al.*, 2010b
- [13] Louchet-Chauvet *et al.*, 2010
- [14] Chen *et al.*, 2011
- [18] Sewell *et al.*, 2012
- [21] Sewell *et al.*, 2014
- [23] Bohnet *et al.*, 2014
- [26] Barontini *et al.*, 2015
- [27] Hosten *et al.*, 2016a
- [28] Cox *et al.*, 2016

FIG. 2 **Summary of experimental achievements.** Gain over the standard quantum limit $(\Delta\theta_{\text{SQL}})^2 = 1/N$ achieved experimentally with trapped ions (black symbols), Bose-Einstein condensates (red) and cold thermal ensembles (blue). The gain is shown on logarithmic [left, dB, $10\log_{10}(\Delta\theta_{\text{SQL}}/\Delta\theta)^2$] and linear [right] scale. The solid thick line is the Heisenberg limit $(\Delta\theta_{\text{HL}})^2 = 1/N^2$. Stars refer to directly measured phase sensitivity gains, performing a full phase estimation experiment. Circles are expected gains based on a characterization of the quantum state, *e.g.* calculated as $(\Delta\theta)^2 = \xi_{\text{R}}^2/N$, where ξ_{R}^2 is the spin-squeezing parameter, or as $(\Delta\theta)^2 = 1/F_{\text{Q}}$, where F_{Q} is the quantum Fisher information, see Sec. II. Filled (open) circles indicate results obtained without (with) subtraction of technical and/or imaging noise. Every symbol is accompanied by a number (in chronological order) giving the reference reported in the side table. Here N is the total number of particles or, in presence of fluctuations, the mean total.

et al., 1986). Quantum metrology studies how to exploit quantum resources, such as squeezing and entanglement, to overcome this classical bound (Giovannetti *et al.*, 2004, 2011). Research on quantum metrology with atomic ensembles also sheds new light on fundamental questions about many-particle entanglement (Amico *et al.*, 2008; Gühne and Tóth, 2009) and related concepts, such as Einstein-Podolsky-Rosen correlations (Reid *et al.*, 2009) and Bell nonlocality (Brunner *et al.*, 2014). Since the systems of interest for quantum metrology often contain thousands or even millions of particles, it is in general not possible to address, detect, and manipulate all particles individually. Moreover, the finite number of measurements limits the possibility to fully reconstruct the gener-

ated quantum states. These limitations call for conceptually new approaches to the characterization of entanglement that rely on a finite number of coarse-grained measurements. In fact, many schemes for quantum metrology require only collective manipulations and measurements on the entire atomic ensemble. Still, the results of such measurements allow one to draw many interesting conclusions about the underlying quantum correlations between the particles.

A. Entanglement useful for quantum metrology

In the context of phase estimation, the idea that quantum correlations are necessary to overcome the standard quantum limit emerged already in pioneering works (Kitagawa and Ueda, 1993; Yurke *et al.*, 1986). A major step forward was to recognize that $\Delta\theta$ can be decreased below $\Delta\theta_{\text{SQL}}$ by using spin-squeezed states (Wineland *et al.*, 1994, 1992), and that such states are entangled (Sørensen *et al.*, 2001; Sørensen and Mølmer, 2001). However, spin-squeezed states are a relatively small class of states useful for quantum-enhanced metrology. Other prominent examples are NOON and Greenberger-Horne-Zeilinger (GHZ) states, which are not spin squeezed but can nevertheless provide phase sensitivities beating the standard quantum limit (Bollinger *et al.*, 1996; Lee *et al.*, 2002). In recent years, it has been clarified that only a special class of quantum correlations can be exploited to estimate an interferometric phase with sensitivity overcoming $\Delta\theta_{\text{SQL}}$. This class of entangled states is fully identified by the quantum Fisher information, F_Q . The condition $F_Q > N$ (Pezzè and Smerzi, 2009) is sufficient for entanglement and *necessary and sufficient for entanglement useful for quantum metrology*: it identifies the class of states that can be used to overcome the standard quantum limit in any interferometer where the phase shift is generated by a local Hamiltonian. Spin-squeezed, GHZ and NOON states fulfill the condition $F_Q > N$. Phase uncertainties down to $\Delta\theta = 1/\sqrt{kN}$ are obtained with metrologically useful k -particle entangled states (Hyllus *et al.*, 2012a; Tóth, 2012). In the absence of noise, the ultimate limit is $\Delta\theta_{\text{HL}} = 1/N$, the so-called Heisenberg limit (Giovannetti *et al.*, 2006; Holland and Burnett, 1993; Yurke *et al.*, 1986), which can be reached with metrologically useful genuine N -particle entangled states ($k = N$).

B. Generation of metrologically useful entanglement in atomic ensembles

A variety of techniques have been used to generate, with atomic ensembles, entangled states useful for quantum metrology. The crucial ingredient is interaction between the particles, for instance atom-atom collisions in Bose-Einstein condensates, atom-light interactions in cold gases, or combined electrostatic and ion-light interaction in ion chains. Figure 2 summarizes the experimental achievements (gain of phase sensitivity relative to the standard quantum limit) as a function of the number of particles. Stars in Fig. 2 show the measured phase-sensitivity gain obtained after a full interferometer sequence using entangled states as input to the atom interferometer. Filled circles report witnesses of metrologically useful entanglement (*i.e.* spin squeezing and quantum Fisher information) measured on experimen-

tally generated states, representing potential improvement in sensitivity. Open circles are inferred squeezing, being obtained after subtraction of detection noise. The Heisenberg limit has been reached with up to ~ 10 trapped ions. Attaining this ultimate bound with a much larger number of particles is beyond current technology as it requires the creation and protection of large amounts of entanglement. Nevertheless, metrological gains up to ~ 100 have been reported with large atomic ensembles. A glance at Fig. 2 reveals how quantum metrology with atomic ensembles is a very active area of research in physics. Moreover, the reported results prove that the field is now mature enough to take the step from proofs-of-principle to technological applications.

C. Outline of this review

This Review presents modern developments of phase-estimation techniques in atomic systems aided by quantum-mechanical entanglement, as well as fundamental studies of the associated entangled states. In Sec. II, we give a theoretical overview of quantum-enhanced metrology with special emphasis on the role of entanglement. We first discuss the concepts of spin squeezing and Fisher information considering spin-1/2 particles. We then illustrate different atomic systems where quantum-enhanced phase estimation—or, at least, the creation of useful entanglement for quantum metrology—has been demonstrated. Sections III and IV review the generation of entangled states in Bose-Einstein condensates. Section V describes the generation of entangled states of many atoms through the common coupling to an external light field. Section VI describes metrology with ensembles of trapped ions. Trapped ions (Blatt and Wineland, 2008; Leibfried *et al.*, 2003a), cold (Hammerer *et al.*, 2010; Ritsch *et al.*, 2013; Tanji-Suzuki *et al.*, 2011) and ultracold (Bloch *et al.*, 2008; Inguscio and Fallani, 2013; Leggett, 2001; Morsch and Oberthaler, 2006; Stamper-Kurn and Ueda, 2013) gases have their own unique characteristics regarding the generation and manipulation of entanglement. Finally, Sec. VII gives an overview of the experimentally realized entanglement-enhanced interferometers and the realistic perspective to increase the sensitivity of state-of-the-art atomic clocks and magnetometers. This section also discusses the impact of noise in the different interferometric protocols.

II. FUNDAMENTALS

In this Review we consider systems and operations involving N particles, and assume that all of their degrees of freedom are restricted to only two modes (single-particle states) that we identify as $|a\rangle$ and $|b\rangle$. These can be two hyperfine states of an atom, as in a Ramsey inter-




FIG. 3 Two-mode interferometers. In a Mach-Zehnder interferometer (a) two spatial modes $|a\rangle$ and $|b\rangle$ are combined on a balanced beam splitter, followed by a relative phase shift $\theta = \theta_a - \theta_b$ between the two arms, and finally recombined on a second balanced beam splitter. In a Ramsey interferometer (b) a resonant Rabi rotation creates a balanced superposition between two internal states $|a\rangle$ and $|b\rangle$, followed by a relative phase shift given by the energy difference between these states multiplied by the interrogation time, $\theta = (\Delta E/\hbar) \times T_R$. Finally a second resonant Rabi rotation recombines the two modes. (c) Equivalent representation of Mach-Zehnder and Ramsey interferometer operations as rotations of the collective spin on the Bloch sphere. The full sequence is equivalent to the rotation of an angle θ around the y axis.

ferometer (Ramsey, 1963), two energy levels of a trapping potential, or two spatially separated arms, as in a Mach-Zehnder interferometer (Mach, 1892; Zehnder, 1891), see Fig. 3. The interferometer operations are collective, acting on all particles in an identical way. The idealized formal description of these interferometer models is mathematically equivalent (Lee *et al.*, 2002; Wineland *et al.*, 1994): as discussed in Sec. II.A, it corresponds to the rotation of a many-particle state on the generalized Bloch sphere of maximum radius $N/2$.

A. Collective spin systems

a. Single spin. By identifying mode $|a\rangle$ with spin-up and mode $|b\rangle$ with spin-down, a (two-mode) atom can be described as an effective spin-1/2 particle: a qubit (Nielsen and Chuang, 2000; Peres, 1995). Any pure state of a single qubit can be written as $|\vartheta, \varphi\rangle = \cos \frac{\vartheta}{2} |a\rangle + e^{i\varphi} \sin \frac{\vartheta}{2} |b\rangle$, with $0 \leq \vartheta \leq \pi$ and $0 \leq \varphi < 2\pi$ the polar and azimuthal angle, respectively, in the Bloch sphere. Pure states are fully specified by $\langle \hat{\sigma} \rangle = \langle s \cdot \hat{\sigma} \rangle$, where $\hat{\sigma} = \{\hat{\sigma}_x, \hat{\sigma}_y, \hat{\sigma}_z\}$ is the Pauli vector and $s = \{\sin \vartheta \cos \varphi, \sin \vartheta \sin \varphi, \cos \vartheta\}$

is the mean spin direction. Mixed qubit states can be expressed as $\hat{\rho} = (1 + rs \cdot \hat{\sigma})/2$, and have an additional degree of freedom given by the length of the spin vector $0 \leq r \leq 1$, such that the effective state vector rs lies inside the Bloch sphere.

b. Many spins. To describe an ensemble of N distinguishable qubits, we can introduce the collective spin vector $\hat{J} = \{\hat{J}_x, \hat{J}_y, \hat{J}_z\}$, where

$$\hat{J}_x = \frac{1}{2} \sum_{l=1}^N \hat{\sigma}_x^{(l)}, \quad \hat{J}_y = \frac{1}{2} \sum_{l=1}^N \hat{\sigma}_y^{(l)}, \quad \hat{J}_z = \frac{1}{2} \sum_{l=1}^N \hat{\sigma}_z^{(l)}, \quad (1)$$

and $\hat{\sigma}^{(l)}$ is the Pauli vector of the l th particle. In particular, $\hat{J}_z$ is half the difference in the populations of the two modes. The operators (1) satisfy the angular-momentum commutation relations

$$[\hat{J}_x, \hat{J}_y] = i\hat{J}_z, \quad [\hat{J}_z, \hat{J}_x] = i\hat{J}_y, \quad [\hat{J}_y, \hat{J}_z] = i\hat{J}_x, \quad (2)$$

and have a linear degenerate spectrum spanning the 2^N -dimensional Hilbert space. The well-known set of states $|J, M\rangle$ forms a basis, where $(\hat{J}_x^2 + \hat{J}_y^2 + \hat{J}_z^2)|J, M\rangle = J(J+1)|J, M\rangle$ and $\hat{J}_z|J, M\rangle = M|J, M\rangle$, and $J \in \{N/2, N/2 - 1, \dots\}$ as well as $M \in \{-J, -J+1, \dots, +J\}$ (Zare, 1988).

c. Many spins in a symmetrized state. The Hilbert space spanned by many-qubit states symmetric under particle exchange is that of total spin $J = N/2$, which is the maximum allowed spin length for N particles.¹ It has dimension $N+1$, linearly increasing with the number of qubits. Symmetric qubit states are naturally obtained for N indistinguishable bosons, and are described by the elegant formalism developed by Schwinger in the 1950s (Biedenharn and Louck, 1981). Angular momentum operators are expressed in terms of bosonic creation, $\hat{a}^\dagger$ and $\hat{b}^\dagger$, and annihilation, $\hat{a}$ and $\hat{b}$, operators for the two modes $|a\rangle$ and $|b\rangle$:

$$\hat{J}_x = \frac{\hat{a}^\dagger \hat{b} + \hat{b}^\dagger \hat{a}}{2}, \quad \hat{J}_y = \frac{\hat{a}^\dagger \hat{b} - \hat{b}^\dagger \hat{a}}{2i}, \quad \hat{J}_z = \frac{\hat{a}^\dagger \hat{a} - \hat{b}^\dagger \hat{b}}{2}. \quad (3)$$

They satisfy the commutation relations (2) and commute with the total number of particles $\hat{N} = \hat{a}^\dagger \hat{a} + \hat{b}^\dagger \hat{b}$. The common eigenstates of $\hat{J}_z$ and $\hat{J}^2 = (\hat{N}/2)(\hat{N}/2 + 1)$

¹ Here we distinguish symmetric states $\hat{\rho}$, which satisfy $\hat{\Pi}\hat{\rho} = \hat{\rho}\hat{\Pi} = \hat{\rho}$ for any particle permutation operator $\hat{\Pi}$, from permutation-invariant states, which satisfy the weaker constraint $\hat{\rho} = \hat{\Pi}\hat{\rho}\hat{\Pi}^\dagger$ for any particle permutation operator $\hat{\Pi}$ (Tobias *et al.*, 2012; Tóth *et al.*, 2010).

are called Dicke states (Dicke, 1954) or two-mode Fock states,

$$|m_z\rangle = |N/2 + m\rangle_a |N/2 - m\rangle_b \\ = \frac{(\hat{a}^\dagger)^{N/2+m}}{\sqrt{(N/2+m)!}} \frac{(\hat{b}^\dagger)^{N/2-m}}{\sqrt{(N/2-m)!}} |0\rangle, \quad (4)$$

where $|0\rangle$ is the vacuum. They correspond to the symmetrized combinations of $N/2 + m$ particles in mode $|a\rangle$ and $N/2 - m$ particles in mode $|b\rangle$, where $m = -N/2, -N/2 + 1, \dots, N/2$. The eigenstates $|m_n\rangle$ along an arbitrary spin direction $\hat{J}_n = n \cdot \hat{J}$ can be obtained by a proper rotation of $|m_z\rangle$: $|m_x\rangle = e^{-i\frac{\pi}{2}\hat{J}_y}|m_z\rangle$ and $|m_y\rangle = e^{i\frac{\pi}{2}\hat{J}_x}|m_z\rangle$, for instance. Finally, it is useful to introduce raising and lowering operators, $\hat{J}_\pm = \hat{J}_x \pm i\hat{J}_y$ ($\hat{J}_+ = \hat{a}^\dagger\hat{b}$ and $\hat{J}_- = \hat{b}^\dagger\hat{a}$), transforming the Dicke states as $\hat{J}_\pm|m_z\rangle = \sqrt{(N/2)(N/2+1-m(m\pm 1))}|(m\pm 1)_z\rangle$.

d. Collective rotations. Any unitary transformation of a single qubit is a rotation $e^{-i\frac{\theta}{2}\hat{\sigma}_n}$ on the Bloch sphere, where n and θ are the rotation axis and rotation angle, respectively. Collective rotations of many (non-interacting) qubits are thus identified as $\otimes_{l=1}^N e^{-i\frac{\theta}{2}\hat{\sigma}_n^{(l)}} = e^{-i\theta\hat{J}_n}$ (Feynman *et al.*, 1957), and are the idealized model of most of the interferometric transformations discussed in this Review. In the collective-spin language, a balanced beam splitter is described by $e^{-i\frac{\pi}{2}\hat{J}_x}$, and a relative phase shift by $e^{-i\theta\hat{J}_z}$. Combining the three transformations, $e^{i\frac{\pi}{2}\hat{J}_x}e^{-i\theta\hat{J}_z}e^{-i\frac{\pi}{2}\hat{J}_x} = e^{-i\theta\hat{J}_y}$, the whole interferometer sequence (Mach-Zehnder or Ramsey), is equivalent to a collective rotation around the y -axis (Yurke *et al.*, 1986), see Fig. 3(c). The most sensitive states lie in a subspace with maximum spin $J = N/2$. Therefore, the most interesting states for phase estimation are the symmetric ones, invariant under particle exchange.

B. Phase estimation

Broadly speaking, an interferometer is any apparatus that transforms a probe state $\hat{\rho}_0$ depending on the value of an unknown phase parameter θ . The parameter θ cannot be measured directly and its estimation proceeds from the results of measurements performed on identical copies of the output state $\hat{\rho}_\theta$. There are good and bad choices for a measurement observable. Good ones (that we will quantify and discuss in more details below) are those characterized by a statistical distribution of measurement results that is maximally sensitive to changes of θ . We indicate as $P(\mu|\theta)$ the probability of a result²

μ given that the parameter has the value θ . The probability of observing the sequence $\mu = \{\mu_1, \dots, \mu_\nu\}$ of ν independent measurements is $P(\mu|\theta) = \prod_{i=1}^\nu P(\mu_i|\theta)$. An estimator $\Theta(\mu)$ is a generic function associating each set of measurement outcomes μ with an estimate of θ . Interference fringes of a Ramsey interferometer are a familiar example of such an estimation (they belong to a more general estimation technique known as the method of moments, discussed in II.B.6). Since the estimator is a function of random outcomes, it is itself a random variable. It is thus characterized by a θ -dependent statistical mean value $\bar{\Theta} = \sum_\mu P(\mu|\theta) \Theta(\mu)$ and variance

$$(\Delta\theta)^2 = \sum_\mu P(\mu|\theta) [\Theta(\mu) - \bar{\Theta}]^2, \quad (5)$$

the sum extending over all possible sequences of measurement results. Different estimators can yield very different results when applied to the same measured data. In the following, we will be interested in locally-unbiased estimators, for which $\bar{\Theta} = \theta$ and $\partial\bar{\Theta}/\partial\theta = 1$, so that the statistical average yields the true parameter value.

1. Cramér-Rao bound and Fisher information

How precise can a statistical estimation be? Are there any fundamental limits? A first answer came in the 1940s with the works of Cramér (1946), Rao (1945), and Fréchet (1943), who independently found a lower bound to the variance (5) of any arbitrary estimator. The Cramér-Rao bound is one of the most important results in parameter-estimation theory. For an unbiased estimator and ν independent measurements, the Cramér-Rao bound reads

$$\Delta\theta \geq \Delta\theta_{\text{CR}} = \frac{1}{\sqrt{\nu F(\theta)}}, \quad (6)$$

where

$$F(\theta) = \sum_\mu \frac{1}{P(\mu|\theta)} \left(\frac{\partial P(\mu|\theta)}{\partial\theta} \right)^2, \quad (7)$$

is the Fisher information (Fisher, 1922, 1925), the sum extending over all possible values of μ . The factor $1/\sqrt{\nu}$ in Eq. (6) is the statistical improvement when performing independent measurements on identical copies of the probe state, and, formally, it is a consequence of the additivity of the Fisher information. The Cramér-Rao bound assumes mild differentiability properties of the likelihood

set of Hermitian operators $\{\hat{E}(\mu)\}$ parametrized by μ (Nielsen and Chuang, 2000). They are positive $\hat{E}(\mu) \geq 0$ to guarantee non-negative probabilities $P(\mu|\theta) = \text{Tr}[\hat{\rho}_\theta \hat{E}(\mu)] \geq 0$, and satisfy $\sum_\mu \hat{E}(\mu) = \mathbb{1}$ to ensure normalization $\sum_\mu P(\mu|\theta) = 1$.

² In a simple scenario, μ is the eigenvalue of an observable $\hat{M}$. In a more general situation, the measurement is described by a

function $P(\mu|\theta)$ and thus holds under very general conditions, see for instance [Kay \(1993\)](#). Difficulties in saturating the Cramér-Rao bound are only present for relatively small ν . In the central limit, $\nu \gg 1$, at least one efficient estimator exists: the maximum likelihood estimator ([Fisher, 1925](#)), see Sec. [II.B.5](#).

2. Lower bound to the Fisher information: average moments

A lower bound to the Fisher information can be obtained from the rate of change with θ of specific moments of the probability distribution ([Pezzè and Smerzi, 2009](#)):

$$F(\theta) \geq \frac{1}{(\Delta\mu)^2} \left(\frac{d\bar{\mu}}{d\theta} \right)^2, \quad (8)$$

where $\bar{\mu} = \sum_{\mu} P(\mu|\theta)\mu$, and $(\Delta\mu)^2 = \sum_{\mu} P(\mu|\theta)(\mu - \bar{\mu})^2$. Equation (8) indicates that the Fisher information is more sensitive to the change of θ than the average moments of $P(\mu|\theta)$.

3. Upper bound to the Fisher information: the quantum Fisher information

An upper bound to the Fisher information is obtained by maximizing Eq. (7) over all possible generalized measurements in quantum mechanics ([Braunstein and Caves, 1994](#)), $F_Q[\hat{\rho}_\theta] = \max_{\{\hat{E}\}} F(\theta)$, called the quantum Fisher information. We have $F(\theta) \leq F_Q[\hat{\rho}_\theta]$, and the corresponding bound on the phase sensitivity for unbiased estimators and ν independent measurements is

$$\Delta\theta_{\text{CR}} \geq \Delta\theta_{\text{QCR}} = \frac{1}{\sqrt{\nu F_Q[\hat{\rho}_\theta]}}, \quad (9)$$

called the quantum Cramér-Rao bound ([Helstrom, 1967](#)). The quantum Fisher information and the quantum Cramér-Rao bound are fully determined by the interferometer output state $\hat{\rho}_\theta$. Hence they allow to calculate the optimal phase sensitivity of any given probe state and interferometer transformation ([Giovannetti et al., 2011](#); [Helstrom, 1976](#); [Holevo, 1982](#); [Paris, 2009](#); [Pezzè and Smerzi, 2014](#); [Spehner, 2014](#)). In general, the quantum Fisher information can be expressed as the variance $F_Q[\hat{\rho}_\theta] = (\Delta\hat{L})^2$ of a θ -dependent Hermitian operator $\hat{L}$ called the symmetric logarithmic derivative and defined as the solution of $\partial\hat{\rho}_\theta/\partial\theta = (\hat{\rho}_\theta\hat{L} + \hat{L}\hat{\rho}_\theta)/2$ ([Helstrom, 1967](#)). For unitary transformations generated by some Hermitian operator $\hat{H}$, $\partial\hat{\rho}_\theta/\partial\theta = i[\hat{\rho}_\theta, \hat{H}]$, an explicit expression for the quantum Fisher information³ can be found in terms of the eigenvalues $q_\kappa \geq 0$

and associated eigenvectors $|\kappa\rangle$ of the mixed probe state $\hat{\rho}_0 = \sum_{\kappa} q_\kappa |\kappa\rangle\langle\kappa|$ ([Braunstein and Caves, 1994](#); [Braunstein et al., 1996](#)):

$$F_Q[\hat{\rho}_0, \hat{H}] = 2 \sum_{\substack{\kappa, \kappa' \\ q_\kappa + q_{\kappa'} > 0}} \frac{(q_\kappa - q_{\kappa'})^2}{q_\kappa + q_{\kappa'}} |\langle\kappa|\hat{H}|\kappa'\rangle|^2. \quad (10)$$

For pure states $|\psi_0\rangle$, Eq. (10) reduces to⁴ $F_Q[|\psi_0\rangle, \hat{H}] = 4(\Delta\hat{H})^2$. For mixed states, $F_Q[\hat{\rho}_0, \hat{H}] \leq 4(\Delta\hat{H})^2$.

a. Optimal measurements. The equality $F(\theta) = F_Q[\hat{\rho}_\theta]$ can always be achieved. A possible optimal choice of measurement for both pure and mixed states is given by the set of projectors onto the eigenstates of $\hat{L}$ ([Braunstein and Caves, 1994](#)). This set of observables is necessary and sufficient for the saturation of the quantum Fisher information whenever $\hat{\rho}_\theta$ is invertible, and only sufficient otherwise. In particular, for pure states and unitary transformations, the quantum Cramér-Rao bound can be saturated, in the limit $\theta \rightarrow 0$, by a dichotomic measurement given by the projection onto the probe state itself, $|\psi_0\rangle\langle\psi_0|$, and onto the orthogonal subspace, $\mathbb{1} - |\psi_0\rangle\langle\psi_0|$. It should be noted that the symmetric logarithmic derivative, and thus also the optimal measurement, generally depends on θ , even for unitary transformations. Nevertheless, without any prior knowledge of θ , the quantum Cramér-Rao bound can be saturated in the asymptotic limit of large ν using only local measurements and adaptive estimators ([Hayashi, 2005](#)): entangled resources at the measurement stage are not necessary to achieve the quantum Cramér-Rao bound.

b. Optimal rotation direction. Given a probe state, and considering unitary transformation generated by $\hat{H} = \hat{J}_n$, it is possible to optimize the rotation direction n in order to maximize the quantum Fisher information ([Hyllus et al., 2010](#)). This optimum is given by the maximum eigenvalue of the 3×3 matrix $[\Gamma_Q]_{ij} = 2 \sum_{\kappa, \kappa' (q_\kappa + q_{\kappa'} > 0)} \frac{(q_\kappa - q_{\kappa'})^2}{q_\kappa + q_{\kappa'}} \langle\kappa'|\hat{J}_i|\kappa\rangle\langle\kappa|\hat{J}_j|\kappa'\rangle$, $i, j = x, y, z$, and the optimal direction by the corresponding eigenvector. For pure states, $[\Gamma_Q]_{ij} = 2(\langle\hat{J}_i\hat{J}_j\rangle + \langle\hat{J}_j\hat{J}_i\rangle) - 4\langle\hat{J}_i\rangle\langle\hat{J}_j\rangle$.

4. Phase sensitivity and statistical distance

Parameter estimation is naturally related to the problem of distinguishing neighboring quantum states along

³ We use the notation $F_Q[\hat{\rho}_\theta]$ to indicate the quantum Fisher information for a generic transformation of the probe state, and $F_Q[\hat{\rho}_0, \hat{H}]$ for unitary transformations.

⁴ We recall that $\Delta\hat{H} \leq (h_{\text{max}} - h_{\text{min}})/2$, where h_{max} and h_{min} are the maximum and minimum eigenvalues of $\hat{H}$ with eigenvectors $|h_{\text{max}}\rangle$ and $|h_{\text{min}}\rangle$, respectively. This bound is saturated by the state $(|h_{\text{max}}\rangle + e^{i\phi}|h_{\text{min}}\rangle)/\sqrt{2}$, with arbitrary real ϕ .




FIG. 4 **Schematic representation of different phase estimation protocols.** (a) *Maximum Likelihood estimation.* Θ_{MLE} is the absolute maximum of the likelihood function $P(\mu|\varphi)$ (black line) corresponding to the observed sequence of results μ . The phase sensitivity is identified as the mean square fluctuation $\Delta\theta_{\text{MLE}}$ of the statistical distribution of Θ_{MLE} (inset) obtained by repeating the measurements several times (at a fixed value of θ). (b) *Method of moments.* The blue line is $\bar{\mu}$ as a function of the parameter θ . A phase estimate Θ_{mom} is associated (dashed arrow) to the average value $\bar{\mu}_\nu$. The statistical uncertainty $\Delta\mu/\sqrt{\nu}$ of $\bar{\mu}_\nu$ (grey region) corresponds to a phase uncertainty $\Delta\theta_{\text{mom}}$. Equation (15) is obtained by error propagation, which locally approximates $\bar{\mu}$ by the tangent to the curve (black line). (c) *Bayesian estimation.* Posterior probability distribution $P(\varphi|\mu)$ (black line). The phase estimate Θ_B can be chosen to be the weighted averaged phase. The integral $\int_{\Delta\theta_B} d\varphi P(\varphi|\mu)$ gives the probability that the true phase value falls into the confidence interval $\Delta\theta_B$ (grey region) around Θ_B .

a path in parameter space (Braunstein and Caves, 1994; Wootters, 1981). Heuristically, the phase sensitivity of an interferometer can be understood as the smallest phase shift for which the output state $\hat{\rho}_\theta$ of the interferometer can be distinguished from the input $\hat{\rho}_0$. We introduce a statistical distance between probability distributions,

$$d_H^2(P_0, P_\theta) = 1 - \mathcal{F}_{\text{cl}}(P, P_\theta), \quad (11)$$

called the Hellinger distance, where $\mathcal{F}_{\text{cl}}(P_0, P_\theta) \equiv \sum_\mu \sqrt{P(\mu|0)P(\mu|\theta)}$ is the statistical fidelity, or overlap, between probability distributions, also known as Bhattacharyya coefficient (Bhattacharyya, 1943). d_H is non-negative, $0 \leq d_H \leq 1$, and its Taylor expansion reads

$$d_H^2(P_0, P_\theta) = \frac{F(0)}{8}\theta^2 + \mathcal{O}(\theta^3). \quad (12)$$

This equation reveals that the Fisher information is the square of a statistical speed, $v_H = \partial d_H / \partial \theta = \sqrt{F(0)}/8$. It measures the rate of change at which a probability distribution varies when tuning the phase parameter θ . Equation (12) has been used to extract the Fisher information experimentally (Strobel et al., 2014), see Sec. III.C. As Eq. (11) depends on the specific measurement, it is possible to associate different statistical distances to the same quantum states. This justifies the introduction of a distance between quantum states by maximizing $d_H^2(P_0, P_\theta)$ over all possible generalized observables, $d_B(\hat{\rho}_0, \hat{\rho}_\theta) = \max_{\{\hat{E}\}} d_H(P_0, P_\theta)$, called the Bures distance (Bures, 1969). Hübner (1992) showed that

$$d_B^2(\hat{\rho}_0, \hat{\rho}_\theta) = 1 - \mathcal{F}_Q(\hat{\rho}_0, \hat{\rho}_\theta), \quad (13)$$

where $\mathcal{F}_Q(\hat{\rho}_0, \hat{\rho}_\theta) = \sqrt{\sqrt{\hat{\rho}_\theta} \hat{\rho}_0 \sqrt{\hat{\rho}_\theta}}$ is the transition probability (Uhlmann, 1976) or the quantum fidelity between

states (Jozsa, 1994). In particular, $\mathcal{F}_Q(|\psi\rangle, |\phi\rangle) = |\langle\psi|\phi\rangle|$ for pure states. A Taylor expansion gives

$$d_B^2(\hat{\rho}_0, \hat{\rho}_\theta) = \frac{F_Q[\hat{\rho}_0]}{8}\theta^2 + \mathcal{O}(\theta^3). \quad (14)$$

The quantum Fisher information is thus the square of a quantum statistical speed, $v_Q = \partial d_B / \partial \theta = \sqrt{F_Q[\hat{\rho}_0]}/8$, maximized over all possible generalized measurements. The quantum Fisher information has also been related to the dynamical susceptibility (Hauke et al., 2016), while lower bounds have been derived by Apellaniz et al. (2015b).

5. The maximum likelihood estimator

The maximum likelihood is one of the most important estimators. It is the phase value that maximizes the likelihood of the observed measurement sequence μ , see Fig. 4(a): $\Theta_{\text{MLE}}(\mu) = \text{argmax}_\varphi P(\mu|\varphi)$. In other words, if we would set the true phase θ to Θ_{MLE} , then the probability to observe μ would be the highest. The key role played by $\Theta_{\text{MLE}}(\mu)$ in parameter estimation is due to its asymptotic properties for independent measurements. For sufficiently large ν (in the central limit), the distribution of the maximum likelihood estimator tends to a Gaussian centered at the true value θ and of variance equal to the inverse Fisher information (Lehmann and Casella, 2003): $P(\Theta_{\text{MLE}}|\theta) = \sqrt{\frac{\nu F(\theta)}{2\pi}} e^{-\frac{\nu F(\theta)}{2}(\theta - \Theta_{\text{MLE}})^2}$. Therefore, the maximum likelihood estimator is asymptotically unbiased and its variance saturates the Cramér-Rao bound: $\Delta\theta_{\text{MLE}} = 1/\sqrt{\nu F(\theta)}$. In the central limit any estimator is as good as—or worse than—the maximum likelihood estimate.

6. Method of moments

The method of moments exploits the variation of collective properties of the probability distribution—such as the mean value $\bar{\mu}$ and variance $(\Delta\mu)^2$ —with the phase shift θ . Let us take the average $\bar{\mu}_\nu = \frac{1}{\nu} \sum_{i=1}^{\nu} \mu_i$ of ν measurements results $\mu_1, \dots, \mu_\nu$. The estimator Θ_{mom} is the value for which $\bar{\mu}$ is equal to $\bar{\mu}_\nu$, see Fig. 4(b). Applying this method requires $\bar{\mu}$ to be a monotonous function of the parameter θ , at least in a local region of parameter values determined from prior knowledge. The sensitivity of this estimator can be calculated by error propagation,⁵ giving

$$\Delta\theta_{\text{mom}} = \frac{\Delta\mu}{\sqrt{\nu} |d\bar{\mu}/d\theta|}, \quad (15)$$

As expected on general grounds and proved by Eq. (8), the method of moments is not optimal in general, $\Delta\theta_{\text{mom}} \leq \Delta\theta_{\text{CR}}$, with no guarantee of saturation even in the central limit. Nevertheless, due to its simplicity, Eq. (15) is largely used in the literature to calculate the phase sensitivity of an interferometer for various input states and measurement observables (Dowling, 1998; Wineland *et al.*, 1994; Yurke *et al.*, 1986). For instance, in the case of unitary rotations generated by $\hat{H} = \hat{J}_y$ (as in Ramsey and Mach-Zehnder interferometers), and taking $\hat{J}_z$ as measurement observable, Eq. (15) in the limit $\theta \approx 0$ can be rewritten as

$$\Delta\theta_{\text{mom}} = \frac{\Delta\hat{J}_z}{\sqrt{\nu} |\langle \hat{J}_x \rangle|}. \quad (16)$$

This equation is useful to introduce the metrological spin-squeezing in Sec. II.C.5. We finally recall that Eqs. (15)-(16) are valid for a sufficiently large number of measurements (central limit) and the factor ν , often neglected in the literature, is necessarily present.

7. Bayesian estimation

The cornerstone of Bayesian inference is Bayes' theorem. Let the phase φ and the measurement results μ be random variables. Their joint probability density can be expressed as $P(\varphi, \mu) = P(\varphi|\mu)P(\mu)$ in terms of the conditional probability $P(\varphi|\mu)$ and the marginal likelihood $P(\mu) = \int d\varphi P(\varphi, \mu)$. Bayes' theorem

$$P(\varphi|\mu) = \frac{P(\mu|\varphi)P(\varphi)}{P(\mu)} \quad (17)$$

follows from the symmetry of the joint probability $P(\varphi, \mu) = P(\mu|\varphi)P(\varphi)$. When the phase is not a random variable, the Bayesian (subjective) interpretation of probabilities sees $P(\varphi)$ as the prior probability distribution for the interferometric phase to take on the value $\theta = \varphi$, *i.e.* before any measurements were done,⁶ and $P(\varphi|\mu)$ as the posterior probability distribution after taking measurement results into account. Equation (17) allows us to update our knowledge about the interferometric phase θ by including measurement results, since $P(\mu|\varphi)$ can be calculated directly (see the introduction of Sec. II.B) and $P(\mu)$ is determined by the normalization $\int_0^{2\pi} P(\varphi|\mu)d\varphi = 1$. Bayesian probabilities express our (lack of) knowledge of the interferometric phase as a probability distribution $P(\varphi|\mu)$. This is radically different from the standard frequentist view where the probability is defined as the infinite-sample limit of the outcome frequency of an observed event. Having the posterior distribution $P(\varphi|\mu)$, we can consider any phase φ as the estimate. In practice, it is convenient to choose the weighted averaged $\int_0^{2\pi} \varphi P(\varphi|\mu)d\varphi$, or the phase corresponding to the maximum of the probability $\text{argmax}_\varphi P(\varphi|\mu)$, since the corresponding mean square fluctuations saturate the Cramér-Rao bound (see below). We can further calculate the probability that the chosen estimate falls into a certain interval $[\theta_1, \theta_2]$ by integrating $\int_{\theta_1}^{\theta_2} P(\varphi|\mu)d\varphi$. To take into account the periodicity of the probability, quantities like $\int_0^{2\pi} e^{i\varphi} P(\varphi|\mu)d\varphi$ can be calculated. Remarkably, Bayesian estimation is asymptotically consistent: as the number of measurements increases, the posterior probability distribution assigns more weight in the vicinity of the true value. The Laplace-Bernstein-von Mises theorem (Gill, 2008; Lehmann and Casella, 2003; Pezzè and Smerzi, 2014) demonstrates that, under quite general conditions, $P(\varphi|\theta) = \sqrt{\frac{\nu F(\theta)}{2\pi}} e^{-\frac{\nu F(\theta)}{2}(\theta-\varphi)^2}$, to leading order in ν , for $\nu \gg 1$. In this limit, the posterior probability becomes normally distributed, centered at the true value of the parameter, and with a variance inversely proportional to the Fisher information. See (Van Trees and Bell, 2007) for a review of bounds in Bayesian phase estimation.

C. Entanglement and phase sensitivity

In this section we show how entanglement between probe particles can offer a precision enhancement in quantum metrology. We start with a formal definition of multiparticle entanglement and then clarify, via the Fisher information introduced in the previous section,

⁵ A Taylor expansion of $\bar{\mu}_\nu$ around the true value θ gives $\bar{\mu}_\nu \approx \bar{\mu} + \frac{d\bar{\mu}}{d\theta}(\Theta_{\text{mom}} - \theta)$. We obtain Eq. (15) by identifying $\bar{\mu}_\nu - \bar{\mu} \approx \Delta\mu/\sqrt{\nu}$ (valid in the central limit $\nu \gg 1$), and $\Theta_{\text{mom}} - \theta \approx \Delta\theta_{\text{mom}}$.

⁶ One often has no prior knowledge on the phase, which is expressed with a flat prior distribution $P(\varphi) = 1/(2\pi)$.

the notion of useful entanglement for quantum metrology.

1. Multiparticle entanglement

A quantum state of N particles (labeled as $l = 1, 2, \dots, N$) is fully separable if it can be written as

$$|\psi_{\text{prod}}\rangle = |\psi^{(1)}\rangle \otimes |\psi^{(2)}\rangle \otimes \dots \otimes |\psi^{(N)}\rangle, \quad (18a)$$

$$\hat{\rho}_{\text{sep}} = \sum_q p_q |\psi_{\text{prod},q}\rangle \langle \psi_{\text{prod},q}|, \quad (18b)$$

where $|\psi^{(l)}\rangle$ is the state of the l th particle. Separable states are characterized by classical correlations, established via local operations and classical communication (Werner, 1989), just by mixing product states. States that are not separable are called entangled (Horodecki *et al.*, 2009). In the case of two parties, any quantum state is either separable or entangled. For $N > 2$, we need further classifications (Dür *et al.*, 2000; Gühne and Tóth, 2009). A multiqubit state possesses bipartite entanglement if, for some qubits i and j , the two-qubit reduced density matrix is entangled. Multiparticle entanglement is quantified by means of the so-called entanglement depth (Sørensen and Mølmer, 2001), defined as the number of particles in the largest non-separable subset. In analogy with Eq. (18), a state of N particles is k -separable if it can be written as a mixture of k -producible pure states (Gühne *et al.*, 2005):

$$|\psi_{k\text{-prod}}\rangle = |\psi_{N_1}\rangle \otimes |\psi_{N_2}\rangle \otimes \dots \otimes |\psi_{N_M}\rangle \quad (19a)$$

$$\hat{\rho}_{k\text{-sep}} = \sum_q p_q |\psi_{k\text{-prod},q}\rangle \langle \psi_{k\text{-prod},q}|, \quad (19b)$$

where $|\psi_{N_l}\rangle$ is the state of $N_l \leq k$ particles and $\sum_{l=1}^M N_l = N$. A state is k -particle entangled if it is k -separable but not $(k-1)$ -separable, *i.e.* if it contains at least one state $|\psi_{N_l}\rangle$ of $N_l = k$ particles that does not factorize. Using another terminology (Sørensen and Mølmer, 2001), it has an entanglement depth larger than $k-1$. Genuine N -particle entangled states ($k=N$) cannot be factorized, each particle being entangled with all the others. Finally, note that k -producible states form a convex set containing the set of k' -producible states with $k' < k$ (Gühne and Tóth, 2009).

2. Sensitivity bound for separable states: the standard quantum limit

The quantum Fisher information of any separable state of N qubits is upper-bounded (Pezzè and Smerzi, 2009):

$$F_Q[\hat{\rho}_{\text{sep}}, \hat{J}_n] \leq N. \quad (20)$$

This inequality follows from the convexity and additivity of the quantum Fisher information and uses $4(\Delta\hat{\sigma}_n)^2 \leq 1$ (Pezzè and Smerzi, 2014). As a consequence of Eqs. (9) and (20), the maximum phase sensitivity achievable with separable states is (Giovannetti *et al.*, 2006)

$$\Delta\theta_{\text{SQL}} = \frac{1}{\sqrt{N\nu}}, \quad (21)$$

generally indicated as the shot-noise or standard quantum limit. This bound is independent of the specific measurement and estimator, and refers to unitary transformations that are local in the particles. Here, the number of particles N plays the role of a statistical gain, in complete analogy to the number of independent repetitions ν of the measurement.

3. Coherent spin states

The notion of coherent spin states was introduced by Arecchi *et al.* (1972) and Radcliffe (1971) as a generalization of the field coherent states first discussed by Glauber (1963), see Zhang *et al.* (1990) for a review. Coherent spin states are constructed as the product of N qubits (spins-1/2) in pure states all pointing along the same mean-spin direction $s = \{\sin\vartheta \cos\varphi, \sin\vartheta \sin\varphi, \cos\vartheta\}$:

$$|\vartheta, \varphi, N\rangle = \bigotimes_{l=1}^N \left[\cos\frac{\vartheta}{2} |a\rangle_l + e^{i\varphi} \sin\frac{\vartheta}{2} |b\rangle_l \right]. \quad (22)$$

Equation (22) is the eigenstate of $\hat{J}_s$ with the maximum eigenvalue of $N/2$. The coherent spin state is a product state and no quantum entanglement is present between the particles. $|\vartheta, \varphi, N\rangle$ can also be written as a binomial sum of Dicke states with $\langle m_z | \vartheta, \varphi, N \rangle = \sqrt{\binom{N}{N/2+m}} (\cos\frac{\vartheta}{2})^{N/2-m} (\sin\frac{\vartheta}{2})^{N/2+m} e^{-i(m+N/2)\varphi}$ (Arecchi *et al.*, 1972). When measuring the spin component of $|\vartheta, \varphi, N\rangle$ along any direction $\perp$ orthogonal to s , each individual atom is projected with equal probability into the up and down eigenstates along this axis, with eigenvalues $\pm 1/2$, respectively: we have $\langle \hat{J}_{\perp} \rangle = 0$, and $(\Delta\hat{J}_{\perp})^2 = N/4$, the so-called projection noise (Itano *et al.*, 1993).

Coherent spin states are optimal separable states for metrology. They saturate the equality sign in Eq. (20) and thus reach the standard quantum limit. Let us consider the rotation of $|\vartheta, \varphi, N\rangle$ around a direction n perpendicular to the mean spin direction s (here s , n and $\perp$ are mutually orthogonal). This rotation displaces the coherent spin state on the surface of the Bloch sphere, see Fig. 5(a). The initial and final states become distinguishable after rotating by an angle $\theta_{\min}$ heuristically giving the phase sensitivity of the state. This rotation angle can be obtained from a geometric reasoning (Yurke *et al.*, 1986): we have $\Delta\hat{J}_{\perp} \approx \langle \hat{J}_s \rangle \sin\theta_{\min}$, giving




FIG. 5 **Rotation of different quantum states.** Wigner distribution (normalized to its maximum value—left column), see Sec. II.D, and spin probability $P_y(m) = |\langle m_y | \psi \rangle|^2$ along the y direction (red histograms - right column) of different quantum states $|\psi\rangle$: (a) a coherent spin state pointing along the positive x -axis, $|\pi/2, 0, N\rangle$, (Sec. II.C.3); (b) a spin-squeezed state with $\xi_R^2 = 0.1$ (Sec. II.C.5); (c) a twin-Fock state (Sec. II.C.6) and (d) a NOON state (Sec. II.C.7). The thin blue histogram is $P_y(m) = |\langle m_y | e^{-i\theta \hat{J}_z} | \psi \rangle|^2$ obtained after a rotation of the state by an angle $\theta = 2/\sqrt{N}$ in (a), $\theta = 2\xi_R/\sqrt{N}$ in (b), $\theta = 2/N$ in (c), and $\theta = \pi/N$ in (d). Here $N = 100$.

$\theta_{\min} \approx 1/\sqrt{N}$ for $N \gg 1$. More rigorously, the squared Bures distance, Eq (13), between $|\vartheta, \varphi, N\rangle$ and the rotated $e^{-i\theta \hat{J}_n} |\vartheta, \varphi, N\rangle$ is

$$d_B^2(|\vartheta, \varphi, N\rangle, e^{-i\theta \hat{J}_n} |\vartheta, \varphi, N\rangle) = 1 - \cos^N(\theta/2), \quad (23)$$

that is $d_B^2 = N\theta^2/8 + \mathcal{O}(\theta^4)$ for small values of θ . According to Eq. (10) we obtain a quantum Fisher information $F_Q[|\vartheta, \varphi, N\rangle, \hat{J}_n] = 4(\Delta \hat{J}_n)^2 = N$. With the




FIG. 6 **Useful k -particle entanglement for quantum metrology.** k -separable states have a quantum Fisher information bounded by the solid blue line, Eq. (25). The dashed line is $F_Q/N = k$. Here, $N = 100$. Adapted from (Hyllus *et al.*, 2012a).

method of moments, Eq. (16), we find a phase sensitivity $\Delta\theta_{\text{mom}} = \Delta \hat{J}_\perp / (\sqrt{\nu} |\langle \hat{J}_s \rangle|) = 1/\sqrt{\nu N}$ (Itano *et al.*, 1993; Yurke *et al.*, 1986). For any rotation around an axis orthogonal to the mean spin direction, coherent spin states thus satisfy $\Delta\theta_{\text{QCR}} = \Delta\theta_{\text{SQL}}$.

4. Useful entanglement for quantum metrology

The violation of Eq. (20), *i.e.* the case where the quantum Fisher information is

$$F_Q[\hat{\rho}, \hat{J}_n] > N, \quad (24)$$

is a sufficient condition for particle-entanglement (Pezzè and Smerzi, 2009). To be more precise, inequality (24) is the condition of useful entanglement for quantum metrology (Pezzè and Smerzi, 2009): it is a necessary and sufficient condition for a quantum state to be useful in the estimation of a phase shift θ —with an interferometer implementing the transformation $e^{-i\theta \hat{J}_n}$ —with a sensitivity overcoming the standard quantum limit. Not all entangled states are useful for quantum metrology. Yet, useless entangled states for quantum metrology might be useful for other quantum technologies. It should also be noted that not all useful entangled states for quantum metrology are equally useful: large quantum Fisher information requires large entanglement depth. For states of type (19b), we have (Hyllus *et al.*, 2012a; Tóth, 2012)

$$F_Q[\hat{\rho}_{k\text{-sep}}, \hat{J}_n] \leq \mathfrak{s}k^2 + \mathfrak{r}^2, \quad (25)$$

where $\mathfrak{s} = \lfloor \frac{N}{k} \rfloor$ is the integer part of $\frac{N}{k}$, and $\mathfrak{r} = N - \mathfrak{s}k$ (note that $\mathfrak{s}k^2 + \mathfrak{r}^2 = Nk$ when N is divisible by k). If the bound (25) is surpassed, then the probe state contains metrologically useful $(k+1)$ -particle entanglement: when used as input state of the interferometer defined by the transformation $e^{-i\theta J_n}$, this state enables a phase sensitivity better than any k -separable state. The bound (25) increases monotonically with k (see Fig. 6), in particular $F_Q[\hat{\rho}_{k\text{-sep}}, \hat{J}_n] \leq Nk$. The maximum value of the quantum Fisher information is obtained for genuine N -particle entangled states, $k = N$, giving (Pezzè and Smerzi, 2009)

$$F_Q[\hat{\rho}, \hat{J}_n] \leq N^2. \quad (26)$$

Equation (26) defines the ultimate Heisenberg limit^{7 8} of phase sensitivity (Giovannetti *et al.*, 2006),

$$\Delta\theta_{\text{HL}} = \frac{1}{N\sqrt{\nu}}. \quad (27)$$

The difference between Eq. (21) and Eq. (27) is a faster scaling of the phase sensitivity with the number of particles, which cannot be obtained by classical means.⁹ We finally note that not all N -particle entangled states reach the Heisenberg limit, as exemplified by the generalized W state $|W\rangle = |\pm N/2 \mp 1\rangle_z$, which corresponds to a Dicke state with one excitation. While the W state is N -particle entangled (Dür *et al.*, 2000), its quantum Fisher information only amounts to $3N - 2$.

5. Metrological spin squeezing

Spin-squeezed states are a class of states having squeezed spin variance along a certain direction, at the cost of anti-squeezed variance along an orthogonal direction. Spin squeezing is one of the most successful

approaches to witness large-scale quantum entanglement beating the standard quantum limit in interferometry.

Let us consider the unitary rotation of a state on the Bloch sphere around an axis n perpendicular to the mean spin direction s , see Fig. 5(b), and calculate the phase sensitivity according to the error propagation formula, Eq. (16). We can write $\Delta\theta_{\text{mom}} = \xi_R/\sqrt{\nu N}$,^{10 11} where

$$\xi_R^2 = \frac{N(\Delta\hat{J}_\perp)^2}{\langle\hat{J}_s\rangle^2}, \quad (28)$$

and $\perp$ is orthogonal to both s and n . ξ_R is the spin-squeezing parameter introduced by Wineland *et al.* (1994, 1992). If $\xi_R^2 < 1$ holds, the state is said to be (metrologically) spin squeezed along the $\perp$ -axis (Wineland *et al.*, 1994, 1992) and it can be used to overcome the standard quantum limit (*i.e.* reaching $\Delta\theta_{\text{mom}} < 1/\sqrt{N}$). This requires states having spin fluctuations orthogonal to the rotation axis smaller than the projection noise, *i.e.* $(\Delta\hat{J}_\perp)^2 < N/4$, and sufficiently large spin length $\langle\hat{J}_s\rangle$.

a. Optimal spin-squeezed states. Optimal spin-squeezed states are searched among the so-called minimum uncertainty states (Aragone *et al.*, 1974; Rashid, 1978; Wódkiewicz and Eberly, 1985). These states saturate the Heisenberg uncertainty relation $(\Delta\hat{J}_n)^2(\Delta\hat{J}_\perp)^2 \geq |\langle\hat{J}_s\rangle|^2/4$, since $[\hat{J}_n, \hat{J}_\perp] = i\hat{J}_s$. We thus have $\xi_R^2 = N/[4(\Delta\hat{J}_n)^2]$ and a lower bound to ξ_R^2 is obtained by maximizing $(\Delta\hat{J}_n)^2$, giving (Agarwal and Puri, 1994; Hillery and Mlodinow, 1993)

$$\xi_R^2 \geq \frac{2}{N+2}. \quad (29)$$

This bound can be saturated by the state $\sqrt{1-\alpha^2}|0_\perp\rangle + \alpha(|1_\perp\rangle + |-1_\perp\rangle)/\sqrt{2}$ in the limit $\alpha \mapsto 0$ (Brif and Mann, 1996), where $|m_\perp\rangle$ are Dicke states defined in Sec. II.A. Notice that spin-squeezed states can achieve a Heisenberg scaling of phase sensitivity, $\Delta\theta_{\text{mom}} = \mathcal{O}(N^{-1})$, but not the Heisenberg limit (27) for $N > 2$. Optimal spin-squeezed states for even values of N and fixed values of $\langle\hat{J}_s\rangle$ are given by the ground state of the Hamiltonian $\hat{J}_\perp^2 - \lambda\hat{J}_s$, where $\lambda \geq 0$ is a Lagrange multiplier (Sørensen and Mølmer, 2001).

⁷ The name ‘‘Heisenberg limit’’ was first introduced in (Hol-land and Burnett, 1993) referring to the heuristic number-phase Heisenberg uncertainty relation $\Delta\theta\Delta N \geq 1$. We refer to the Heisenberg scaling of phase sensitivity when $\Delta\theta = \mathcal{O}(N^{-1})$.

⁸ It is possible to maximize the phase sensitivity by optimizing the number of particles N entering into the interferometer multiplied by the times ν that the measurement is performed. This provides a definition of Heisenberg limit $1/N_T$, where $N_T = N \times \nu_{\text{opt}}$, and ν_{opt} is the optimal number of measurements that maximize the phase sensitivity for a fixed number of particles N . Since ν_{opt} may depend on N , there might be, in principle, states having a Fisher information larger than N but a phase variance above the standard quantum limit $1/\sqrt{N_T}$.

⁹ Equations (21) and (27) can be generalized for transformation $e^{-i\theta \sum_{l=1}^N \hat{h}^{(l)}}$, where $\hat{h}^{(l)}$ is an arbitrary local Hamiltonian for the l th particles (that can be a generic qudit), and we take $\hat{h}^{(l)} = \hat{h}$ for all N particles. We have $\Delta\theta_{\text{SQL}} = \frac{1}{\delta h \sqrt{N\nu}}$, and $\Delta\theta_{\text{HL}} = \frac{1}{\delta h N \sqrt{\nu}}$ (Giovannetti *et al.*, 2006), where $\delta h = |h_{\text{max}} - h_{\text{min}}|$, and h_{max} and h_{min} are the maximum and minimum eigenvalues of $\hat{h}$, respectively.

¹⁰ Since this equation assumes that the phase-encoding transformation is unitary, see Sec. II.B.6, it gives an expected phase sensitivity based on squeezing properties of the states.

¹¹ In the literature, it is possible to find different notation for the metrological spin-squeezing parameter (ξ and ξ_S are also commonly used). Here we follow the notation first introduced by Wineland *et al.* (1994), and used in a previous review (Ma *et al.*, 2011). In particular, here ξ_S refers to the Kitagawa and Ueda (1993) spin-squeezing parameter, see Sec. II.C.8.




FIG. 7 **Relation between spin squeezing and quadrature squeezing.** For $\langle \hat{J}_z \rangle \approx N/2 \gg 1$, the Bloch sphere can be approximated locally as a plane orthogonal to the mean spin direction (here, the z -axis). Spin squeezing along the $\hat{J}_x$ axis is equivalent to squeezing of the $\hat{X}$ quadrature. A rotation around the y -axis is equivalent to a displacement in the $\hat{X}$ direction.

b. Spin squeezing and bosonic quadrature squeezing. In the case of quantum probe states having $N \gg 1$ and a strong population imbalance between the two modes, spin squeezing can be well approximated by single-mode quadrature-squeezing. Let $|a\rangle$ be the highly-populated mode (continuous-variable limit, $\langle \hat{a}^\dagger \hat{a} \rangle \approx N \gg 1$) and perform the approximate Holstein-Primakoff transformation (Duan *et al.*, 2002; Madsen and Mølmer, 2004; Wang and Sanders, 2003a) $\hat{J}_+/\sqrt{N} \mapsto \hat{b}$, $\hat{J}_-/\sqrt{N} \mapsto \hat{b}^\dagger$, and $2\hat{J}_z/N \mapsto 1$, formally equivalent to the mean-field replacement $\hat{a} \mapsto \sqrt{N}$. Within this approximation, the spin operators $\hat{J}_x$ and $\hat{J}_y$ map onto the position $\hat{X} = (\hat{b} + \hat{b}^\dagger)/\sqrt{2} = \sqrt{2/N}\hat{J}_x$ and momentum $\hat{P} = (\hat{b} - \hat{b}^\dagger)/(i\sqrt{2}) = \sqrt{2/N}\hat{J}_y$ quadrature operators (Scully and Zubairy, 1997), respectively, rescaled by $\sqrt{N/2}$. We thus find

$$\xi_R^2 = 2(\Delta\hat{Q})^2, \quad (30)$$

where $\hat{Q} = (\hat{b}e^{-i\phi} + \hat{b}^\dagger e^{i\phi})/\sqrt{2} = \hat{X} \cos \phi + \hat{P} \sin \phi$ and ϕ is an angle in the X - P plane. Equation (30) shows the equivalence between the metrological spin-squeezing parameter and the quadrature variance, within the approximations. In particular, the rotation $e^{-i\theta\hat{J}_y}$ maps onto a displacement of the state along the X direction in the quadrature plane by an amount $\theta \times \sqrt{N/2}$, described by $e^{-i\theta\sqrt{N/2}\hat{P}}$, see Fig. 7. When squeezing the quadrature variance below the vacuum noise limit, *i.e.* $(\Delta\hat{Q})^2 < 1/2$, it is possible to overcome the standard quantum limit of phase sensitivity, *i.e.* $\xi_R^2 < 1$. The sensitivity of interferometers using a probe state with all atoms in a single mode can be increased by feeding the other mode with a quadrature-squeezed state, as first proposed by Caves (1981) for an optical interferometer.

c. Spin squeezing and Fisher information. Spin squeezing $\xi_R < 1$ is a sufficient condition for useful particle entanglement in metrology (Sørensen *et al.*, 2001). Furthermore, Sørensen and Mølmer (2001) showed that the degree of spin squeezing is related to metrologically useful k -particle entanglement (for a given spin length, smaller and smaller values of ξ_R can be obtained by increasing the entanglement depth). The quantum Fisher information detects entanglement in a larger number of states than those recognized by metrological spin squeezing (Pezzè and Smerzi, 2009):

$$\frac{N}{F_Q[\hat{\rho}, \hat{J}_n]} \leq \xi_R^2. \quad (31)$$

This inequality, which follows from Eq. (8), shows that if a state is spin squeezed, $\xi_R^2 < 1$, it also satisfies the condition of metrologically useful entanglement $F_Q[\hat{\rho}, \hat{J}_n] > N$. The contrary is not true: there are states that are not spin squeezed and yet entangled and useful for quantum metrology. The Dicke and NOON states, discussed below, are notable examples.

6. Dicke states

Dicke states, Eq. (4), have a precise relative number of particles between the two modes and a completely undefined phase. They are not spin squeezed (Wang and Mølmer, 2002). A direct calculation of the quantum Fisher information gives

$$F_Q[|m_z\rangle, \hat{J}_\perp] = \frac{N^2}{2} - 2m^2 + N \quad (32)$$

for any rotation direction $\perp$ orthogonal to the z axis. Dicke states with $m = \pm N/2$ are coherent spin states; those with $m \neq \pm N/2$ are metrologically usefully entangled. From the perspective of quantum metrology, the most interesting Dicke state is the twin-Fock state (Holland and Burnett, 1993; Sanders and Milburn, 1995), $|N/2\rangle_a |N/2\rangle_b = |0_z\rangle$, corresponding to $N/2$ particles in each mode. It can be visualized as a ring on the equator of the Bloch sphere, see Fig. 5(c). A rotation around any axis in the x - y plane converts the well-defined number difference into a well defined relative phase between the two modes. It should be noted that the twin-Fock state has zero mean spin length. Therefore, the metrologically useful entanglement of the twin-Fock state cannot be exploited when measuring the relative number of particles. A possible phase-sensitive signal is the variance of the relative population (Kim *et al.*, 1998), see Sec. IV.B.2. The phase sensitivity calculated via the method of moments strongly depends on θ : for $\theta \approx 0$ and $N \gg 1$, we have $(\Delta\theta)^2 = \frac{2}{\nu(N^2+2N)} + \mathcal{O}(\theta^2)$, which is a factor two above the Heisenberg limit at $\theta = 0$ and remains sub-standard quantum limit for $\theta \lesssim 1/\sqrt{N}$. Similar results

can be obtained with error propagation when estimating the phase shift from the measurement of the parity operator (Campos *et al.*, 2003; Gerry *et al.*, 2004), which measures the difference in populations between even and odd eigenstates of $\hat{J}_z$ and is very difficult to implement for large spins. A θ -independent phase sensitivity can be reached when measuring the number of particles at the output ports of the interferometer and using a maximum likelihood estimator or a Bayesian method (Holland and Burnett, 1993; Krschek *et al.*, 2011; Pezzè and Smerzi, 2006).

Squeezing the number of particles at both inputs of the interferometer is not necessary to reach sub-standard quantum limit sensitivities (Pezzè and Smerzi, 2013). Let us consider a probe state $\hat{\rho} = \hat{\rho}_a \otimes |N\rangle_b \langle N|$, where $\hat{\rho}_a$ is an arbitrary state in mode $|a\rangle$ with mean particle number N_a and $|N\rangle_b$ is a Fock state of $N/2$ particles in mode $|b\rangle$. We find

$$F_Q[\hat{\rho}_a \otimes |N/2\rangle_b \langle N/2|, \hat{J}_y] = NN_a + \frac{N}{2} + N_a. \quad (33)$$

Heisenberg scaling is achieved when $N_a = N/2$, without any assumptions on $\hat{\rho}_a$. In particular, existing interferometers that operate with uncorrelated atoms can be improved by simply replacing the vacuum state in one of the two input ports by a Fock state.

7. NOON states

The Heisenberg limit can be saturated by the state

$$|\text{NOON}\rangle = \frac{|N\rangle_a |0\rangle_b + e^{i\phi} |0\rangle_a |N\rangle_b}{\sqrt{2}}, \quad (34)$$

given by a coherent superposition of all particles in mode $|a\rangle$ and all particles in mode $|b\rangle$, where ϕ is an arbitrary phase. This state is called NOON state (Lee *et al.*, 2002) when considering indistinguishable bosonic particles. When considering distinguishable particles, as ions in a Paul trap for instance, see Sec. VI, the state (34) is generally called a ‘‘Schrödinger cat’’ (Bollinger *et al.*, 1996; Leibfried *et al.*, 2005) or Greenberger-Horne-Zeilinger state (Monz *et al.*, 2011), originally introduced in (Greenberger *et al.*, 1990) for three particles. A look at the Wigner distribution of the NOON state, see Fig. 5(d,left), reveals substructures of angular size $1/N$ given by spherical harmonic contributions Y_k^q with the maximum allowed value $k = N$ (Schmied and Treutlein, 2011). Rotating the NOON state around the z -axis, the initial and final states becomes distinguishable after a rotation angle $\theta_{\min} \approx 1/N$. The squared Bures distance between the probe and the rotated state is

$$d_B^2(|\text{NOON}\rangle, e^{-i\theta \hat{J}_z} |\text{NOON}\rangle) = 1 - \cos(N\theta/2), \quad (35)$$

which oscillates in phase N time faster than the corresponding overlap for a coherent spin state, Eq. (23), and

is $d_B^2 = N^2\theta^2/8 + \mathcal{O}(\theta^4)$ for small θ (Pezzè and Smerzi, 2007). Note that the relative spin probability distribution of the NOON state, $P(m_y) = |\langle m_y | e^{-i\theta \hat{J}_z} |\text{NOON}\rangle|^2$, see Fig. 5(d,right), shows a comb-like structure as a function of m . These substructure change quickly with θ : for $\theta = 2\pi n/N$ only even values of m are populated, for $\theta = \pi(1 + 2n)/N$ only odd values of m are populated, with $n = 0, 1, \dots, N-1$. According to Eq. (10), we obtain $F_Q[|\text{NOON}\rangle, \hat{J}_z] = 4(\Delta \hat{J}_z)^2 = N^2$, and thus $\Delta\theta_{\text{QCR}} = \Delta\theta_{\text{HL}}$ for the NOON state. It is possible to reach this sensitivity via the method of moments by measuring the parity of the relative number of particles among the two modes (Bollinger *et al.*, 1996), see Sec. VI.

8. Further notions of spin squeezing and their relation to entanglement

When addressing individual qubits is not possible, or in presence of low counting statistics, entanglement criteria based on mean and variance of spin operators—as the condition $\xi_R < 1$ introduced in Sec. II.C.5—are experimentally relevant. Moreover, states of a large number of particles cannot be characterized via full state tomography: the reconstruction of the full density matrix is hindered and finally prevented by the exponential increase in the required number of measurements. In the literature, different definitions of spin squeezing for collective angular momentum operators can be found (Ma *et al.*, 2011; Tóth and Apellaniz, 2014). In the following we review the ones most relevant for the present context.

a. Squeezing parameter of Kitagawa and Ueda. A spin-1/2 particle is characterized by isotropic spin fluctuations, equal to $1/4$, along any direction orthogonal to the mean spin direction s . By adding N uncorrelated spins all pointing along the same direction s (as in a coherent spin state), we have $(\Delta \hat{J}_\perp)^2 = N/4$, where $\hat{J}_\perp$ points along an arbitrary direction orthogonal to s . Quantum correlations between spins may result in reduced fluctuations in one direction, $(\Delta \hat{J}_\perp)^2 < N/4$, at the expense of enhanced fluctuations along the other direction orthogonal to s . This suggests the introduction of the spin-squeezing parameter (Kitagawa and Ueda, 1993)

$$\xi_S^2 = \frac{4 \min_\perp (\Delta \hat{J}_\perp)^2}{N}, \quad (36)$$

$\xi_S^2 < 1$ being the spin-squeezing condition. Equation (36) is related to metrological spin squeezing via the relation $\min_\perp \xi_R^2 = [N/(2\langle \hat{J}_s \rangle)]^2 \xi_S^2$. Since $|\langle \hat{J}_s \rangle| \leq N/2$, we obtain

$$\xi_S^2 \leq \xi_R^2. \quad (37)$$

In other words, metrological spin squeezing, $\xi_R^2 < 1$, implies spin squeezing according to the definition of Kitagawa and Ueda. The converse is not true: there is




FIG. 8 ξ_S^2 versus ξ_R^2 . The metrological spin-squeezing parameter ξ_R^2 is given by the ratio between the Kitagawa-Ueda spin-squeezing parameter ξ_S^2 and the squared spin length (Ramsey contrast) $2|\langle \hat{J}_s \rangle|/N$. $\xi_R^2 < 1$ is found in the blue region upper-bounded by the standard quantum limit (upper thick black line) and lower-bounded by Eq. (29) (dashed line). The lower thick black line is the Heisenberg limit. The orange region highlights the regime of parameters showing $\xi_S^2 < 1$ but no metrological spin squeezing. The solid blue lines are bounds to ξ_R^2 obtained for k -particle entangled states (Sørensen and Mølmer, 2001). Here, $N = 1000$.

no direct relation between $\xi_S^2 < 1$ and the improvement of metrological sensitivity, as illustrated in Fig. 8. It is worth noting that the minimum in Eq. (36) is given by the smallest eigenvalue of the covariance matrix $\text{Cov}(\hat{J}_{n_i}, \hat{J}_{n_j}) = \{\langle \hat{J}_{n_i}, \hat{J}_{n_j} \rangle\}/2 - \langle \hat{J}_{n_i} \rangle \langle \hat{J}_{n_j} \rangle$, where n_1, n_2 are two mutually orthogonal directions in the plane perpendicular to s and $\{\hat{J}_{n_1}, \hat{J}_{n_2}\} = \hat{J}_{n_1} \hat{J}_{n_2} + \hat{J}_{n_2} \hat{J}_{n_1}$. Taking, without loss of generality, $\langle \hat{J}_{n_1} \rangle = \langle \hat{J}_{n_2} \rangle = 0$, we have (Wang and Sanders, 2003b)

$$\xi_S^2 = \frac{\langle \hat{J}_{n_1}^2 + \hat{J}_{n_2}^2 \rangle - \sqrt{\langle \hat{J}_{n_1}^2 - \hat{J}_{n_2}^2 \rangle^2 + \{\langle \hat{J}_{n_1}, \hat{J}_{n_2} \rangle\}^2}}{N/2}. \quad (38)$$

b. Entanglement witnessed by mean values and variances of spin operators. For separable states, the inequalities

$$N(\Delta \hat{J}_{n_1})^2 \geq \langle \hat{J}_{n_2}^2 \rangle + \langle \hat{J}_{n_3}^2 \rangle, \quad (39a)$$

$$(\Delta \hat{J}_{n_1})^2 + (\Delta \hat{J}_{n_2})^2 + (\Delta \hat{J}_{n_3})^2 \geq \frac{N}{2}, \quad (39b)$$

$$(N-1)(\Delta \hat{J}_{n_1})^2 \geq \langle \hat{J}_{n_2}^2 \rangle + \langle \hat{J}_{n_3}^2 \rangle - \frac{N}{2}, \quad (39c)$$

$$(N-1)[(\Delta \hat{J}_{n_1})^2 + (\Delta \hat{J}_{n_2})^2] \geq \langle \hat{J}_{n_3}^2 \rangle + \frac{N(N-2)}{4} \quad (39d)$$

are all fulfilled, where n_1, n_2 and n_3 are three mutually orthogonal directions. The violation of at least one of the above inequalities signals that the state is entangled. Equation (39a) is equivalent to $\xi_R^2 \geq 1$, see Sec. II.C.5, and was introduced by Sørensen *et al.* (2001). The inequalities (39b)-(39d) have been introduced by Tóth *et al.* (2007, 2009). The third condition, Eq. (39c), can be rewritten as $\xi_D^2 \geq 1$, where

$$\xi_D^2 = \frac{N(\Delta \hat{J}_{n_1})^2}{\langle \hat{J}^2 \rangle - N/2 - \langle \hat{J}_{n_1} \rangle^2}. \quad (40)$$

In particular, the condition $\xi_D^2 < 1$ detects entanglement close to Dicke states (Tóth *et al.*, 2007), see also Raghavan *et al.* (2001). The detection of multiparticle entanglement close to Dicke states has been studied by Duan (2011) and Lücke *et al.* (2014); conditions for multipartite entanglement close to Dicke states of spin- J particles with $J > 1/2$ have been considered by Vitagliano *et al.* (2016). A violation of the condition (39b) can be used to detect entanglement in singlet states (Behbood *et al.*, 2014; Tóth and Mitchell, 2010). The inequalities (39b)-(39d) and the further inequality $\langle \hat{J}^2 \rangle \leq (N/2)(N/2 + 1)$, which is valid for all quantum states (not only for separable states), form a system of conditions that defines a polytope in the three dimensional space with coordinates $\langle \hat{J}_{n_1}^2 \rangle$, $\langle \hat{J}_{n_2}^2 \rangle$, and $\langle \hat{J}_{n_3}^2 \rangle$ (Tóth *et al.*, 2007, 2009). The polytope encloses all separable states. It has been demonstrated that Eqs. (39) form a complete set (Tóth *et al.*, 2007, 2009), meaning that it is not possible to add new entanglement conditions based on mean values and variances of spin moments that detect more entangled states. The inequalities (39) have been further generalized to arbitrary spin systems (Vitagliano *et al.*, 2014, 2011) and to systems of a fluctuating number of particles (Hyllus *et al.*, 2012b). Furthermore, Korbicz *et al.* (2005, 2006) have shown that, if the inequality

$$\left(\langle \hat{J}_{n_1}^2 \rangle + \frac{N(N-2)}{4} \right)^2 < \left(\langle \hat{J}_{n_2}^2 \rangle + \langle \hat{J}_{n_3}^2 \rangle - \frac{N}{2} \right)^2 + (N-1)^2 \langle \hat{J}_{n_1} \rangle^2 \quad (41)$$

holds, then the state possesses bipartite entanglement.

c. Spin squeezing and entanglement of symmetric states. We emphasize that none of the entanglement witnesses above require any assumptions on the symmetry of the state. For states that are symmetric under particle exchange, we have $\langle \hat{J}^2 \rangle = (N/2)(N/2 + 1)$. In this case, Eq. (41) can be rewritten as

$$\xi_N^2 = \frac{4(\Delta \hat{J}_n)^2}{N} < 1 - \frac{4\langle \hat{J}_n \rangle^2}{N^2}, \quad (42)$$

where ξ_N^2 is called the number-squeezing parameter. The inequality (42) is necessary and sufficient for bipartite

entanglement (Korbicz *et al.*, 2005, 2006). It should be noted that if $\xi_S^2 < 1$ holds, then the inequality (42) is satisfied as well (taking $n = \perp$ and $\langle \hat{J}_n \rangle = 0$). Hence, symmetric spin-squeezed states possess two-qubit entanglement. The converse is not true: since n in Eq. (42) is not necessarily orthogonal to the mean spin direction s , number squeezing ($\xi_N^2 < 1$) does not imply spin squeezing ($\xi_S^2 < 1$) or entanglement.

The relationship between Kitagawa-Ueda spin squeezing and bipartite entanglement has also been studied by Ulam-Orgikh and Kitagawa (2001) and Wang and Sanders (2003b). For an arbitrary symmetric state of N qubits, the spin-squeezing parameter (36) can be written in terms of the two-spin correlation function (Ulam-Orgikh and Kitagawa, 2001)

$$\xi_S^2 = 1 + (N-1)\langle \hat{\sigma}_\perp^{(i)} \otimes \hat{\sigma}_\perp^{(j)} \rangle, \quad i \neq j. \quad (43)$$

This equation shows that spin squeezing $\xi_S^2 < 1$ is equivalent to negative pairwise spin-spin correlations that, in turn, are sufficient for pairwise entanglement.¹² Furthermore, ξ_S^2 is related to the concurrence $\mathcal{C}$ (Ulam-Orgikh and Kitagawa, 2001; Wang and Sanders, 2003b):

$$\xi_S^2 = 1 - (N-1)\mathcal{C}. \quad (44)$$

We recall that $\mathcal{C} > 0$ is a necessary and sufficient condition of—and quantifies—entanglement of a pair of qubits (Hill and Wootters, 1997; Wootters, 1998). For symmetric pure states of two qubits, Eq. (44) always holds (Ulam-Orgikh and Kitagawa, 2001): these states are entangled if and only if they satisfy $\xi_S < 1$. For symmetric states of many qubits, Eq. (44) holds under certain conditions,¹³ see (Ma *et al.*, 2011; Wang and Sanders, 2003b), and in particular only if $\xi_S^2 \leq 1$. The relation (44) tells us that $\xi_S^2 < 1$ implies $\mathcal{C} > 0$ and thus bipartite entanglement. When $\xi_S^2 > 1$, Eq. (44) does not hold and we cannot draw any conclusion about bipartite entanglement: a state might have a positive concurrence and not be spin squeezed, as exemplified by Dicke states (Wang and Mølmer, 2002).

9. Einstein-Podolsky-Rosen entanglement and Bell correlations

a. Continuous variable and Einstein-Podolsky-Rosen entanglement Let us consider two bosonic modes, $|a_+\rangle$ and $|a_-\rangle$, and introduce the corresponding annihilation $\hat{a}_{\pm 1}$

and creation $\hat{a}_{\pm 1}^\dagger$ operators. Mode-separable quantum states are defined as $\hat{\rho}_{\text{msep}} = \sum_q p_q \hat{\rho}_q^{(+1)} \otimes \hat{\rho}_q^{(-1)}$, where $p_q > 0$, and $\hat{\rho}_q^{(\pm 1)}$ is the state in the $|a_{\pm 1}\rangle$ mode. Mode entanglement, *i.e.* $\hat{\rho} \neq \hat{\rho}_{\text{msep}}$ can be revealed by correlations between bosonic position $\hat{X}_{\pm 1}(\phi) = (\hat{a}_{\pm 1}e^{-i\phi} + \hat{a}_{\pm 1}^\dagger e^{i\phi})/\sqrt{2}$ and momentum $\hat{P}_{\pm 1}(\phi) = \hat{X}_{\pm 1}(\phi + \pi/2)$ quadratures (Reid *et al.*, 2009). Mode entanglement finds several applications in quantum technologies (Braunstein and van Loock, 2005). Mode-separable states fulfill (Duan *et al.*, 2000a; Simon, 2000)

$$V_{X(\phi)}^\pm + V_{P(\phi)}^\mp \geq 2, \quad (45)$$

where $V_{X(\phi)}^\pm = \text{Var}[\hat{X}_{+1}(\phi) \pm \hat{X}_{-1}(\phi)]$ and $V_{P(\phi)}^\mp = \text{Var}[\hat{P}_{+1}(\phi) \mp \hat{P}_{-1}(\phi)]$ are variances. A violation of this condition detects entanglement between the $|a_{\pm 1}\rangle$ modes. It is also a necessary and sufficient condition for mode entanglement in Gaussian states (Duan *et al.*, 2000a; Simon, 2000). For further (and sharper) conditions of mode entanglement see (Giovannetti *et al.*, 2003; Shchukin and Vogel, 2005; Walborn *et al.*, 2009).

Correlations between quadrature variances are at the heart of the Einstein-Podolsky-Rosen (EPR) paradox (Einstein *et al.*, 1935). When the quadratures $\hat{X}_{+1}$ and $\hat{P}_{+1}$ are measured in independent realizations of the same state, the correlations allow for a prediction of $\hat{X}_{-1}$ and $\hat{P}_{-1}$ with inferred variances violating the Heisenberg uncertainty relation $\Delta \hat{X}_{-1}^{\text{inf}} \Delta \hat{P}_{-1}^{\text{inf}} < 1/2$, known as EPR criterion (Reid, 1989; Reid *et al.*, 2009). This extends the original EPR discussion that was limited to perfect quadrature correlations. Non-steerable states, including separable states, fulfill (Reid, 1989)

$$V_{X(\phi)}^- \times V_{P(\phi)}^+ \geq 1/4. \quad (46)$$

The violation of this condition witnesses a strong form of entanglement (“EPR entanglement”) necessary to fulfill the EPR criterion. With atoms, continuous-variable entanglement has been first proved with room-temperature vapor cells (Julsgaard *et al.*, 2001). With spinor Bose-Einstein condensates, mode entanglement (Gross *et al.*, 2011) and EPR entanglement (Peise *et al.*, 2015a) have been demonstrated, see Sec. IV.C.

b. Bell correlations. The strongest form of correlations between particles are those that violate a Bell inequality (Bell, 1964). The existence of Bell correlations has profound implications for the foundations of physics and underpins quantum technologies such as quantum key distribution and certified randomness generation (Brunner *et al.*, 2014). Bell correlations have been observed in systems of at most a few (usually two) particles (Eibl *et al.*, 2003; Hensen *et al.*, 2015; Hofmann *et al.*, 2012; Lanyon *et al.*, 2014; Zhao *et al.*, 2003), but their role in many-body systems is largely unexplored.

¹² The two-qubit reduced density matrix of a separable symmetric state of N qubits has positive pairwise correlations: $\langle \hat{\sigma}_\perp^{(i)} \otimes \hat{\sigma}_\perp^{(j)} \rangle > 0$ for any $i \neq j$ (Wang and Sanders, 2003b).

¹³ Equation (44) has been derived in (Wang and Sanders, 2003b) for symmetric states having $\langle \hat{J}_s \rangle \neq 0$ and $\langle \hat{J}_{n_i} \rangle = \langle \hat{J}_s \hat{J}_{n_i} \rangle = \langle \hat{J}_{n_i} \hat{J}_s \rangle = 0$, $i = 1, 2$, where n_1 and n_2 are vectors orthogonal to the mean spin direction s .

While entanglement is necessary for Bell correlations, it is not sufficient (Brunner *et al.*, 2014; Werner, 1989). Because of this, the above entanglement criteria cannot be used to determine whether a system could violate a Bell inequality. In Schmied *et al.* (2016) a criterion in the spirit of a spin-squeezing parameter is derived to determine whether Bell correlations are present in an N -particle quantum system. For any two axes n_1 and n_2 , the inequality

$$\mathcal{W} = -\frac{|\langle \hat{J}_{n_2} \rangle|}{N/2} + (n_1 \cdot n_2)^2 \frac{\langle \hat{J}_{n_1}^2 \rangle}{N/4} + 1 - (n_1 \cdot n_2)^2 \geq 0 \quad (47)$$

is satisfied for all states that are not Bell correlated. States that satisfy $\mathcal{W} < 0$ can violate the many-particle Bell inequality of Tura *et al.* (2014). By optimizing Eq. (47) over the angle between the two axes, a criterion follows that facilitates comparison with well-known spin-squeezing criteria: for any two axes n_1 and n_2 perpendicular to each other,

$$\frac{\langle \hat{J}_{n_1}^2 \rangle}{N/4} \geq \frac{1}{2} \left[1 - \sqrt{1 - \left(\frac{\langle \hat{J}_{n_2} \rangle}{N/2} \right)^2} \right] \quad (48)$$

is satisfied for all states that are not Bell correlated. Detecting Bell correlations by violating inequality (47) or (48) requires only collective manipulations and measurements on the entire N -particle system. While this does not provide a loophole-free and device-independent Bell test, it is a powerful tool for characterizing correlations in many-body systems state-independently. Bell correlations according to Eqs. (47) and (48) have been observed with Bose-Einstein condensates, see Sec. III.B.3.

D. Tomography of spin states

1. Spin-noise tomography

Spin-noise tomography is a widely-used technique to gain information about spin-squeezed states, whose main characteristics are captured by their second spin moments along the squeezing and anti-squeezing directions. For this tomography, the state is rotated by an angle ϑ using resonant Rabi rotations, followed by projective spin measurements along the z -axis. The k th moment of these spin projections can be fit to a linear combination of $\cos(n\vartheta)$ and $\sin(n\vartheta)$ with $n \in \{k, k-2, k-4, \dots\}$, which allows interpolating these projective measurements to arbitrary angles ϑ . Also, technical noise sources can be characterized and their influence subtracted from the resulting moments; any spin squeezing concluded from these inferred moments will be called *inferred* spin squeezing. Since this method estimates spin projection moments separately, they do not necessarily fulfill all consistency criteria imposed by the positive-semidefiniteness of the density operator (Schmied, 2016).




FIG. 9 **Quasi-probability representations.** Upper row: P , W , and Q representations of a coherent state in the $+x$ direction, see Table I, with $s = \{1, 0, 0\}$. Lower row: P , W , and Q representations of a spin-squeezed state reached by one-axis twisting with $\chi t = 0.01\pi$ (see Fig. 14). The color scale for each panel is normalized to its maximum value. Here $N = 100$.

In practical situations concerning spin-squeezed states, however, this restriction is often irrelevant. The information captured from low-order spin moments through spin-noise tomography may be insufficient to characterize and visualize general spin states. Different techniques have been developed for full state tomography (Blume-Kohout, 2010; Paris and Řeháček, 2004; Schmied and Treutlein, 2011), estimating all spin moments up to order N .

2. Visualizing spin states

State representations on the Bloch sphere are very useful to gain an intuition about the properties of quantum states. In the following we consider symmetric states of N spin-1/2 particles (see Sec. II.A). There are various representations in the form of pseudo-probability densities (Schleich, 2001). These are based on the decomposition of a general density operator $\hat{\rho} = \sum_{k=0}^{2J} \sum_{q=-k}^k \rho_{kq} \hat{T}_{kq}^{(J)}$ into the orthonormalized spherical tensor (or multipole) operators $\hat{T}_{kq}^{(J)} = \sum_{\mu, \mu'=-J}^J (-1)^{J-\mu'} \langle J, \mu; J, -\mu' | k, q \rangle | J, \mu_z \rangle \langle J, \mu'_z |$ defined in terms of Clebsch-Gordan coefficients, with $\rho_{kq} = \text{Tr}[\hat{\rho}(\hat{T}_{kq}^{(J)})^\dagger]$ (Agarwal, 1981; Arecchi *et al.*, 1972; Dowling *et al.*, 1994; Schmied and Treutlein, 2011). This decomposition has properties similar to the Fourier transform in Euclidean space: it separates low-frequency components (small k) from high-frequency components (large k), which are affected very differently by experimental noise. Further, it allows us to define the family of spher-

TABLE I **Distributions of a coherent spin state.** Different representations of a coherent spin state with mean spin direction s ; the last column is the approximate expression for $N \gg 1$. Here $x = \{\sin \vartheta \cos \varphi, \sin \vartheta \sin \varphi, \cos \vartheta\} \cdot s$, $P_k(x)$ are Legendre polynomials, $J_1(x)$ is a Bessel function of the first kind, and $\zeta = \cos^{-1}(x)$.

exact	$N \gg 1$
$P(x) = \sum_{k=0}^N \frac{2k+1}{4\pi} P_k(x)$	$\approx \frac{(N+1)^2}{2\pi} \frac{J_1[(N+1)\zeta]}{(N+1)\zeta}$
$W(x) = \sum_{k=0}^N \frac{2k+1}{4\pi} \sqrt{\frac{N!(N+1)!}{(N-k)!(N+k+1)!}} P_k(x)$	$\approx \frac{N+\frac{3}{2}}{2\pi} \exp[-\frac{N+\frac{3}{2}}{2}\zeta^2]$
$Q(x) = \frac{N+1}{4\pi} \left(\frac{1+x}{2}\right)^N$	$\approx \frac{N+2}{4\pi} \exp[-\frac{N+2}{4}\zeta^2]$

ical functions (Agarwal, 1981)

$$f(\vartheta, \varphi) = \sqrt{\frac{N+1}{4\pi}} \sum_{k=0}^N f_k \sum_{q=-k}^k \rho_{kq} Y_{kq}(\vartheta, \varphi) \quad (49)$$

in terms of spherical harmonics $Y_{kq}(\vartheta, \varphi)$. All of these functions are linear in the density operator. The following representations (corresponding to different choices for the coefficients f_k) are often used, with examples given in Table I and Fig. 9.

a. Wigner distribution. The Wigner quasi-probability distribution $W(\vartheta, \varphi)$ (Wigner, 1932) corresponds to the case $f_k = 1$ in Eq. (49). It is found by replacing the spherical tensor operators in the decomposition of $\hat{\rho}$ by spherical harmonics of the same order, which obey the same ortho-normalization. Because of this close similarity, the Wigner quasi-probability distribution is equivalent to the density operator. $W(\vartheta, \varphi)$ is not a true probability density, as it can take negative values (Leibfried *et al.*, 1996; Lvovsky *et al.*, 2001; McConnell *et al.*, 2015). For continuous variables, this is often understood as a sign of nonclassical behavior. Note however that in the present finite-dimensional space, even the Wigner distribution of a coherent spin state has negative parts of amplitude $\sim 2^{-N}$, exponentially decreasing with the number of particles.

b. Husimi-Kano representation. The Husimi-Kano Q representation $Q(\vartheta, \varphi) = \frac{N+1}{4\pi} \langle \vartheta, \varphi, N | \hat{\rho} | \vartheta, \varphi, N \rangle$ (Barontini *et al.*, 2015; Haas *et al.*, 2014; Husimi, 1940) corresponds to the case $f_k = \sqrt{\frac{N!(N+1)!}{(N-k)!(N+k+1)!}}$ in Eq. (49). It is non-negative and proportional to the probability of finding the system in the coherent spin state $|\vartheta, \varphi, N\rangle$. Since $Q(\vartheta, \varphi)$ is the convolution of $W(\vartheta, \varphi)$ with the Wigner distribution of a coherent spin state (Agarwal, 1981), in practice the former contains much less information than the latter. Furthermore, recovering the Wigner distribution (and hence the density matrix) from an experimentally determined Q representation is generally not feasible.

c. Glauber-Sudarshan representation. The Glauber-Sudarshan P representation $P(\vartheta, \varphi)$ (Glauber, 1963; Kiesel *et al.*, 2008; Sudarshan, 1963) is obtained for $f_k = \sqrt{\frac{(N-k)!(N+k+1)!}{N!(N+1)!}}$ in Eq. (49). It is the deconvolution of $W(\vartheta, \varphi)$ with the Wigner distribution of a coherent spin state (Agarwal, 1981), and serves to construct the density operator from coherent spin states: $\hat{\rho} = \int_0^\pi d\vartheta \sin \vartheta \int_0^{2\pi} d\varphi P(\vartheta, \varphi) |\vartheta, \varphi, N\rangle \langle \vartheta, \varphi, N|$. While in infinite-dimensional Hilbert spaces the P representation is often of limited practical use because of its singular behavior (Scully and Zubairy, 1997), this is not the case for the representation of a spin. Indeed, in this case, partial-wave contributions are limited to angular momenta $k \leq N$ in Eq. (49) in order to match the number of degrees of freedom of the density operator. In contrast to continuous variables, here even classical (separable) states have P representations with negative parts, as can be seen in Table I.

III. ENTANGLEMENT VIA ATOMIC COLLISIONS: THE BOSONIC JOSEPHSON JUNCTION

Tunable elastic atom-atom collisions are naturally present in Bose-Einstein condensates and represent a well-established method to generate metrologically useful entanglement in these systems. Furthermore, Bose-Einstein condensates have a weak coupling to the environment and can be restricted to occupy two modes only. These can be two “internal” hyperfine atomic states or two “external” motional states of a trapping potential, see below, and are successfully described by the bosonic Josephson junction model¹⁴

$$\hat{H}_{\text{BJJ}} = -\hbar\Omega\hat{J}_x + \hbar\chi\hat{J}_z^2 + \hbar\delta\hat{J}_z. \quad (50)$$

Here, $\hat{J}_x$ describes linear coupling between the two modes, $\hat{J}_z^2$ accounts for the interaction of each atom with all the other particles in the system, and $\hat{J}_z$ models an energy imbalance between the modes. The interaction χ and the coupling Ω parameters (in the following we assume $\Omega \geq 0$, without loss of generality) can be precisely and independently tuned, and, in particular, switched on and off at will. Furthermore, Bose-Einstein condensates offer the possibility to control the trapping geometry and to count atoms directly using established techniques such as absorption or fluorescence imaging. Useful entanglement for quantum metrology can be found in the ground state of Eq. (50), for $\chi \neq 0$ (either positive and negative), see Sec. III.A. The ground state can be experimentally

¹⁴ The Hamiltonian (50) belongs to a class of models first introduced by Lipkin, Meshkov and Glick in nuclear physics (Glick *et al.*, 1965; Lipkin *et al.*, 1965; Meshkov *et al.*, 1965), see Ulyanov and Zaslavskii (1992) for a review.




FIG. 10 **Realizations of the bosonic Josephson junction with an atomic Bose-Einstein condensate.** (a) “External” bosonic Josephson junction realized by an ultracold gas trapped in a double-well potential (thick black line). The blue and red lines are the amplitude of the single-particle symmetric ground state and anti-symmetric first-excited state, vertically displaced at the corresponding eigenenergy (dashed line). (b) “Internal” bosonic Josephson junction made by a trapped Bose-Einstein condensate in two different hyperfine states. For each state, the thick black line is the harmonic trap and the thin colored line is the amplitude of the single particle ground state.

addressed by adiabatically tuning interaction and/or coupling parameters. The nonlinear interaction can also be exploited for the dynamical generation of entanglement starting from particles prepared in a separable state, see Secs. III.B and III.C. In particular, for $\Omega, \delta = 0$, Eq. (50) is equivalent to the one-axis twisting model first introduced by Kitagawa and Ueda (1993). A limitation is that the contact interaction, which is the ingredient to entangle the atoms, is also, via particle losses induced by non-elastic scattering, a main source of decoherence in these systems.

The *external bosonic Josephson junction* is made of two spatially separated modes. It can be realized with a dilute Bose-Einstein condensate confined in a double-well potential (Javanainen, 1986; Milburn *et al.*, 1997; Smerzi *et al.*, 1997), see Fig. 10(a). For a sufficiently-high barrier, the two lowest-energy single-particle states are almost degenerate and well separated from higher levels of the potential. Weak interparticle interactions do not significantly modify this situation and justify a two-mode approximation of the many-body wavefunction. The parameters in Eq. (50) are identified as (Ananikian and Bergeman, 2006; Milburn *et al.*, 1997; Spekkens and Sipe, 1999)

$$\hbar\Omega = E_e - E_g, \quad (51a)$$

$$\hbar\chi = \frac{8\pi\hbar^2 a_s}{M} \int d^3r |\psi_g(r)|^2 |\psi_e(r)|^2, \quad (51b)$$

where $\psi_g(r)$ and $\psi_e(r)$ are the spatial wavefunctions of the symmetric ground and antisymmetric first-excited states of the double-well potential with energies E_g and E_e , respectively, a_s is the s-wave scattering length (posi-

tive for repulsive interactions and negative for attractive interactions) (Landau and Lifchitz, 1975), and M the atomic mass. For weak interaction, the calculation of these coefficients can be improved using the first symmetric and antisymmetric eigenstates of the Gross-Pitaevskii equation in the double-well trap (Raghavan *et al.*, 1999; Zapata *et al.*, 1998). Experimentally, the external weak link has been realized on atom chips (Berrada *et al.*, 2013; Hall *et al.*, 2007; Jo *et al.*, 2007b; LeBlanc *et al.*, 2011; Maussang *et al.*, 2010; Schumm *et al.*, 2005), or optically (Albiez *et al.*, 2005; Gati *et al.*, 2006; Hadzibabic *et al.*, 2006; Levy *et al.*, 2007; Shin *et al.*, 2004). The spatial separation allows for devising sensors for a variety of fields and forces (Cronin *et al.*, 2009). The experimental challenges are the required high degree of stability of the potential difference and the relative phase between the two wells, as well as the precise control of the tunneling coupling between the two wells in order to induce Rabi rotations.

The *internal bosonic Josephson junction* is created with a trapped Bose-Einstein condensate in two different hyperfine states (Cirac *et al.*, 1998; Steel and Collett, 1998), see Fig. 10(b). The Josephson-like coupling is provided by an electromagnetic field that coherently transfers particles between the two states via Rabi rotations (Böhi *et al.*, 2009; Hall *et al.*, 1998a,b). Assuming that the external motion of the atoms is not influenced by the internal dynamics, one can apply a single-mode approximation for each atomic species. The many-body Hamiltonian becomes Eq. (50) with coefficients

$$\hbar\Omega = \hbar\Omega_R \int d^3r \psi_a^*(r) \psi_b(r), \quad (52a)$$

$$\hbar\chi = U_{aa} + U_{bb} - 2U_{ab}, \quad (52b)$$

$$U_{ij} = \frac{2\pi\hbar^2 a_s^{(i,j)}}{M} \int d^3r |\psi_i(r)|^2 |\psi_j(r)|^2 \quad (52c)$$

where Ω_R is the Rabi frequency, $a_s^{(a,a)}$, $a_s^{(b,b)}$ and $a_s^{(a,b)}$ are the intra- and inter-species s-wave scattering lengths, and $\psi_{a,b}(r)$ are single-particle mode functions of the two internal states, which can be determined in a mean-field description from the Gross-Pitaevskii equation. A more accurate value for χ is obtained if one also takes into account the change of the mode functions with particle number as in Li *et al.* (2008b). Since the phase and amplitude of the coupling field Ω_R can be switched on and off on nanosecond time scales, arbitrary rotations on the Bloch sphere can be implemented that are helpful to read out and characterize the internal state. Furthermore, during Rabi coupling pulses it is possible to reach the regime $\Omega \gg N\chi$, where interaction effects can be neglected.

A. Metrologically useful entanglement in the ground state of the bosonic Josephson junction

Several approaches to the bosonic Josephson junction model Hamiltonian (50) have been discussed in the literature. A semiclassical (mean-field) approximation provides useful insights (Raghavan *et al.*, 1999; Smerzi *et al.*, 1997). It is obtained by replacing mode operators with complex numbers: $\hat{a} \mapsto \sqrt{N_a}e^{-i\varphi_a}$ and $\hat{b} \mapsto \sqrt{N_b}e^{-i\varphi_b}$, where $N_{a,b}$ and $\varphi_{a,b}$ are the numbers of particles and phases of the condensate in the $|a\rangle$ and $|b\rangle$ modes, respectively. The spin operators are replaced by $\hat{J}_x \rightarrow \frac{N}{2}\sqrt{1-z^2}\cos\varphi$, $\hat{J}_y \rightarrow \frac{N}{2}\sqrt{1-z^2}\sin\varphi$, and $\hat{J}_z \rightarrow \frac{N}{2}z$, where $\varphi = \varphi_a - \varphi_b$ is the relative phase between the two condensate modes, and $z = (N_a - N_b)/N$ the fractional population difference ($-1 \leq z \leq 1$). The Hamiltonian (50) becomes

$$H(z, \varphi) = \frac{\Lambda z^2}{2} - \sqrt{1-z^2}\cos\varphi + \Xi z, \quad (53)$$

where energies are in units of $N\hbar\Omega/2$, $\Lambda = N\chi/\Omega$, and $\Xi = \delta/\Omega$. In the following we mainly focus on the case $\Xi = 0$, unless explicitly stated. For $\Lambda \geq 1$, the minimum of $H(z, \varphi)$ is found at the point $\Upsilon_0 = (z = 0, \varphi = 0)$.

A common method to extend the analysis beyond the mean-field approximation consists of quantizing the conjugate number and phase semiclassical variables (Leggett and Sols, 1991), *i.e.* replacing z and φ with operators $\hat{z}$ and $\hat{\varphi}$ obeying the commutation relation $[\hat{z}, \hat{\varphi}] = 2i/N$, where $2/N$ plays the role of a Planck constant. We can thus write $\hat{\varphi} = -\frac{2i}{N}\hat{\partial}_z$ and $\hat{z} = \frac{2i}{N}\hat{\partial}_\varphi$ by canonical quantization. Note that a rigorous phase operator is lacking (Carruthers and Nieto, 1968; Lynch, 1995), and that the above phase quantization may bring notable artifacts when predicting large phase fluctuations. Series-expanding the Hamiltonian (53) to second order around Υ_0 , $H(z, \varphi) = \frac{\Lambda+1}{2}z^2 + \frac{\varphi^2}{2}$, we obtain the quantum Hamiltonian

$$\hat{H}_z = -\frac{2}{N^2}\hat{\partial}_z^2 + \frac{\Lambda+1}{2}\hat{z}^2 \quad (54)$$

in number representation, and

$$\hat{H}_\varphi = -\frac{2(\Lambda+1)}{N^2}\hat{\partial}_\varphi^2 + \frac{1}{2}\hat{\varphi}^2 \quad (55)$$

in phase representation. Equations (54) and (55) predict Gaussian number and phase ground state wave functions with variances

$$\sigma_z^2 = \frac{1}{N\sqrt{1+\Lambda}} \text{ and } \sigma_\varphi^2 = \frac{\sqrt{1+\Lambda}}{N}, \quad (56)$$

respectively (Paraoanu *et al.*, 2001; Smerzi and Raghavan, 2000). Improvements over Eqs. (54) and (55) have been discussed in the literature. Considering again a second-order expansion of Eq. (53) around $\varphi = 0$ and

rewriting $\frac{1}{2}\sqrt{1-z^2}\varphi^2 = \varphi\frac{1}{2}\sqrt{1-z^2}\varphi$, one obtains the Hermitian Hamiltonian

$$\hat{H}_z = -\frac{2}{N^2}\hat{\partial}_z\sqrt{1-\hat{z}^2}\hat{\partial}_z + W_0(\hat{z}), \quad (57)$$

describing a fictitious quantum particle with z -dependent effective mass moving in an anharmonic one-dimensional potential

$$W_0(z) = \frac{\Lambda z^2}{2} - \sqrt{1-z^2} + \Xi z. \quad (58)$$

Shchesnovich and Trippenbach (2008) have derived the above Hamiltonian using a continuous approximation of the relative population difference, and avoiding phase-number commutation relations (see also Javanainen and Ivanov (1999) and Spekkens and Sipe (1999)). For $\Xi = 0$, the effective potential (58) changes from harmonic for $\Lambda > -1$, to quartic at $\Lambda = -1$, to a double-well shape with wells centered at $z_\pm = \pm\sqrt{1-1/\Lambda^2}$ for $\Lambda < -1$. The fictitious quantum particle description is expected to give accurate results provided that the wavefunction is sufficiently smooth and vanishes at the boundaries of z (Juliá-Díaz *et al.*, 2012), and phase fluctuations remain small. Anglin *et al.* (2001) proposed a complementary approach consisting of a projection of the bosonic Josephson junction Hamiltonian (50) over the overcomplete Bargmann basis. This leads to an exact quantum phase model $\hat{H}_{\text{BJJ}}|\psi\rangle = \int_{-\pi}^{\pi} \frac{d\varphi}{2\pi} |\varphi\rangle e^{(N/2\Lambda)\cos\varphi} \hat{H}_{\text{EQPM}}\psi(\varphi)$, where $|\varphi\rangle$ are Bargmann phase states, $|\psi\rangle$ is an arbitrary two-mode state, $\psi(\varphi)$ is an effective phase wavefunction, and

$$\hat{H}_{\text{EQPM}} = -\frac{2\Lambda}{N^2}\partial_\varphi^2 - \left(1 + \frac{1}{N}\right)\cos(\varphi) - \frac{1}{4\Lambda}\cos(2\varphi). \quad (59)$$

This equation is exact in the sense that the lowest $N+1$ frequencies in its eigenspectrum are exactly the spectrum of Eq. (50) (Anglin *et al.*, 2001).

1. Ground state for positive nonlinearities

A positive effective nonlinearity ($\Lambda > 0$) makes number fluctuations between the two modes energetically unfavorable. The ground state $|\psi_\Lambda\rangle$ of Eq. (50) with $\delta = 0$ is thus characterized by a squeezed atom-number distribution at the expense of increased phase fluctuations (Javanainen and Ivanov, 1999; Leggett and Sols, 1998; Spekkens and Sipe, 1999; Steel and Collett, 1998). Following Leggett (2001), we distinguish three regimes, see Fig. 11.

a. Rabi regime, $0 < \Lambda \ll 1$. At $\Lambda = 0$ the ground state is given by the coherent spin state pointing in the positive- x direction (for $\Omega > 0$), $|\psi_{\Lambda=0}\rangle \propto (\hat{a}^\dagger + \hat{b}^\dagger)^N|\text{vac}\rangle$,




FIG. 11 **Entanglement in the ground states of the BJJ.** Upper panels: normalized Wigner distribution of the ground state $|\psi_\Lambda\rangle$ of the Hamiltonian (50), in different regimes and for $\delta = 0$. The histograms are probability distributions $P_y(m) = |\langle m_y | \psi_\Lambda \rangle|^2$ and $P_z(m) = |\langle m_z | \psi_\Lambda \rangle|^2$. Crossing from the Rabi to the Fock regime (for positive nonlinearities), P_z narrows while P_y broadens. In the Fock regime, P_y vanishes for odd values of m . In the transition from the disordered to the ordered phase (for negative nonlinearities), P_y narrows while P_z broadens and splits when crossing the critical value $\Lambda = -1$. (Lower panels) Normalized quantum Fisher information (F_Q/N , red line) and inverse spin-squeezing parameter ($1/\xi_R^2$, blue line) as a function of Λ . For $\Lambda > 0$, the solid black line is Eq. (60). For $\Lambda < 0$, the solid black lines are Eq. (64) in the disordered regime, and Eq. (65) in the ordered regime. Here, $N = 100$, and the color scale of the Wigner function is as in Fig. (5).

with binomial occupation of each mode. In the Rabi regime, the Josephson junction is dominated by tunneling, which keeps a well-defined relative phase between the two modes. The coherence is high, $\langle \hat{J}_x \rangle \approx N/2$, and number (phase) fluctuations slightly decrease (increase) with respect to the noninteracting case. Using $(\Delta \hat{J}_y)^2 = N^2 \sigma_\varphi^2/4$, $(\Delta \hat{J}_z)^2 = N^2 \sigma_z^2/4$, and Eq. (56), we find

$$\xi_R^2 = \frac{1}{\sqrt{1+\Lambda}} \text{ and } \frac{F_Q[|\psi_\Lambda\rangle, \hat{J}_y]}{N} = \sqrt{\Lambda+1}. \quad (60)$$

In particular, $F_Q/N = 1/\xi_R^2$ in this regime.¹⁵ The ground state of Eq. (50) is spin-squeezed along the z axis, and useful for sensing rotations around the y axis.

b. Josephson regime, $1 \ll \Lambda \ll N^2$. In the Josephson regime number (phase) fluctuations are further reduced (increased) while the coherence remains high. The approximations leading to Eq. (56) are still very good. We

thus expect Gaussian number and phase distributions with width $\sigma_z^2 = 1/(N\sqrt{\Lambda})$ and $\sigma_\varphi^2 = \sqrt{\Lambda}/N$, respectively, giving (Pezzè *et al.*, 2005)

$$\xi_R^2 = \frac{1}{\sqrt{\Lambda}} \text{ and } \frac{F_Q[|\psi_\Lambda\rangle, \hat{J}_y]}{N} = \sqrt{\Lambda}. \quad (61)$$

For a fixed ratio $\chi/\Omega = \Lambda/N$, we have $F_Q/N = 1/\xi_R^2 = \sqrt{\chi/\Omega\sqrt{N}}$, predicting a scaling of phase sensitivity $(\Delta\theta_{\text{QCR}})^2 = \mathcal{O}(N^{-3/2})$ intermediate between the standard quantum limit and the Heisenberg limit.

c. Fock regime, $\Lambda \gg N^2$. In the Fock regime, interaction dominates over tunneling, giving rise to a fragmented Bose-Einstein condensate (Spekkens and Sipe, 1999). The phase fluctuates so strongly that it becomes ill defined within a 2π period, and the approach leading to Eq. (56) breaks down. The ground state is approximatively obtained by putting an equal number of particles in both modes, with vanishing number fluctuations. For $\Lambda \rightarrow \infty$ and even values of N , we have $|\psi_\Lambda\rangle \approx (\hat{a}^\dagger)^N (\hat{b}^\dagger)^N |\text{vac}\rangle = |N/2\rangle_a |N/2\rangle_b$ (Dunningham *et al.*, 2002). In the limit of large but finite Λ , we find

$$\xi_R^2 = \frac{2}{N+2} \text{ and } \frac{F_Q[|\psi_\Lambda\rangle, \hat{J}_\perp]}{N} = \frac{N+2}{2}, \quad (62)$$

¹⁵ In Eqs. (60)-(63), *i.e.* for positive nonlinearities, the spin squeezing is calculated as $\xi_R^2 = N(\Delta \hat{J}_z)^2 / \langle \hat{J}_x \rangle^2$. For negative nonlinearities, Eq. (64), as $\xi_R^2 = N(\Delta \hat{J}_y)^2 / \langle \hat{J}_x \rangle^2$.

where $\perp$ is any axis on the equatorial plane of the Bloch sphere. In this limit, both $(\Delta\hat{J}_z)^2$ and $\langle\hat{J}_x\rangle$ vanish, such the ratio $\xi_R^2 = N(\Delta\hat{J}_z)^2/\langle\hat{J}_x\rangle$ is finite but very sensitive to experimental noise. For odd values of N , we have $|\psi_\Lambda\rangle \approx \frac{1}{\sqrt{2}}(|\frac{N+1}{2}\rangle_a|\frac{N-1}{2}\rangle_b + |\frac{N-1}{2}\rangle_a|\frac{N+1}{2}\rangle_b)$ (Jääskeläinen *et al.*, 2004), which is spin-squeezed with

$$\xi_R^2 = \frac{4N}{N(N+2)+1}, \text{ and } \frac{F_Q[|\psi_\Lambda\rangle, \hat{J}_\perp]}{N} = \frac{(N^2-1)}{2N} + 1. \quad (63)$$

2. Ground state for negative nonlinearities

A negative effective nonlinearity ($\Lambda < 0$) favors localization of particles in one mode and, in a symmetric Josephson junction ($\delta = 0$), enhances number fluctuations.¹⁶ The interplay of linear coupling and nonlinear interaction in the Hamiltonian (50) gives rise to a second-order quantum phase transition with discrete symmetry breaking and mean-field critical exponents. This quantum phase transition is due to the competition between interaction and coupling (Botet *et al.*, 1982; Gilmore and Feng, 1978; Ulyanov and Zaslavskii, 1992), and occurs in the thermodynamic limit, $N \rightarrow \infty$ and $\chi \rightarrow 0$ such that $\Lambda = -1$. The order parameter is given by the absolute value of the population imbalance, $|z|$, with $W_0(z)$ in Eq. (58) playing the role of an effective Ginzburg-Landau potential. Mode- and particle-entanglement (Buonsante *et al.*, 2012; Dusuel and Vidal, 2005; Ma and Wang, 2009; Orús *et al.*, 2008) and enhanced fluctuations of the order parameter (Ziń *et al.*, 2008) approaching the transition point have been extensively studied. This quantum phase transition has been experimentally investigated by Trenkwalder *et al.* (2016) in a double-well potential. The dynamics following the sudden quench into the vicinity of the quantum critical point (in a two-component Bose gas) has been studied by Nicklas *et al.* (2015).

a. Disordered phase, $-1 < \Lambda < 0$. The approximations leading to Eq. (56) remain valid also for weak attractive interactions. We thus find phase squeezing and number anti-squeezing (Steel and Collett, 1998):

$$\xi_R^2 = \sqrt{1+\Lambda}, \text{ and } \frac{F_Q[|\psi_\Lambda\rangle, \hat{J}_z]}{N} = \frac{1}{\sqrt{1+\Lambda}}. \quad (64)$$

The ground state is entangled and useful for sensing rotations around the z -axis, see Fig. 11. At $\Lambda \rightarrow -1$ the harmonic-oscillator approximation breaks down and Eq. (56) predicts a divergence of number fluctuations. At the transition point one finds $F_Q/N \approx 1/\xi_R^2 = \mathcal{O}(N^{-1/3})$.

b. Ordered phase, $\Lambda < -1$. This phase is characterized by the presence of macroscopic-superposition state (Cirac *et al.*, 1998; Ho and Ciobanu, 2004; Huang and Moore, 2006). The mean spin length $|\langle\hat{J}\rangle|$ vanishes and spin squeezing is lost, see Fig. 11. For strong nonlinearities, $\Lambda \ll -1$, the ground state is given by the symmetric superposition (*i.e.* a Schrödinger cat-like state) of states localized on the right and the left wells of the effective double-well Ginzburg-Landau potential $W_0(z)$ for $\Xi = 0$ in Eq. (58) (Shchesnovich and Trippenbach, 2008), giving

$$\frac{F_Q[|\psi_\Lambda\rangle, \hat{J}_z]}{N} = N\left(1 - \frac{1}{\Lambda^2}\right). \quad (65)$$

For $\Lambda \ll -\sqrt{N}$, the ground state is approximatively given by a NOON state and the quantum Fisher information tends toward the ultimate bound N^2 . The macroscopic superposition is lost for infinitesimally small energy imbalances $\Xi \neq 0$, breaking the symmetry of $W_0(z)$.

3. Adiabatic splitting

Experimentally, the Rabi and Josephson regimes for repulsive interactions and the disordered phase for attractive interactions can be reached by the adiabatic splitting of a Bose-Einstein condensate initially prepared in the strong-tunneling (Rabi) regime (Bodet *et al.*, 2010; Isella and Ruostekoski, 2005; Javanainen and Ivanov, 1999; Menotti *et al.*, 2001; Pezzè *et al.*, 2005; Streltsov *et al.*, 2007). By changing the plasma frequency $\omega_p = \Omega\sqrt{1+\Lambda}$ adiabatically in time, the system follows its ground state if (Schaff *et al.*, 2014)

$$\left|\frac{d\omega_p}{dt}\right| \lesssim \omega_p^2. \quad (66)$$

Approaching the Fock regime and the ordered phase, the required timescale for adiabaticity eventually diverges, setting a limit to the attainable entanglement. Shortcuts to adiabaticity (Juliá-Díaz *et al.*, 2012; Lapert *et al.*, 2012; Yuste *et al.*, 2013) and optimal control techniques (Grond *et al.*, 2009a,b; Huang and Moore, 2008; Pichler *et al.*, 2016) for the fast production of highly-entangled states have been also studied. It should be noted that Eq. (66) assumes that the gas is prepared in the ground states configuration. Experimentally, finite temperature of the gas is the main factor limiting the squeezing via the adiabatic splitting technique (Estève *et al.*, 2008).

4. Experimental spin squeezing in the ground state of the BJJ

Estève *et al.* (2008) reported the first direct experimental demonstration of spin squeezing and entanglement in the ground state of the bosonic Josephson junction. A ^{87}Rb Bose-Einstein condensate confined in a shallow

¹⁶ For attractive Bose gases we consider here a density range in which the condensate remains stable.




FIG. 12 **Spin squeezing in the ground states of the bosonic Josephson junction.** Symbols report number squeezing (with photon shot-noise subtracted) and phase coherence obtained by splitting a Bose-Einstein condensate. Open and filled symbols correspond to different barrier heights (larger for the open symbols). The shaded areas show systematic error bounds. Solid lines are reference values for ξ_R^2 , the orange line being the standard quantum limit. Measurements are shown for the two main well pairs of a six-well lattice (red and blue circles) and for a double-well potential (green diamonds). The total atom number in each pair is approximately $N = 2200$ in the six-well case and $N = 1600$ in the double-well case. The inset shows single-site-resolving absorption images of atoms trapped in an optical lattice superposed on an atomic dipole trap. Taken from (Estève *et al.*, 2008).

harmonic trap is split adiabatically by ramping up a one-dimensional optical lattice. The particles are distributed over two and six lattice sites, see inset of Fig. 12. Direct atom detection better than atomic shot noise is implemented by absorption imaging of the atomic density, with a spatial resolution well below the lattice spacing. This gives direct access to the number-squeezing parameter $\xi_N^2 = 4(\Delta\hat{J}_z)^2/N$. Additionally, the phase coherence $\langle\cos\varphi\rangle^2$ is directly obtained from the interference pattern observed after releasing the atomic cloud from the trap. Noticing that $\langle\hat{J}_x\rangle^2 = \langle\hat{N}_a\rangle\langle\hat{N}_b\rangle\langle\cos\varphi\rangle^2$ (valid for $\langle\hat{N}_a\rangle, \langle\hat{N}_b\rangle \approx N/2 \gg 1$ and $\sigma_\varphi^2 \ll 2\pi$), metrological spin squeezing is given by $\xi_R^2 = \xi_N^2/\langle\cos\varphi\rangle^2$. The simultaneous presence of number squeezing and high phase coherence allowed to reach an inferred spin squeezing of $\xi_R^2 = -3.8$ dB for the two main well pairs of a six-well lattice, and $\xi_R^2 = -2.3$ dB for the double-well configuration. The results are summarized in Fig. 12.




FIG. 13 **Number and phase distributions of a split Bose-Einstein condensate.** Number (a) and phase (b) distributions after adiabatic splitting a Bose-Einstein condensate in a double-well potential. (a) Distribution of photon imbalance $s = s_L - s_R$ between fluorescence images of the left and right cloud (inset), proportional to the imbalance of atom numbers in the two wells. The solid black line is the normal distribution corresponding to the measured number-squeezing factor, the dotted red line is the expected distribution when detection noise is subtracted, and the dashed blue line is the distribution expected for a coherent spin state in the absence of detection noise. (b) The curves indicate a normal distribution with the measured $\Delta\phi$ (solid black) and the distributions expected for a coherent state in the absence (dashed blue) and in the presence (dashed-dot green) of detection noise. The inset shows a typical matter-wave interference pattern from which the phase is extracted. Taken from (Berrada *et al.*, 2013).

More recent experiments have studied number squeezing (Maussang *et al.*, 2010) and spin squeezing (Berrada *et al.*, 2013) when splitting a double well trap on an atom chip. Maussang *et al.* (2010) report a detailed investigation of the optimal splitting time of a Bose-Einstein condensate, given by the interplay of the barrier raising rate (which must be slow in order to avoid excitations to higher modes), heating effects and atom loss. Berrada *et al.* (2013) studied the coherent splitting of a Bose-Einstein condensate of $N = 1200$ ^{87}Rb atoms in a double well on an atom chip. The splitting is based on radio-frequency dressing (Hofferberth *et al.*, 2006) that dynamically deforms the static magnetic trap into a double-well potential (Schumm *et al.*, 2005). The quantum state gen-




FIG. 14 **One-axis twisting dynamics.** The left panel reports the inverse spin-squeezing parameter ($1/\xi_R^2$, blue line) and normalized QFI (F_Q/N , red line) as function of $\chi t/\pi$. The right panels show snapshots of the Wigner distribution at different times. For $\chi t = \pi/2$ we plot both the Wigner distributions for $N = 100$ (left) and $N = 101$ (right). Here $N = 100$ (unless indicated), and the color scale is as in Fig. 5.

erated after the splitting is characterized via the measurement of the number of particles in each well and the relative phase, see Fig. 13, showing number squeezing and phase anti-squeezing. From these results, using $\xi_R^2 = \xi_N^2 / \langle \cos \varphi \rangle^2$, it is possible to obtain $\xi_R^2 = -5.6$ dB ($\xi_R^2 = -7.8$ dB inferred). This corresponds to a useful (for metrology) entanglement depth of 150 atoms, excluding useful entanglement of less than 67 particles with 90% probability. The experiment has also investigated the reduced phase diffusion associated to number squeezing (Castin and Dalibard, 1997; Javanainen and Wilkens, 1997; Lewenstein and You, 1996), and fast recombination (Berrada *et al.*, 2016; Jo *et al.*, 2007a; Negretti and Henkel, 2004; Negretti *et al.*, 2008; Scott *et al.*, 2008) of the condensates in the two wells.

B. One-axis twisting dynamics

One-axis twisting is a benchmark model for studying the generation of spin-squeezed states (Kitagawa and Ueda, 1993), Schrödinger cats (Mølmer and Sørensen, 1999) and useful entanglement for quantum metrology (Pezzè and Smerzi, 2009). The one-axis twisting Hamiltonian (here, along the z -axis) is

$$\hat{H}_{\text{OAT}} = \hbar \chi \hat{J}_z^2. \quad (67)$$

This model can be realized via atom-atom elastic collisions in a Bose-Einstein condensate (Sørensen *et al.*, 2001), see Eq. (50), trapped ions, see Sec. VI.B, and cold atoms in an optical cavity, see Sec. V.B.

The quantum dynamics $e^{-i\hat{H}_{\text{OAT}}t/\hbar}$ of a localized spin wavepacket can be roughly viewed as a rotation $e^{-i\chi t \langle \hat{J}_z \rangle \hat{J}_z}$ around the z axis. The $\langle \hat{J}_z \rangle$ -dependent angular velocity (whose sign differs on the two hemispheres of the Bloch sphere) leads to a twisting of the state on the Bloch sphere. To see this effect quantitatively,

let us consider the dynamical evolution of an initial coherent spin state pointing along the positive x -axis, $|\psi_{\text{OAT}}(t)\rangle = e^{-i\chi t \hat{J}_z^2} |\pi/2, 0, N\rangle$, see Fig. 14. The coherent spin state initially stretches in the $y-z$ plane tangential to the Bloch sphere. Kitagawa and Ueda (1993) identified the squeezing angle $\delta = \frac{1}{2} \arctan(\frac{B}{A})$ in terms of $A = 1 - \cos^{N-2}(2\chi t)$ and $B = 4 \sin(\chi t) \cos^{N-2}(\chi t)$, as well as the squeezed and anti-squeezed spin components $\hat{J}_s = \hat{J}_z \cos(\delta) - \hat{J}_y \sin(\delta)$ and $\hat{J}_{\text{as}} = \hat{J}_y \cos(\delta) + \hat{J}_z \sin(\delta)$, respectively. Squeezing is accompanied by loss of contrast, $\langle \hat{J}_x \rangle = \frac{N}{2} \cos^{N-1}(\chi t)$, as the state spreads on the Bloch sphere. The spin squeezing $\xi_R^2 = N(\Delta \hat{J}_s)^2 / \langle \hat{J}_x \rangle^2$ and the quantum Fisher information $F_Q[|\psi_{\text{OAT}}(t)\rangle, \hat{J}_{\text{as}}] = 4(\Delta \hat{J}_{\text{as}})^2$ are readily calculated, giving (Kitagawa and Ueda, 1993; Sørensen *et al.*, 2001)

$$\xi_R^2 = \frac{4 + (N-1)(A - \sqrt{A^2 + B^2})}{4 \cos^{2N-2}(\chi t)}, \quad (68)$$

and (Pezzè and Smerzi, 2009)

$$\frac{F_Q[|\psi_{\text{OAT}}(t)\rangle, \hat{J}_{\text{as}}]}{N} = 1 + \frac{(N-1)}{4} (A + \sqrt{A^2 + B^2}), \quad (69)$$

respectively.¹⁷ For $\chi t \lesssim 1/\sqrt{N}$ the state is spin squeezed, $\xi_R^2 < 1$, reaching $\xi_R^2 = \mathcal{O}(N^{-2/3})$ at an optimal time $\chi t = \mathcal{O}(N^{-2/3})$. For $\chi t \gtrsim 1/\sqrt{N}$ the states wraps around the Bloch sphere and spin squeezing is lost, $\xi_R^2 > 1$. Yet, the state is still entangled. The quantum Fisher information reaches a plateau $F_Q[|\psi_{\text{OAT}}(t)\rangle, \hat{J}_{\text{as}}] = N(N+1)/2$ for $2/\sqrt{N} \lesssim \chi t \lesssim \pi/2 - 2/\sqrt{N}$ signaling entanglement,

¹⁷ The quantum Fisher information optimized over rotation directions in the Bloch sphere is given by the maximum between $F_Q[|\psi_{\text{OAT}}(t)\rangle, \hat{J}_{\text{as}}]/N$, Eq. (69), and $F_Q[|\psi_{\text{OAT}}(t)\rangle, \hat{J}_x]/N = N[1 - \cos^{2N-2}(\chi t)] - (N-1)A/2$.

according to Eq. (24), in the over-squeezed state. The state evolves, at time $\chi t = \pi/n$, into a coherent superposition of $2 \leq n \lesssim \pi\sqrt{N}/2$ coherent spin states distributed evenly on the equator of the Bloch sphere (Agarwal *et al.*, 1997), see Fig. 14. At $\chi t = \pi/2$ we observe the dynamical creation of a NOON state along the x axis, $|\psi_{\text{OAT}}(\pi/2\chi)\rangle = e^{-i\frac{\pi}{2}\hat{J}_y}|\text{NOON}\rangle$, if N is even, and along the y axis, $|\psi_{\text{OAT}}(\pi/2\chi)\rangle = e^{-i\frac{\pi}{2}\hat{J}_x}|\text{NOON}\rangle$, if N is odd (Agarwal *et al.*, 1997; Mølmer and Sørensen, 1999). For the NOON state, the quantum Fisher information, optimized over the spin direction, reaches its maximum value $F_Q = N^2$ [see Sec. II.C.7 for a discussion of the properties of the $|\text{NOON}\rangle$ state]. For even (odd) values of N , the dynamics is reversed for $\chi t \geq \pi/2$ ($\chi t \geq \pi$) and we have a complete revival of the initial condition at $\chi t = \pi$ ($\chi t = 2\pi$). The one-axis twisting dynamics is modified by particle loss (Li *et al.*, 2008a; Sinatra *et al.*, 2013, 2012a, 2011; Spehner *et al.*, 2014) and other imperfections such as phase noise (Ferrini *et al.*, 2008, 2011) and finite temperature of the Bose gas (Sinatra *et al.*, 2011), limiting the achievable squeezing. We will further comment on these works in Sec. VII.

1. Spin squeezing and particle entanglement

Spin squeezing obtained via one-axis twisting dynamics in a Bose-Einstein condensate has been first experimentally demonstrated by Gross *et al.* (2010) and Riedel *et al.* (2010). These experiments used a ^{87}Rb condensate prepared in a coherent spin state with equal mean populations in two hyperfine states. With ^{87}Rb atoms, the interaction parameter $\hbar\chi = U_{aa} + U_{bb} - 2U_{ab}$, see Eq. 52b, is small due to an almost perfect compensation of inter- and intra-species collisional interactions, *i.e.* $2U_{ab} \approx U_{aa} + U_{bb}$. Two different approaches have been used to increase χ artificially: Riedel *et al.* (2010) used a state-dependent trapping potential to change the wave function overlap between the two internal states and thus reduce U_{ab} (Böhi *et al.*, 2009; Li *et al.*, 2008b), while Gross *et al.* (2010) exploited a Feshbach resonance (Chin *et al.*, 2010) to change the scattering length $a_s^{(a,b)}$.

Riedel *et al.* (2010) used a spatially-inhomogeneous microwave field to manipulate the trapping potentials of the atom cloud in the two hyperfine states $|a\rangle = |F=1, m_F=-1\rangle$ and $|b\rangle = |2, 1\rangle$ by microwave level shifts (Treutlein *et al.*, 2006). The trapping potential for atoms in one state is suddenly shifted by such a state-selective force, and the states coherently demix and begin to oscillate in space. In this way the overlap of the wavefunctions of the two states changes dynamically and modulates collisional effects (Li *et al.*, 2008b): according to Eq. (52c), when the two components are spatially separated, the inter-species interaction U_{ab} vanishes. The parameter χ is then determined solely by intra-species inter-




FIG. 15 **Spin-noise tomography and reconstructed Wigner function of a spin-squeezed Bose-Einstein condensate.** a) Reconstructed Wigner distribution of a spin-squeezed state of $N \approx 1250$ atoms (see Sec. II.D.2). The black contour line indicates where the Wigner function has fallen to $1/e$ of its maximum. For comparison, the circular $1/e$ contour of an ideal coherent spin state is shown. b) Observed spin fluctuations of a spin-squeezed state (solid circles) and of a coherent spin state (open circles), as a function of the turning angle of the Bloch sphere (see Sec. II.D.1). Solid lines are results of dynamical simulations including losses and technical noise. Taken from Riedel *et al.* (2010).

actions and reaches sufficiently-high values to induce fast spin-squeezing dynamics. After each full oscillation, the two states coherently remix and the collisional squeezing dynamics stops. Spin-noise tomography and the reconstructed Wigner function of the squeezed state are shown in Fig. 15. The results demonstrate a squeezing parameter $\xi_R^2 = -1.2 \text{ dB}$ ($\xi_R^2 = -2.5 \text{ dB}$ inferred). More recent experiments using this technique achieved up to $\xi_R^2 = -7.0 \text{ dB}$ with detection noise subtracted (Ockeloen *et al.*, 2013; Schmied *et al.*, 2016).

Gross *et al.* (2010) used a Bose-Einstein condensate of $N \approx 400$ atoms in the $|a\rangle = |1, 1\rangle$ and $|b\rangle = |2, -1\rangle$ hyperfine levels of the electronic ground state of Rubidium. A bias magnetic field is used to bring the system near a Feshbach resonance, to reduce the inter-species s-wave scattering length and thus enhance the effective nonlinearity χ leading to squeezing and entanglement. Spin-noise tomography leads to an inferred $\xi_R^2 = -8.2 \text{ dB}$, predicting an entanglement depth excluding less than 80 particles within three standard deviations (and a mean of 170 entangled atoms).




FIG. 16 **Direct experimental demonstration of a phase sensitivity beyond the standard quantum limit.** Left: Ramsey fringe scanned over a full 2π period. The blue data are obtained with a coherent spin state, the red data with a spin-squeezed state, showing a visibility of 92%. Solid lines are $\pm N/2$, measured for each phase setting, as a reference. Right: Ramsey fringe around the maximum-slope optimal point and averaged over several experimental realizations. The solid lines are fits through the lower and upper ends of the two standard deviation error bars. The gray shaded areas are the uncertainty regions for an ideal coherent spin state, leading to the standard quantum limit. Red data show a phase error 15% below the standard quantum limit. Taken from [Gross *et al.* \(2010\)](#).

2. Quantum interferometry

[Gross *et al.* \(2010\)](#) have also demonstrated a full Ramsey interferometer sequence with spin-squeezed states, see Fig. 16. Squeezed-state creation is followed by a rotation around the x axis to prepare a phase-squeezed state (*i.e.* squeezed along the y axis)—in effect constructing a nonlinear beam-splitter. After a short interrogation time $\chi t \ll 1$, during which the interaction is still active, a second $\pi/2$ beam splitter closes the Ramsey sequence. The directly measured phase sensitivity gain is $(\Delta\theta/\Delta\theta_{\text{SQL}})^2 = -1.4\text{ dB}$, corresponding to a reduction of phase variance 15% below the standard quantum limit. In a more recent version of the experiment, [Muessel *et al.* \(2014\)](#) have been able to scale the generation of spin-squeezed states up to $N = 12500$ particles using a chain of trapped independent condensates and exploiting a differential estimation of the phase. They demonstrated a spin squeezing $\xi_R^2 = -3.4\text{ dB}$ and a sensitivity 24% below the standard quantum limit. Ramsey interferometry using spin-squeezed states generated via one-axis twisting in Bose-Einstein condensate has recently found application to sense magnetic fields ([Muessel *et al.*, 2014](#); [Ockeloen *et al.*, 2013](#)), as discussed in more detail in Sec. VII.




FIG. 17 **Bell correlations in a Bose-Einstein condensate.** Measurement of the Bell correlation witness $\mathcal{W}$ of Eq. (47) on a spin-squeezed state as a function of the angle ϑ between the squeezing axis a and the axis n lying in the squeezing plane, with 1σ error bars. The red continuous line is the value of $\mathcal{W}$ computed from the measurement of $\zeta_a^2 = 4\langle\hat{J}_a^2\rangle/N$ and the fitted Rabi oscillation $C_n = C_b \sin(\vartheta)$ with $C_b = 2\langle\hat{J}_b\rangle/N$. Bell correlations are present in the blue shaded region. Right inset: representation of the same data (ζ_a^2 and C_b) as a single black point, with 1σ error bars. Red shaded region: entanglement detected by spin squeezing $\xi_R^2 < 1$, Eq. (28). Blue shaded region: Bell correlations detected by violation of the witness inequality (48). Adapted from [Schmied *et al.* \(2016\)](#).

3. Bell correlations

[Schmied *et al.* \(2016\)](#) have observed Bell correlations in spin-squeezed Bose-Einstein condensates of about $N = 480$ ^{87}Rb atoms trapped on an atom chip. Following the generation of the spin-squeezed state by one-axis twisting, the Bell correlation witness $\mathcal{W}$ of Eq. (47) is determined by measuring the second spin moment along the squeezed direction and combining it with spin contrast measurements along other projection axes, see Fig. 17. The measured values of $\mathcal{W}$ violate the inequality (47) and thus also the inequality (48). This proves the presence of Bell correlations and implies that the spin-squeezed states can violate the many-particle Bell inequality of [Tura *et al.* \(2014\)](#). The detection of Bell correlations in spin-squeezed Bose-Einstein condensates indicates that spin-squeezed states are useful beyond quantum metrology: they also contain the resource for quantum information tasks such as certified randomness generation or quantum key distribution ([Brunner *et al.*, 2014](#)).

Because there is no spatial separation between the atoms, the interpretation of the experiment of [Schmied *et al.* \(2016\)](#) relies on the common assumption that atoms do not communicate through unknown channels. This assumption can be relaxed if the atoms are spatially separated in an optical lattice potential, as proposed in [Pelis-](#)




FIG. 18 **Twist-and-turn dynamics.** Normalized quantum Fisher information (F_Q/N , red line) and inverse spin-squeezing parameter ($1/\xi_R^2$, blue line) as a function of $\omega_\pi t/\pi$ for the twist-and-turn model, where $\omega_\pi = \Omega\sqrt{\Lambda - 1}$. The inset shows the effective potential $W_\pi(z)$ initially (dotted line) and after a quench to a finite Λ (solid line, $\Lambda = 1.5$). The initial coherent spin state corresponds to a Gaussian wavepacket at the top of the barrier. The bottom panels show snapshots of the Wigner distribution at different times. The solid line is the mean-field separatrix and the dots are fixed points of the semiclassical model. Here $N = 100$ and the color scale is as in Fig. 5.

son *et al.* (2016). Many-body Bell tests with BECs have also been proposed in Laloë and Mullin (2009) and Mullin and Laloë (2008). In section III.D we discuss proposals to create entanglement between spatially separated spinor BECs, which could also be employed for Bell tests.

C. Twist-and-turn dynamics

The dynamical creation of entanglement in the one-axis twisting model is enriched—and accelerated, to some extent (Muessel *et al.*, 2015; Sorelli *et al.*, 2015)—by using the turning term $\hat{J}_x$ simultaneously with the twisting term J_z^2 in the Hamiltonian (50). The resulting twist-and-turn dynamics is experimentally studied from the starting point of a coherent spin state pointing in the $+x$ or $-x$ direction. The interaction is suddenly switched to a finite value of Λ in presence of linear coupling.

The twist-and-turn can be described as the dynamics of an effective relative-number wavepacket (Juliá-Díaz *et al.*, 2012). For an initial coherent spin state polarized along the positive x -axis, the effective Hamiltonian is given by Eq. (57). $W_0(z)$ can be well approximated as

a harmonic potential of frequency $\omega_0 = \Omega\sqrt{1 + \Lambda}$. Suddenly switching the interaction from $\Lambda = 0$ to a positive value (thus corresponding to squeezing the harmonic oscillator potential, *i.e.* $\omega_0/\Omega > 1$) gives rise to a breathing mode of the effective wavepacket, predicting number squeezing of the relative population. For a coherent spin state polarized along the negative x -axis, the effective Hamiltonian is $\hat{H}_z = -\frac{2}{N^2}\hat{\partial}_z\sqrt{1 - \hat{z}^2}\hat{\partial}_z + W_\pi(\hat{z})$ with $W_\pi(z) = -\frac{\Lambda z^2}{2} - \sqrt{1 - z^2}$. For $0 < \Lambda < 1$, $W_\pi(z)$ is harmonic with frequency $\omega_\pi = \Omega\sqrt{1 - \Lambda}$. Since $\omega_\pi/\Omega < 1$, a sudden switch of the interaction leads to a breathing motion predicting number anti-squeezing (and phase squeezing). Damping of the oscillations is observed for increasing values of Λ (Choi and Bigelow, 2005; Gordon and Savage, 1999) corresponding to the wavepacket feeling the anharmonicity of the effective potential. These predictions are recovered, for $0 < \Lambda \ll 1$, by a frozen-spin approximation (Law *et al.*, 2001) that consists of neglecting fluctuations of the mean spin, $\hat{J}_x$ being replaced by $\pm N/2$. The variances

$$(\Delta \hat{J}_z)^2 = \frac{N}{4} \left(\cos^2(\omega t) + \frac{\Omega^2}{\omega^2} \sin^2(\omega t) \right), \quad (70a)$$

$$(\Delta \hat{J}_y)^2 = \frac{N}{4} \left(\cos^2(\omega t) + \frac{\omega^2}{\Omega^2} \sin^2(\omega t) \right), \quad (70b)$$

show periodic oscillations and the spin-squeezing parameter reaches $\xi_R^2 = 1/(1 + \Lambda)$ ($\xi_R^2 = 1 - \Lambda$) for the coherent spin state pointing along the positive (negative) x axis.

The situation changes completely if $\Lambda > 1$ (Gordon and Savage, 1999; Micheli *et al.*, 2003). The potential $W_\pi(z)$ turns from a single-well to a double-well shape, with minima at $z_\pm = \pm\sqrt{1 - 1/\Lambda^2}$. The initial Gaussian wavepacket corresponding to the coherent spin state sits at the top of the barrier, see Fig. 18. The repulsive double-well potential tends to split the state giving rise to a macroscopic superposition corresponding to the wavepacket localized in the left and right well of $W_\pi(z)$. The semiclassical equations of motion $\dot{z} = -\frac{\partial H(z, \varphi)}{\partial \varphi}$ and $\dot{\varphi} = \frac{\partial H(z, \varphi)}{\partial z}$, where $H(z, \varphi)$ is given by Eq. (53), offer an alternative view. The fixed point $\Upsilon_\pi = (z = 0, \varphi = \pi)$ becomes unstable for $\Lambda > 1$, and two stable fixed points $\Upsilon_\pm = (z_\pm, \varphi = \pi)$ appear (Raghavan *et al.*, 1999; Smerzi *et al.*, 1997). The exact quantum dynamics shows that, at short times, the state stretches along the semiclassical separatrix passing through Υ_π , see Fig. 18. At longer times the quantum state wraps around $\Upsilon_\pm$. In this regime, spin squeezing is lost, while the quantum Fisher information—sensitive to the creation of macroscopic superposition states—continues increasing. For $\Lambda = 2$ the semiclassical separatrix reaches the poles of the Bloch sphere: the dynamics creates close to NOON states on a time scale $t \approx \ln(8N)/(\chi N)$ (Micheli *et al.*, 2003). For $\Lambda > 2$ the separatrix winds around the Bloch sphere, the maximum distance between the separatrix and the equator decreases, and the dynamics approaches that of




FIG. 19 Entanglement of non-spin-squeezed states. (a) Experimental Husimi distributions at different evolution times of twins-and-turn dynamics. (b) Normalized Fisher information (F/N , red diamonds) and inverse spin-squeezing parameter ($1/\xi_R^2$, blue circles) as a function of the tomography angle. The gray shaded area is only accessible by entangled states. At 26 ms, spin squeezing cannot identify the entanglement that is detected by the Fisher information. (c) Squared Hellinger distance as a function of the rotation angle θ at three evolution times: $t = 0$ corresponding to a separable state (black points), $t = 15$ ms corresponding to an optimal spin-squeezed state (gray points), and $t = 26$ ms corresponding to a state which is entangled but not spin squeezed (red points). The curvature of the quadratic fit is proportional to F/N . Taken from [Strobel *et al.* \(2014\)](#).

one-axis twisting.

1. Entanglement of non-spin-squeezed states

Twist-and-turn dynamics has been investigated by [Muessel *et al.* \(2015\)](#) and [Strobel *et al.* \(2014\)](#) using two internal hyperfine levels $|b\rangle = |F = 1, m_F = 1\rangle$ and $|a\rangle = |2, -1\rangle$ of ^{87}Rb atoms. A narrow Feshbach resonance is used to reduce the inter-species interaction. Coupling between the internal levels is provided by radiofrequency and microwave drive. The experiments start with $N \approx 400$ atoms in the $|b\rangle$ mode, followed by a $\pi/2$ Rabi pulse that prepares a coherent spin state aligned with the x -axis. Subsequently, the Rabi coupling is decreased to reach $\Lambda \approx 1.5$, and its phase adjusted to orient it along the negative x -axis. [Muessel *et al.* \(2015\)](#) and [Strobel *et al.* \(2014\)](#) have investigated the short-time spin-squeezing dynamics. For an evolution of 15 ms, [Strobel *et al.* \(2014\)](#) demonstrated $\xi_R^2 = -4.5$ dB [and inferred $\xi_R^2 = -7.1$ dB ([Muessel *et al.*, 2015](#))]. [Muessel *et al.* \(2015\)](#) demonstrated a spin squeezing $\xi_R^2 = -4$ dB using ~ 30 independent condensates in parallel (each condensate experiencing independently a twist-and-turn dynamics), with a total of $N = 10^4$ particles. For longer times, spin squeezing is quickly lost and the experimental Husimi distribution of the reconstructed state shows

the characteristic S-shape, see Fig. 19. [Strobel *et al.* \(2014\)](#) extracted the Fisher information, reaching values $F > N$ and thus demonstrating that the state is entangled. The experimental extraction of the Fisher information requires to rotate the state around the y -axis, and to collect the probability distributions of the relative particle number in $|a\rangle$ and $|b\rangle$ at different rotation angles. The Fisher information is obtained from a quadratic fit of the Hellinger distance (11), according to Eq. (12), between a reference distribution (at $\theta = 0$) and the distribution obtained at a finite θ . Experimental results are shown in Fig. 19(c).

D. Entanglement of two spatially-separated spinor Bose-Einstein condensates

Creating entanglement between two spatially separated, individually addressable spinor Bose-Einstein condensates offers interesting possibilities: in such a system, local manipulations and measurements can be performed on the spin state of each condensate separately and non-local quantum correlations between the measurement results can be directly observed. This is particularly relevant for experiments on EPR entanglement and Bell tests, where the spatial separation can be used to rule out unknown causal influences between the clouds.

[Bar-Gill *et al.* \(2011\)](#); [He *et al.* \(2011\)](#); and [Kurkjian *et al.* \(2013\)](#) have proposed schemes to generate EPR entanglement between two spatially separated Bose-Einstein condensates using elastic collisions. The condensates represent collective spins $\hat{J}^{(1)}$ and $\hat{J}^{(2)}$, respectively, which can be individually addressed and prepared in a coherent spin state. [Bar-Gill *et al.* \(2011\)](#) and [He *et al.* \(2011\)](#) have considered schemes where EPR entanglement is generated by first spin-squeezing each condensate and then interfering the squeezed states on a beam splitter. In the scheme of [Kurkjian *et al.* \(2013\)](#), on the other hand, a state-dependent potential is turned on such that the wave function of state $|a\rangle$ of the first condensate overlaps with the wave function of state $|b\rangle$ of the second condensate, picking up a collisional phase shift. Dropping constant and linear terms in $\hat{J}^{(i)}$, this realizes the Hamiltonian $H_{\text{BEC}} = \chi_1(\hat{J}_z^{(1)})^2 + \chi_2(\hat{J}_z^{(2)})^2 - \chi_{12}\hat{J}_z^{(1)}\hat{J}_z^{(2)}$, which generates entanglement between the two condensates in addition to spin-squeezing in each condensate. After an interaction time, the state-dependent potential is turned off and the condensates are spatially separated again for detection, revealing EPR entanglement between them ([Kurkjian *et al.*, 2013](#)). For long interaction times, macroscopic entangled states can be created, as also analyzed by [Byrnes \(2013\)](#), who investigates applications of entangled Bose-Einstein condensates in quantum information processing ([Byrnes *et al.*, 2012](#)). While such experiments have not yet been reported, we note that continuous-variable entanglement has been cre-

ated between spatially separated atomic vapor cells using a measurement-based scheme (Julsgaard *et al.*, 2001).

IV. ENTANGLEMENT VIA ATOMIC COLLISIONS: SPIN-MIXING DYNAMICS

The generation of correlated photon pairs via optical four-wave mixing, *e.g.* spontaneous parametric down-conversion in nonlinear crystals (Kwiat *et al.*, 1995), is one of the most widely used sources of entangled quantum states in optics (O'Brien *et al.*, 2009). When post-selecting the total number of photons in the pair distribution (let us indicate with $|+1\rangle$ and $|-1\rangle$ the signal and idler), the corresponding state is a twin-Fock, $|n\rangle_{+1}|n\rangle_{-1}$. The twin-Fock state has been used to overcome the standard quantum limit in an optical interferometer using up to $n = 2$ photons per pair (Krischek *et al.*, 2011; Nagata *et al.*, 2007; Xiang *et al.*, 2011). Without post-selection, four-wave mixing creates quadrature-squeezed light (Breitenbach *et al.*, 1997; Ou *et al.*, 1992; Slusher *et al.*, 1985; Walls, 1983; Wu *et al.*, 1986). Following the proposal of Caves (1981), squeezed light created in this way has been successfully used for optical interferometry beyond the standard quantum limit (Vahlbruch *et al.*, 2005), with direct application to gravitational wave detectors (Aasi *et al.*, 2013; Schnabel *et al.*, 2010). In the same spirit, the generation of correlated pairs of atoms has thus attracted large interest: many experiments have been proposed and performed. In the following, we review the most successful of these techniques, namely, spin-mixing dynamics in a spinor Bose-Einstein condensate (Kawaguchi and Ueda, 2012; Stamper-Kurn and Ueda, 2013), see Sec. IV.A. We discuss the creation of twin-Fock states in Sec. IV.B, and quadrature squeezing in Sec. IV.C. Finally, in Sec. IV.D we review alternative protocols for the creation of atom pairs.

A. Spinor Bose-Einstein condensates

1. Spin-changing collisions

When a Bose-Einstein condensate is confined in a far-off resonant optical dipole trap, the spin degree of freedom of the atoms evolves freely: spin-changing s-wave collisions, see Fig. 20, give rise to a coherent redistribution of atomic populations among Zeeman sub-levels while preserving the total magnetization (Ho, 1998; Ohmi and Machida, 1998). The quantum description of an ultracold gas of spin- F bosons requires introducing a vector order parameter with $2F + 1$ components, $\{\hat{\Psi}_{-F}, \hat{\Psi}_{-F+1}, \dots, \hat{\Psi}_F\}$ (Kawaguchi and Ueda, 2012; Stamper-Kurn and Ueda, 2013). Here $\hat{\Psi}_{m_F}$ is the atomic field annihilation operator associated with the hyperfine spin state $|F, m_F\rangle$. The single-mode approximation (Law *et al.*, 1998; Pu *et al.*, 1999) assumes that




FIG. 20 **Binary s-wave collision of two spin- F bosons.** (a) When two spin- F bosons in internal states $|F, m_F\rangle$ and $|F, m'_F\rangle$ approach each other (here, F is the hyperfine spin and $m_F, m'_F = -F, -F+1, \dots, F$ the magnetic quantum number), they couple to form a total spin $G = F + F'$. The combined internal state is given by $|G, m_G\rangle$. For bosons under elastic s-wave scattering, G is restricted to even values satisfying $0 \leq G \leq 2F$, *e.g.* $G = 0, 2$ for two $F = 1$ particles. (c) After the collision the atoms fly apart and their internal states are again well described by the hyperfine spins F . However, the magnetic quantum numbers may have changed: $|F, m_F\rangle|F, m'_F\rangle \mapsto |F, m''_F\rangle|F, m'''_F\rangle$. The conservation of total angular momentum imposes $m_F + m'_F = m''_F + m'''_F$.

the spatial atomic density distribution of all spin components is approximately equal, and not affected by the spin dynamics. The common scalar wavefunction $\phi(r)$ defining the spatial mode of the condensate is determined as the solution of the Gross-Pitaevskii equation, neglecting contributions from the spin-dependent interactions. In this approximation, the field operators are $\hat{\Psi}_{m_F}(r) = \phi(r)\hat{a}_{m_F}$, where $\hat{a}_{m_F}$ are annihilation operators that obey the usual bosonic commutation relations. We define as $\hat{N}_{m_F} = \hat{a}_{m_F}^\dagger \hat{a}_{m_F}$ the number of particles in the mode $|F, m_F\rangle$, and $\hat{N} = \sum_{m_F=-F}^F \hat{N}_{m_F}$ the total number of particles. The many-body Hamiltonian for a $F = 1$ Bose-Einstein condensate in presence of an external magnetic field B is (Law *et al.*, 1998; Pu *et al.*, 1999)

$$\begin{aligned} \hat{H}_{\text{SM}} = & [\hbar q + \hbar\lambda(2\hat{N}_0 - 1)](\hat{N}_{+1} + \hat{N}_{-1}) + \\ & + 2\hbar\lambda(\hat{a}_{-1}^\dagger \hat{a}_{+1}^\dagger \hat{a}_0 \hat{a}_0 + \hat{a}_0^\dagger \hat{a}_0^\dagger \hat{a}_{-1} \hat{a}_{+1}), \end{aligned} \quad (71)$$

suppressing terms proportional to the conserved magnetization, $\hat{N}_{+1} - \hat{N}_{-1}$, and total number of atoms. Here, $q \propto B^2$ accounts for the quadratic Zeeman effect and $\hbar\lambda = g_2 \int d^3r |\phi(r)|^4$, with $g_2 = \frac{4\pi\hbar^2(a_2 - a_0)}{3M}$, and a_G the scattering lengths for s-wave collisions in the $G = 0, 2$ allowed channels. The parameter λ can be positive, as in ^{23}Na , or negative, as in ^{87}Rb . The single-mode approximation is valid if the system size is much smaller than the spin healing length, $\sqrt{\hbar^2/2M|g_2|\rho_0}$, where ρ_0 is the density of the $m_F = 0$ component, giving the minimum size of a spin-domain (Stamper-Kurn and Ueda, 2013). The second term in Eq. (71) accounts for the collisional shift arising due to elastic s-wave scattering. The last term in Eq. (71) describes the coherent and reversible creation of a pair of atoms in the magnetic sublevels $m_F = \pm 1$ from the scattering of two atoms in $m_F = 0$ (Duan *et al.*,

2000b; Pu and Meystre, 2000). The spin dynamics of an initial condensate in $m_F = 0$ is thus formally analogous to optical spontaneous four-wave mixing (Goldstein and Meystre, 1999): atom-atom interaction plays the role of the nonlinear Kerr medium, the $m_F = 0$ condensate is equivalent to a coherent pump field with the external trapping potential corresponding to a high-finesse cavity, and the condensates in $m_F = \pm 1$ can be identified as signal and idler. Remarkably, spin changing collisions are unaffected by the linear Zeeman shift from a homogeneous magnetic field. Only higher-order effects, such as a quadratic Zeeman shift or a linear Zeeman shift from an inhomogeneous magnetic field, affect it.

It is useful to rewrite the Hamiltonian (71) in terms of spin operators

$$\begin{aligned}\hat{S}_x &= \frac{\hat{a}_0^\dagger \hat{J} + \hat{a}_0 \hat{J}^\dagger}{2}, & \hat{\mathcal{A}}_x &= \frac{\hat{a}_0^\dagger \hat{a} + \hat{a}_0 \hat{a}^\dagger}{2}, \\ \hat{S}_y &= \frac{\hat{a}_0^\dagger \hat{J} - \hat{a}_0 \hat{J}^\dagger}{2i}, & \hat{\mathcal{A}}_y &= \frac{\hat{a}_0^\dagger \hat{a} - \hat{a}_0 \hat{a}^\dagger}{2i}, \\ \hat{S}_z &= \frac{\hat{a}_0^\dagger \hat{a}_0 - \hat{J}^\dagger \hat{J}}{2}, & \hat{\mathcal{A}}_z &= \frac{\hat{a}_0^\dagger \hat{a}_0 - \hat{a}^\dagger \hat{a}}{2},\end{aligned}\quad (72)$$

where $\hat{J} = (\hat{a}_{+1} + \hat{a}_{-1})/\sqrt{2}$ and $\hat{a} = (\hat{a}_{+1} - \hat{a}_{-1})/\sqrt{2}$ are symmetric and antisymmetric combinations of $\hat{a}_{+1}$ and $\hat{a}_{-1}$. The spin operators $\hat{S} = \{\hat{S}_x, \hat{S}_y, \hat{S}_z\}$ and $\hat{\mathcal{A}} = \{\hat{\mathcal{A}}_x, \hat{\mathcal{A}}_y, \hat{\mathcal{A}}_z\}$ do not commute with each other, and define two SU(2) subspaces.¹⁸ With these operators, the Hamiltonian (71) takes the compact form (Duan *et al.*, 2002; Law *et al.*, 1998)

$$\hat{H}_{\text{SM}} = (4\hbar\lambda\hat{S}_x^2 - \frac{2\hbar q}{3}\hat{S}_z) + (4\hbar\lambda\hat{\mathcal{A}}_y^2 - \frac{2\hbar q}{3}\hat{\mathcal{A}}_z), \quad (73)$$

which highlights the presence of nonlinear spin terms. It should be noticed that a symmetric radio-frequency coupling between the $m_F = 0, \pm 1$ modes with Rabi frequency Ω_{rf} corresponds to a rotation of the $\hat{S}$ vector around the x axis, $\hat{H}_{\text{rf}} = \frac{\hbar\Omega_{\text{rf}}}{2\sqrt{2}}(\hat{a}_0^\dagger \hat{a}_{+1} + \hat{a}_0^\dagger \hat{a}_{-1} + h.c.) = \hbar\Omega_{\text{rf}}\hat{S}_x$. A relative phase shift between the $m_F = 0$ and $m_F = \pm 1$ modes corresponds to a rotation of both $\hat{S}$ and $\hat{\mathcal{A}}$ around the z axis, $e^{-i\theta(\hat{N}_0 - \hat{N}_{+1})/2}e^{-i\theta(\hat{N}_0 - \hat{N}_{-1})/2} = e^{-i\theta\hat{S}_z}e^{-i\theta\hat{\mathcal{A}}_z}$.

¹⁸ The operators (72) are related to the angular momentum $\hat{L}$ and quadrupole tensor $\hat{Q}$ operators used to describe spin-1 condensates (Hamley *et al.*, 2012), see also (Law *et al.*, 1998; Müstecaplıoğlu *et al.*, 2002; Sau *et al.*, 2010). We have the following correspondence: $\hat{L}_x = 2\hat{S}_x$, $\hat{L}_y = -2\hat{\mathcal{A}}_y$ and $\hat{L}_z = \hat{N}_{+1} - \hat{N}_{-1}$ are the components of $\hat{L}$, $Q_{yz} = -2\hat{S}_y$, $Q_{zx} = 2\hat{\mathcal{A}}_x$, $Q_{zz} - Q_{yy} = -4\hat{S}_z$, $Q_{xx} - Q_{zz} = 4\hat{\mathcal{A}}_z$ and $Q_{xy} = i(\hat{a}_{-1}^\dagger \hat{a}_{+1} - \hat{a}_{+1}^\dagger \hat{a}_{-1})$ are the components of $\hat{Q}$.

2. Quantum spin-mixing dynamics in the low-depletion limit

We consider the evolution of an initial large ($N \gg 1$) condensate prepared in $m_F = 0$. The low-depletion limit, *i.e.*, $N_0 \approx N$, is analyzed by replacing the mode operator $\hat{a}_0$ in Eq. (71) with $\sqrt{N_0}$. In this approximation, the condensate serves as an unlimited particle resource for the parametric amplification of the $m_F = \pm 1$ modes. We obtain the quadratic Hamiltonian $\hat{H}_{\text{SM}} = \hbar\alpha(\hat{a}_{+1}^\dagger \hat{a}_{+1} + \hat{a}_{-1}^\dagger \hat{a}_{-1}) + \hbar\beta(\hat{a}_{-1}^\dagger \hat{a}_{+1}^\dagger + \hat{a}_{-1} \hat{a}_{+1})$, where $\alpha = q + \lambda(2N_0 - 1)$ and $\beta = 2\lambda N_0$, that can be diagonalized by a Bogoliubov transformation (Duan *et al.*, 2000b; Pu and Meystre, 2000), see also (Truax, 1985). The unitary evolution $|\psi_{\text{SM}}(t)\rangle = e^{-i\hat{H}_{\text{SM}}t/\hbar}|\text{vac}\rangle$ can be calculated exactly, giving

$$|\psi_{\text{SM}}(t)\rangle = \sum_{n=0}^{+\infty} \frac{[-i\beta\tau \sin(\frac{t}{\tau})]^n}{[\cos(\frac{t}{\tau}) + i\alpha\tau \sin(\frac{t}{\tau})]^{n+1}} |n\rangle_{+1} |n\rangle_{-1}, \quad (74)$$

where $\tau = 1/\sqrt{\alpha^2 - \beta^2}$. This state is characterized by a vanishing population difference $\Delta^2(\hat{N}_{+1} - \hat{N}_{-1}) = 0$, while the $m_F = \pm 1$ modes are nonempty, $\langle \hat{N}_{\pm 1} \rangle = \frac{\beta^2}{\alpha^2 - \beta^2} \sin^2(t\sqrt{\alpha^2 - \beta^2})$, and characterized by large (super-Poissonian) population fluctuations, $\Delta^2 N_{\pm 1} = \langle \hat{N}_{\pm 1} \rangle (\langle \hat{N}_{\pm 1} \rangle + 1)$ (Klempt *et al.*, 2010; Leslie *et al.*, 2009; Liu *et al.*, 2009). It is also interesting to consider the quadratures

$$2(\Delta\hat{Q}_s)^2 = V_Q \cos^2(\phi - \phi_Q) + V_Q^{-1} \sin^2(\phi - \phi_Q), \quad (75a)$$

$$2(\Delta\hat{Q}_a)^2 = V_Q \sin^2(\phi - \phi_Q) + V_Q^{-1} \cos^2(\phi - \phi_Q), \quad (75b)$$

where $\hat{Q}_s = \cos\phi\hat{X}_s + \sin\phi\hat{P}_s$, $\hat{X}_s = (\hat{J} + \hat{J}^\dagger)/\sqrt{2}$, $\hat{P}_s = (\hat{J} - \hat{J}^\dagger)/i\sqrt{2}$, and an analogous definitions of $\hat{Q}_a$ in terms of the operators $\hat{a}$ and $\hat{a}^\dagger$ introduced above. The time-dependent coefficients are $V_Q = 1 + 2\langle \hat{N}_{\pm 1} \rangle - 2\Delta\hat{N}_{\pm 1}$ and $\cos(2\phi_Q) = \frac{\alpha \langle \hat{N}_{\pm 1} \rangle}{\beta \Delta\hat{N}_{\pm 1}}$.

For $\alpha^2 < \beta^2$, τ is imaginary and the condensate is dynamically unstable (Sadler *et al.*, 2006). The instability is characterized by the exponential increase of population in the $m_F = \pm 1$ modes, $\langle \hat{N}_{\pm 1} \rangle = \frac{\beta^2}{\beta^2 - \alpha^2} \sinh^2(t\sqrt{\beta^2 - \alpha^2})$. On resonance ($\alpha = 0$) the spin-mixing Hamiltonian becomes

$$\hat{H}_{\text{SM}} = 2\hbar\lambda N_0 (\hat{a}_{-1}^\dagger \hat{a}_{+1}^\dagger + \hat{a}_{-1} \hat{a}_{+1}), \quad (76)$$

which generates the familiar two-mode squeezed-vacuum state (Walls and Milburn, 1994),

$$|\psi(t)\rangle = \sum_{n=0}^{+\infty} \frac{(-i \tanh r)^n}{\cosh r} |n\rangle_{+1} |n\rangle_{-1}, \quad (77)$$

where $r = |\beta|t = 2|\lambda|N_0 t$ and $\langle \hat{N}_{\pm 1} \rangle = \sinh^2 r$. Using the operators $\hat{J}$ and $\hat{a}$, the Hamiltonian (76) can be rewritten

as $\hat{H}_{\text{SM}} = \frac{\hbar\beta}{2}(\hat{J}^\dagger\hat{J}^\dagger + \hat{J}\hat{J}) - \frac{\hbar\beta}{2}(\hat{a}^\dagger\hat{a}^\dagger + \hat{a}\hat{a})$, which generates a single-mode squeezed-vacuum (Walls and Milburn, 1994) in each mode. In the low-depletion limit, Eq. (73) becomes

$$\hat{H}_{\text{SM}} = 2\hbar\lambda[(\hat{S}_x^2 - \hat{S}_y^2) - (\hat{\mathcal{A}}_x^2 - \hat{\mathcal{A}}_y^2)], \quad (78)$$

corresponding to two-axis counter-twisting (Kitagawa and Ueda, 1993) for the $\mathcal{S}$ and $\mathcal{A}$ spins. In this case, Eq. (75a) reduces to $2(\Delta\hat{Q}_s)^2 = e^{2r}\sin^2(\phi - \frac{\pi}{4}) + e^{-2r}\cos^2(\phi - \frac{\pi}{4})$, and Eq. (75b) to $2(\Delta\hat{Q}_a)^2 = e^{2r}\cos^2(\phi - \frac{\pi}{4}) + e^{-2r}\sin^2(\phi - \frac{\pi}{4})$. The quadrature $(\Delta\hat{Q}_a)^2$ is squeezed by a factor e^{-2r} below the vacuum level at an optimal angle $\phi = \pi/4$ [$\phi = 3\pi/4$], and anti-squeezed by a factor e^{2r} at $\phi = 3\pi/4$ [$\phi = \pi/4$]. For a sufficiently long evolution time, $t \gtrsim 1/(2|\lambda|N_0)$ spin dynamics if modified by the depletion of the condensate. The initially exponential growth of population in $m_F = \pm 1$ slows down and finally stops (Law *et al.*, 1998; Mias *et al.*, 2008)

For $\alpha^2 \geq \beta^2$ the spin-mixing dynamics is stable and the population of $m_F = \pm 1$ oscillates in time, $\langle \hat{N}_{\pm 1} \rangle = \frac{\beta^2}{\alpha^2 - \beta^2} \sin^2(t\sqrt{\alpha^2 - \beta^2})$. For $\alpha^2 \gg \beta^2$ the amplitude of this oscillation becomes negligible and the system remains stable in $m_F = 0$.

Typical experiments exploring the quantum spin-mixing dynamics (Chang *et al.*, 2004; Klempt *et al.*, 2010; Leslie *et al.*, 2009; Liu *et al.*, 2009; Peise *et al.*, 2015b; Schmaljohann *et al.*, 2004) start with a condensate in the stable configuration. The condensate is suddenly quenched below the quantum critical point by lowering the magnetic field, or using microwave dressing to reach the resonance condition (Gerbier *et al.*, 2006). After spin-mixing dynamics, the trap is turned off and a Stern-Gerlach field is used to separate the m_F components during a time-of-flight expansion.

Finally, it is worth recalling that the discussion so far has been focused on one spatial eigenmode. A rich resonance structure is expected when taking into account many spatial modes, as experimentally shown by Klempt *et al.* (2010) and Scherer *et al.* (2010).

B. Population correlations and twin-Fock state

Quantum spin-mixing dynamics can turn a large fraction of an atomic Bose-Einstein condensate into a mixture of perfectly correlated pairs of atom, *i.e.* the twin Fock state $|n\rangle_{+1}|n\rangle_{-1}$ (with large fluctuations of n), in the $m_F = \pm 1$ Zeeman states. This state is particle entangled and useful in a Ramsey interferometer to reach sensitivities beyond the standard quantum limit, see Sec. II.C.6.




FIG. 21 **Population correlations after spin-mixing dynamics.** Left panel: $\langle \hat{N}_{+1} + \hat{N}_{-1} \rangle$ as a function of the total atom number N (red circles). Error bars are fluctuations (1 standard deviation). Blue circles are $\Delta^2(\hat{N}_{+1} - \hat{N}_{-1})$. The inset shows the distribution of $\hat{N}_{+1} + \hat{N}_{-1}$ for $250 < N < 300$. The black line is a fitted squeezed-vacuum distribution corresponding to $r \approx 2$. Taken from Gross *et al.* (2011). Right panel: standard deviation of the population difference $(\hat{N}_{+1} - \hat{N}_{-1})/2$ (red line), compared to the projection noise $\sqrt{N_{+1} + N_{-1}}/2$ (dashed line), and $\sqrt{(N_{+1} + N_{-1})/4 + \sigma_{\text{dn}}^2}$ (dot-dashed line) taking into account a number-independent detection noise level $\sigma_{\text{dn}} = 20$ (dotted line). The blue line is the experimental result for unentangled atoms. The shaded area indicates the standard deviation. Adapted from Lücke *et al.* (2011)

1. Number squeezing

The presence of correlations in the populations of $m_F = \pm 1$ states after spin dynamics has been first demonstrated by Bookjans *et al.* (2011); Gross *et al.* (2011); and Lücke *et al.* (2011). These experiments have measured the population of $m_F = \pm 1$ states. A number squeezing parameter $\xi_N^2 = \Delta^2(\hat{N}_{+1} - \hat{N}_{-1})/(N_{+1} + N_{-1})$ up to $\xi_N^2 \approx -7$ dB [$\xi_N^2 = -12.4$ dB has been reported more recently by Lücke *et al.* (2014)] below the projection noise level has been demonstrated, see Figs. 21. The achieved squeezing is mainly limited by detection noise, for short times, and particles losses, for longer time evolutions.

2. Quantum interferometry with a twin-Fock state

Lücke *et al.* (2011) have investigated the phase sensitivity of the paired atoms produced via spin dynamics. The experimental sequence starts with the spin-mixing dynamics of a ^{87}Rb condensate prepared in $|\hat{F} = 2, m_F = 0\rangle$. The generated atom pairs in $|2, \pm 1\rangle$ (due to the short evolution time, the populations in $|2, \pm 2\rangle$ can be neglected due to the quadratic Zeeman shift detuning) are coupled via a series of microwave pulses of variable duration, see Fig. 22. This forms an internal-state beam splitter $e^{-i\theta\hat{J}_x}$, where the rotation angle θ is estimated via the measurement of $\hat{J}_z^2$. The output state is post-selected to a total number $N_{+1} + N_{-1} \approx 10^4$. Ideally, one thus expects a twin-Fock state as input of




FIG. 22 **Twin-Fock interferometry.** The inset of panel (a) shows experimental operations: (1) spin dynamics, (2) a resonant microwave π pulse between $|2, -1\rangle$ and $|1, 0\rangle$, (3) a pulse of variable duration, defining θ , and (4) a second π pulse. a) Second moment $\langle \hat{J}_z^2 \rangle$ of the population imbalance $\hat{J}_z = (\hat{N}_{+1} - \hat{N}_{-1})/2$ as a function of θ (dots with error bars) for post-selected numbers of atoms between 6400 and 7600. The solid line is a polynomial fit (with grey uncertainty region); the dotted line is the theoretical prediction including detection noise. b) Same for $\langle (\Delta \hat{J}_z)^2 \rangle = \langle \hat{J}_z^4 \rangle - \langle \hat{J}_z^2 \rangle^2$. c) Phase estimation uncertainty obtained via error propagation (solid orange line with grey uncertainty region) compared to the theoretical prediction (dashed orange line) and Cramér-Rao bound (dotted orange line) including detection noise only. At $\theta = 0.015$ rad, the phase variance lies 1.61 dB below the standard quantum limit (black dashed line). Adapted from Lücke *et al.* (2011).

the beam splitter operations. Applying error propagation $\Delta\theta = (\Delta \hat{J}_z^2)/|d\langle \hat{J}_z^2 \rangle/d\theta|$, it was possible to reach a sensitivity 1.61 dB below the standard quantum limit at an optimal rotation angle, see Fig. 22.

3. Particle entanglement

Lücke *et al.* (2014) have investigated particle entanglement in the atomic twin-Fock state. After spin-mixing dynamics of a ^{87}Rb condensate, the number of particles in $m_F = \pm 1$ is post-selected for a total number of $N \approx 8000$. Entanglement is detected by using the inequality $\xi_D^2 < 1$, Eq. (40), further extended to include the detection of arbitrary k -particle entanglement (Lücke

et al., 2014). Experimental data exclude an entanglement depth of less than 28 atoms with two standard deviations of confidence, and show an average entanglement depth of 68 atoms. It should be noted that $\xi_D^2 < 1$ signals entanglement independently from its metrological usefulness. A condition for useful k -particle entanglement, based on second moments of spin components, was derived by Apellaniz *et al.* (2015a). Useful k -particle entanglement is a more demanding resource than algebraic k -particle entanglement: in Apellaniz *et al.* (2015a), similar experimental data as those reported by Lücke *et al.* (2014) show a useful entanglement depth of about 3 atoms.

C. Quadrature squeezing and squeezed-vacuum state

1. Quadrature squeezing

The experiments (Gross *et al.*, 2011; Hamley *et al.*, 2012; Peise *et al.*, 2015a) have investigated quadrature squeezing in the state generated via spin-mixing dynamics in the low-depletion limit, with negligible population transfer from $m_F = 0$ to $m_F = \pm 1$. In this regime, the quadrature variance Eq. (75a) can be rewritten as $2(\Delta \hat{Q}_\phi)^2 = N\Delta^2(\hat{\mathcal{S}}_x \cos \phi + \hat{\mathcal{S}}_y \sin \phi)/\langle \hat{\mathcal{S}}_z \rangle^2$ and $\langle \hat{\mathcal{S}}_z \rangle \approx N/2$ [and analogous relation between Eq. (75b) and the $\hat{\mathcal{A}}$ spin]: quadrature squeezing corresponds to spin squeezing. Furthermore, in the low-depletion limit, a symmetric radiofrequency coupling between the $m_F = 0, \pm 1$ modes corresponds to a displacement operation, $e^{-i\Omega_{\text{rf}}t\hat{\mathcal{S}}_x} = e^{-i\Omega_{\text{rf}}t\sqrt{N}(\hat{\mathcal{S}}_+ + \hat{\mathcal{S}}_-)/2}$, and can be used to realize an atomic homodyne measurement, as first shown by Gross *et al.* (2011), see also (Peise *et al.*, 2015a), the local oscillator being the condensate in $m_F = 0$. Hamley *et al.* (2012) have used $N = 4.5 \times 10^4$ ^{87}Rb and characterized the spin-squeezed states via noise tomography, see Fig. 23. Values up to $\xi_R^2 = -8.3$ dB (-10.3 dB inferred) below the SQL are reported.

2. Interferometry with squeezed vacuum

Quadrature squeezing is a resource for quantum interferometry, see Sec. II.C.5.b. Within spin-mixing dynamics, squeezing occurs in the quadratures corresponding to the symmetric $|s\rangle = (|1, +1\rangle + |1, -1\rangle)/\sqrt{2}$ and the anti-symmetric $|a\rangle = (|1, +1\rangle - |1, -1\rangle)/\sqrt{2}$ combinations of the $|F = 1, m_F = \pm 1\rangle$ modes. Kruse *et al.* (2016) have realized an atomic clock that exploits the squeezed-vacuum states created in the symmetric mode $|s\rangle$. An initial condensate of $N = 10^4$ atoms undergoes spin-mixing dynamics using the $F = 1$ manifold of ^{87}Rb , in the low-depletion regime ($\langle \hat{N}_0 \rangle \gg \langle \hat{N}_{\pm 1} \rangle$ after spin dynamics). The interferometer consists of a rf coupling (corresponding to a balanced beam splitter between the condensate in $|1, 0\rangle$ and $|s\rangle$, $\hat{H}_{\text{rf}} = \hbar\Omega_{\text{rf}}\hat{\mathcal{S}}_x$), a




FIG. 23 **Quadrature squeezing.** The upper panel shows the quadrature variance $4\Delta^2(\hat{s}_x \cos \phi + \hat{s}_y \sin \phi)/N$ at different evolution times (colored lines and symbols) as functions of the quadrature angle ϕ . Symbols are experimental results obtained via state tomography, whereas the solid lines are theoretical predictions. Squeezing down to -8.3 dB below the vacuum limit is observed. The bottom panels show reconstructed phase space distributions at $t = 15$ ms and $t = 45$ ms; the black ellipses are the $1/\sqrt{e}$ uncertainty ellipse predicted theoretically. Adapted from Hamley *et al.* (2012).

relative phase shift θ implemented by a detuned π pulse between $|1,0\rangle$ and $|2,0\rangle$, and a second balanced beam splitter between $|1,0\rangle$ and $|s\rangle$. The anti-symmetric mode $|a\rangle$ is left unchanged by the interferometer transformation (up to an overall phase), while the $|s\rangle$ is rotated in the Bloch sphere identified by the operators $\{\hat{s}_x, \hat{s}_y, \hat{s}_z\}$. The phase signal is given by $(\hat{N}_{+1} + \hat{N}_{-1})/N$, which shows Ramsey fringes as a function of θ . The experiment of Kruse *et al.* (2016) reports a clock sensitivity reaching 2 dB below the standard quantum limit. The expected sensitivity is $\Delta\theta = e^{-r}/\sqrt{N}$ (Caves, 1981). Ideally, interferometry using the squeezed-vacuum state in one input port and a coherent state in the second port can reach the Heisenberg limit $\Delta\theta = 1/N$ when the input states in each port are equally populated (Pezzè and Smerzi, 2008).

3. Nonlinear SU(1,1) interferometry

Spin-mixing dynamics can be also used to realize a SU(1,1) interferometer, as first proposed by Yurke *et al.* (1986) in optics [realized recently with a bright laser source by Hudelist *et al.* (2014)], and further analyzed by Gabbrielli *et al.* (2015) and Marino *et al.* (2012) for atoms. The basic idea of this interferometric scheme is to exploit spin-mixing dynamics to implement a nonlinear beam splitter, where, on average, $\mathcal{N} = \langle \hat{N}_{+1} + \hat{N}_{-1} \rangle$ atoms are transferred in pairs from an initial condensate prepared in $m_F = 0$ to $m_F = \pm 1$. The system acquires a relative phase $\theta = 2\theta_0 - (\theta_{+1} + \theta_{-1})$ between $m_F = 0$ and $m_F = \pm 1$ modes. A second spin-mixing dynamics closes the interferometer. The final populations in $m_F = 0, \pm 1$ depend on θ . When treating the condensate in $m_F = 0$ as an undepletable source of atomic pairs, the predicted phase sensitivity is

$$\Delta\theta = \sqrt{\frac{\mathcal{N}(\mathcal{N} + 2) \cos^2(\theta/2) + 1}{\mathcal{N}(\mathcal{N} + 2) \sin^2(\theta/2)}}. \quad (79)$$

Equation (79) reaches $\Delta\theta = 1/\sqrt{\mathcal{N}(\mathcal{N} + 2)}$ at the optimal working point $\theta = \pi$ (a dark fringe, where no particle is found in the output $m_F = \pm 1$ states). An analysis beyond the low-depletion limit shows that this interferometer scheme can reach $\Delta\theta = \mathcal{O}(1/N)$ (Gabbrielli *et al.*, 2015), namely, the Heisenberg scaling where N is the total number of particles.

Linnemann *et al.* (2016) have experimentally realized a nonlinear SU(1,1) interferometer within the $F = 2$ manifold of ^{87}Rb . The experiment is performed in the low-depletion regime, $\mathcal{N} = 2.8$, and using $N \approx 400$ atoms. The interferometer is probed by tuning the relative phase between the $m_F = 0$ condensate and the $m_F = \pm 1$ modes to the dark fringe $\theta = \pi$, where maximum sensitivity is expected. The interferometrically measured phase θ is imprinted via a second-order Zeeman shift by applying a magnetic field for varying times, and is read out from the mean total number of atoms in the $m_F = \pm 1$ modes. Results of an error propagation analysis demonstrate a sensitivity $\Delta\theta$ close to the theoretical prediction $1/\sqrt{\mathcal{N}(\mathcal{N} + 2)}$ of Eq. (79).

4. Continuous variable and EPR entanglement

The two-mode squeezed-vacuum state (77) realizes—in field modes—the position-momentum correlations that are at the heart of the Einstein-Podolsky-Rosen (EPR) criterion (Reid, 1989), see Sec. II.C.9.a. We recall that¹⁹

¹⁹ Notice the relations $\hat{Q}_s(\phi) = [\hat{X}_{+1}(\phi) + \hat{X}_{-1}(\phi)]/\sqrt{2}$ and $\hat{Q}_a(\phi) = [\hat{X}_{+1}(\phi) - \hat{X}_{-1}(\phi)]/\sqrt{2}$, where $\hat{X}_{\pm 1}(\phi) = (\hat{a}_{\pm 1}e^{-i\phi} +$




FIG. 24 **Einstein-Podolsky-Rosen entanglement with spinor condensates.** a) Variances $V_{X(\phi)}^+$ and $V_{X(\phi)}^-$ as functions of the local oscillator phase ϕ . The dashed line is the quadrature variance for the vacuum state. The lowest measured quadrature variance is $V_X^- = 0.42$, corresponding to a squeezing of 3.77 dB below the vacuum limit. b) Product $V_{X(\phi)}^- \times V_{X(\phi+\pi/2)}^+$: data points below the dashed line violate the inequality (46) and thus signal Einstein-Podolsky-Rosen entanglement. Data reach $V_{X(\phi)}^- \times V_{X(\phi+\pi/2)}^+ = 0.18(3)$, which is 2.4 standard deviations below the limit of $1/4$. c) Sum $V_{X(\phi)}^+ + V_{X(\phi+\pi/2)}^-$: data points below the dotted line violate the inequality (45) and thus signal entanglement between the $m_F = \pm 1$ modes. Data reach $V_{X(\phi)}^+ + V_{X(\phi+\pi/2)}^- = 0.85(8)$, which is 15 standard deviations below the limit of 2. The dashed line is a sufficient condition for Einstein-Podolsky-Rosen entanglement. Adapted from Peise *et al.* (2015a).

$V_{X(\phi)}^\pm = \cosh(2r) \mp \sin(2\phi) \sinh(2r)$. As the angle ϕ is varied, these quadrature variances oscillate between a maximum value e^{2r} and a minimum value e^{-2r} , which is below the value of 1 of the unsqueezed vacuum state. In the limit $r \rightarrow \infty$, at $\phi = 3\pi/4$ ($\phi = \pi/4$), we have perfect correlations between $\hat{X}_{+1}$ and $\hat{X}_{-1}$ ($\hat{X}_{+1}$ and $-\hat{X}_{-1}$), as well as between $\hat{P}_{+1}$ and $-\hat{P}_{-1}$ ($\hat{P}_{+1}$ and $\hat{P}_{-1}$), where $\hat{P}_{\pm 1}(\phi) = \hat{X}_{\pm 1}(\phi + \pi/2)$. According to the criteria (45) and (46) discussed in Sec. II.C.9, the two-mode squeezed-vacuum state is mode entangled for $r > 0$ and fulfills the EPR criterion for $r > \ln \sqrt{2}$ (Reid *et al.*, 2009). Experimentally, the quadrature modes are accessed via atomic homodyne detection (Gross *et al.*, 2011), in analogy to optical homodyning (Ou *et al.*, 1992), where the condensate in $m_F = 0$ is used as local oscillator.²⁰ The quadra-

tures $X_\pm(\phi)$ are proportional to the number of particles $\hat{N}_{+1} \pm \hat{N}_{-1}$ measured after a radiofrequency pulse coupling $m_F = 0$ with $m_F = \pm 1$ (Gross *et al.*, 2011). With spinor condensates (Duan *et al.*, 2002), continuous variable entanglement, *e.g.* the violation of Eq. (45), was first demonstrated by Gross *et al.* (2011), while EPR entanglement, *e.g.* the violation of Eq. (46) was observed by Peise *et al.* (2015a), see Fig. 24. The experimental violation of Eq. (46) demonstrates a form of entanglement intrinsically connected with local realism. The demonstration of the continuous-variable EPR paradox with massive particles would additionally require space-like separation of measurements to rule out causal influences, which has not yet been experimentally achieved.

D. Other protocols to create correlated atomic pairs

Spin-mixing dynamics is not the only possibility to create correlated atom pairs. Alternative methods have been studied and experimentally implemented. Relative number squeezing between two outgoing Bose-Einstein condensates with opposite momenta was obtained in the collisional de-excitation of a one-dimensional quasi-condensate, where the reduced dimensionality restricted the number of available modes (Bücker *et al.*, 2011). The atoms are initially prepared in a highly non-equilibrium state such that the only allowed de-excitation channel is a two-particle collision process, emitting atom pairs. In Bücker *et al.* (2011) a number squeezing of up to -4.3 dB was observed with $\mathcal{N} = 200$ converted particles. In Jaskula *et al.* (2010), sub-Poissonian atom number fluctuations, with 0.5 dB of number squeezing, were observed between outgoing modes of opposing momenta in the halo produced by the s-wave scattering of two Bose-Einstein condensates of metastable ^4He . The amount of detected squeezing was limited by the large number of outgoing collisional modes (isotropically distributed over the scattering sphere) and the correspondingly small number of atoms per mode. With this scheme (Lewis-Swan and Kheruntsyan, 2014) it was possible to realize the Hong-Ou-Mandel experiment (Lopes *et al.*, 2015) where beam splitters are realized using Bragg scattering on an optical lattice (Bonneau *et al.*, 2013). Each atom of the atomic pair in the motional degree of freedom is sent to the input channel of a beam splitter. When two inputs are indistinguishable, they emerge together in one of the output channels, demonstrating two-particle interference.

²⁰ In the case of a finite number of particles in the $m_F = 0$ mode (Ferris *et al.*, 2008; Raymer *et al.*, 2003), the inequalities (45) and

(46) generalize to $V_{X(\phi)}^\pm + V_{P(\phi)}^\mp \geq 2 - \frac{N_{+1} + N_{-1}}{N_0}$, and $V_{X(\phi)}^- \times V_{P(\phi)}^+ \geq \frac{1}{4} (1 - \frac{N_{-1}}{N_0})^2$, respectively.

V. ENTANGLEMENT CREATION VIA ATOM-LIGHT INTERACTION

The light-matter quantum interface finds important applications in several areas of quantum information processing (Hammerer *et al.*, 2010; Kimble, 2008). Here we review the many successful proposals and experiments that exploit such interfaces for the creation of useful entangled states for quantum metrology. This quest started in the late 1990s with pioneering experiments using room-temperature vapor cells (Kuzmich *et al.*, 2000, 1999) and laser-cooled atomic gases (Hald *et al.*, 1999), which demonstrated the reduction of atomic spin (or pseudospin) fluctuations. In these experiments, the atom-light interface was implemented in free space using optically-thick atomic ensembles (Kuzmich and Polzik, 2003). Further improvements of this approach allowed to realize metrological spin squeezing (Appel *et al.*, 2009) and to use it in proof-of-principle atomic clocks (Louchet-Chauvet *et al.*, 2010) and magnetometers (Sewell *et al.*, 2012, 2014; Wasilewski *et al.*, 2010). Optical cavities can be used to effectively increase the optical depth of the atomic sample and represent a versatile system with rich possibilities for creating entanglement between atoms (Miller *et al.*, 2005; Ritsch *et al.*, 2013; Tanji-Suzuki *et al.*, 2011). Cavity-based atom-light interfaces has allowed the demonstration of metrological spin-squeezing (Chen *et al.*, 2011; Leroux *et al.*, 2010a; Schleier-Smith *et al.*, 2010b), squeezed-state atomic clocks (Leroux *et al.*, 2010b), and the creation of highly non-classical states (Barontini *et al.*, 2015; McConnell *et al.*, 2015).

Key theoretical proposals and experiments on atom-light interfaces for quantum metrology can be classified in three broad categories: (A) nondestructive measurements, including quantum nondemolition measurements and Zeno dynamics, Sec. V.A; (B) light-mediated coherent interactions between distant atoms, Sec. V.B; and (C) transfer of squeezing to atoms via absorption of non-classical light, Sec. V.C. Atom-light interaction currently represents the most successful method for producing large amounts of squeezing and entanglement in atomic ensembles (Cox *et al.*, 2016; Hosten *et al.*, 2016a).

A. Quantum state preparation using nondestructive measurements

1. Quantum nondemolition measurements in free space

In a quantum nondemolition (QND) measurement, a *system* (*e.g.* an atomic cloud) and *meter* (a measurement degree of freedom, *e.g.* a light beam) are coupled, such that a direct measurement of the meter provides indirect information about an observable of the system (Braginsky and Khalili, 1992; Braginsky *et al.*, 1980; Grangier




FIG. 25 **Conditional spin squeezing via QND measurements.** A two-mode light beam (blue arrow) propagates through the atomic ensemble. The light beam becomes entangled with the atomic ensemble. By measuring the light field, one gains information about the atomic state. The atomic distribution is plotted here in the $y - z$ plane. The initial coherent spin state distribution (orange circle) changes after the detection of the light beam (green ellipse). It becomes squeezed in $\hat{J}_z$ and shifted (conditioned by the measurement result), according to Eqs. (82a) and (82b).

et al., 1998). The measurement is called QND if the system and the measured value of the observable are conserved after the measurement. QND measurements are a resource for quantum metrology as they allow the preparation of entangled and spin-squeezed states of many atoms. This is well illustrated by a model Hamiltonian describing a far-off resonant dispersive interaction between the collective spin of an ensemble of atoms and a two-mode light beam in free space (Kuzmich *et al.*, 1998; Takahashi *et al.*, 1999):

$$t_p \hat{H}_{\text{QND}} = \hbar k \hat{S}_z \hat{J}_z. \quad (80)$$

Here, k is the interaction strength, t_p is the light pulse duration, and $\hat{S} = \{\hat{S}_x, \hat{S}_y, \hat{S}_z\}$ is the Stokes vector operator of the light, with components $\hat{S}_x = (\hat{a}_+^\dagger \hat{a}_- + \hat{a}_-^\dagger \hat{a}_+)/2$, $\hat{S}_y = (\hat{a}_+^\dagger \hat{a}_- - \hat{a}_-^\dagger \hat{a}_+)/2i$, and $\hat{S}_z = (\hat{a}_+^\dagger \hat{a}_+ - \hat{a}_-^\dagger \hat{a}_-)/2$. $\hat{a}_\pm$ can be two polarization modes (Hammerer *et al.*, 2004; Takahashi *et al.*, 1999), in which case Eq. (80) describes the paramagnetic Faraday rotation of light; they can also be two spatial modes of an optical Mach-Zehnder interferometer where atoms are placed in one arm and phase-shift the light (Chaudhury *et al.*, 2006; Oblak *et al.*, 2005; Windpassinger *et al.*, 2008). The interaction strength in Eq. (80) is $k \propto \frac{\sigma}{A} \frac{\Gamma}{\Delta}$, where σ is the resonant photon scattering cross section of the probe transition, Γ is the spontaneous emission rate, A is the spatial cross section of the atomic ensemble illuminated by the pulse, and Δ the detuning of the light from res-

onance (Hammerer *et al.*, 2010). The Hamiltonian (80) satisfies the back-action evasion condition $[\hat{J}_z, \hat{H}_{\text{QND}}] = 0$ such that $\hat{J}_z$ is a constant of motion. During the interaction, $\hat{S}$ precesses around the z -axis by an angle $k\hat{J}_z$. To lowest order in this angle, we have

$$\hat{S}_x^{\text{out}} \approx \hat{S}_x^{\text{in}} - k\hat{S}_y^{\text{in}}\hat{J}_z^{\text{in}}, \quad \hat{J}_x^{\text{out}} \approx \hat{J}_x^{\text{in}} - k\hat{J}_y^{\text{in}}\hat{S}_z^{\text{in}}, \quad (81a)$$

$$\hat{S}_y^{\text{out}} \approx \hat{S}_y^{\text{in}} + k\hat{S}_x^{\text{in}}\hat{J}_z^{\text{in}}, \quad \hat{J}_y^{\text{out}} \approx \hat{J}_y^{\text{in}} + k\hat{J}_x^{\text{in}}\hat{S}_z^{\text{in}}, \quad (81b)$$

$$\hat{S}_z^{\text{out}} = \hat{S}_z^{\text{in}}, \quad \hat{J}_z^{\text{out}} = \hat{J}_z^{\text{in}}. \quad (81c)$$

A measurement of $\hat{S}_x^{\text{out}}$ or $\hat{S}_y^{\text{out}}$ thus realizes a QND measurement of $\hat{J}_z$ while preserving the system's quantum coherence. As an illustration, we take atomic and optical systems both initially prepared in coherent spin states polarized along the x -axis. Assuming $\langle \hat{S}_y^{\text{in}} \rangle = 0$ and $\langle \hat{J}_z \rangle = 0$, the average phase precession is zero, $\langle \hat{S}_y^{\text{out}} \rangle = 0$, while the variance $(\Delta \hat{S}_y^{\text{out}})^2 = n(1 + \kappa^2)/4$ increases with $\kappa^2 = nNk^2/4$, where N is the number of atoms, and n the number of photons in the pulse. The mean value and variance of $\hat{J}_z$ after the measurement of the light spin $\hat{S}_y$ (with result m_y) are

$$\langle \hat{J}_z^{\text{out}} \rangle|_{m_y} = \frac{\kappa}{1 + \kappa^2} \sqrt{\frac{N}{n}} m_y, \quad (82a)$$

$$(\Delta \hat{J}_z^{\text{out}})^2|_{m_y} = \frac{1}{1 + \kappa^2} \frac{N}{4}, \quad (82b)$$

respectively,²¹ see Fig. 25. Although the final atomic state depends on the result of the measurement of the light beam (and it is thus termed conditional), the reduction in spin noise according to Eq. (82b) is completely deterministic, with no post-selection necessary. Unconditional spin squeezing can be achieved via quantum feedback that compensates, via a spin rotation, the stochastic shift of $\langle \hat{J}_z \rangle$ in Eq. (82a) due to the random measurement outcome (Berry and Sanders, 2002; Thomsen *et al.*, 2002). Cox *et al.* (2016) and Inoue *et al.* (2013) have experimentally demonstrated such unconditional spin squeezing via feedback control, see also (Vanderbruggen *et al.*, 2013). In the QND scheme, the squeezing is due to the projective measurement of the atomic state upon detection of the light, and its efficacy is determined by the performance of the detector. Yet, the above discussion is highly idealized: far-off resonant dispersive probing is unavoidably accompanied by decoherence due to Raman spin flips.²² Notice that $\kappa^2 = \eta\alpha_0$, where $\alpha_0 \propto N\sigma/A$ is the resonant optical depth and $\eta \propto n\frac{\sigma}{A}(\frac{\Gamma}{\Delta})^2$ is the average photon scattering rate per atom during the probe pulse. Scattered photons carry information about the




FIG. 26 **Spin squeezing via quantum nondemolition measurements in free space.** a) A cloud of cold Cs atoms is confined in an elongated dipole trap placed in one arm of an optical Mach-Zehnder interferometer and aligned with the largest resonant optical depth. The relative populations of the clock levels $|a\rangle$ and $|b\rangle$ are measured via the phase shift ϕ accumulated by two detuned probe beams propagating through the atomic cloud. b) Red diamonds are $\text{Var}(\Phi)$ (with $\Phi = \phi_1 - \zeta\phi_2$) obtained after two consecutive QND measurements of covariance ζ . Blue symbols are $\text{Var}(\Phi)$ obtained for a coherent spin state ($\Phi = \phi_1$, dots, and $\Phi = \phi_2$, stars). The solid line is a quadratic fit and the dashed line is the expected linear scaling with the atom number N due to projection noise. The dot-dashed line is the projection noise scaled down by the factor $(1 - \eta)^2 = 0.64$, corresponding to the reduction by the measured observed loss of atomic coherence. Different color regions are the optical shot noise (light blue), detector noise (dark blue) and projection noise (green). The inset illustrates the trade-off between spin squeezing and loss of coherence due to spontaneous emission. Adapted from Appel *et al.* (2009).

spin state of the atoms: these photons reduce the spin length as $|\langle \hat{J}_x^{\text{out}} \rangle| \approx (1 - \eta)|\langle \hat{J}_x \rangle|$ for $\eta \ll 1$ (Hammerer *et al.*, 2004), and add noise that counteracts the reduction of spin variance (de Echaniz *et al.*, 2005; Hammerer *et al.*, 2004; Madsen and Mølmer, 2004)—unless a suitable choice of atomic levels is used to avoid this effect (Chen *et al.*, 2014; Saffman *et al.*, 2009). The successful implementation of QND measurements is thus based on a proper choice of light power (increasing n reduces the photon shot noise of light detection but increases Raman scattering) and detuning [exploiting the favorable scaling of $k \propto \Gamma/\Delta$ over $\eta \propto (\Gamma/\Delta)^2$]. The most important figure of merit of an atom-light interface is the resonant optical depth α_0 , which should be much larger than unity.

Appel *et al.* (2009) reported the first experimental demonstration of metrological spin squeezing (reaching $\xi_R < 1$) via QND measurements in free space. This experiment used $N = 1.2 \times 10^5$ Cs atoms (with measured $\kappa^2 \approx 3.2$ and optical depth $\alpha_0 \approx 16$). Two equally intense and linearly polarized laser beams of different frequencies enter the arms of an optical Mach-Zehnder interferometer, see Fig. 26(a). The beams off-resonantly

²¹ These equations are obtained considering $\hat{J}_z$ and $\hat{S}_y$ to be Gaussian-distributed random variables (Hammerer *et al.*, 2010).

²² In the literature, such measurements are termed QND provided that the loss of coherence in the atomic ensemble remains small.

probe different atomic transitions and experience phase shifts proportional to the number of atoms in the probed levels (Saffman *et al.*, 2009). The detection of the relative phase shift ϕ accumulated in the optical path performs a QND measurement of the relative population in the two atomic levels (Kuzmich *et al.*, 1998; Saffman *et al.*, 2009). Spin squeezing is quantified by correlations between two consecutive QND measurements. One finds $\text{Var}(\phi_2 - \zeta\phi_1) = \frac{1}{n} + \frac{\kappa^2}{1+\kappa^2} \frac{N}{4}$, where ϕ_1 and ϕ_2 refer to the first and second phase shift detection, respectively, and the covariance $\zeta = \text{Cov}(\phi_1, \phi_2) / \text{Var}(\phi_1) = \frac{\kappa^2}{1+\kappa^2}$ expresses the correlations between the two measurements. The results reveal a spin squeezing $\xi_R^2 = -3.4$ dB, see Fig. 26(b). Furthermore, Louchet-Chauvet *et al.* (2010) have used a similar apparatus to perform a Ramsey sequence (between the two QND measurements) and demonstrated an atomic clock with a measured phase sensitivity 1.1 dB below the standard quantum limit.

Spin squeezing has also been experimentally performed by measuring the Faraday rotation of a linearly polarized probe passing through a cloud of atoms. This technique was pioneered by Kuzmich *et al.* (2000, 1999), and more recently investigated in ensembles of spin-1/2 ^{171}Yb (Inoue *et al.*, 2013; Takano *et al.*, 2009, 2010) and spin-1 ^{87}Rb atoms (Sewell *et al.*, 2012, 2014, 2013). In Sewell *et al.* (2012, 2014) spin squeezing was achieved using a two-polarization probing technique (Koschorreck *et al.*, 2010a,b). These experiments demonstrated $\xi_R^2 = -1.5$ dB (Sewell *et al.*, 2014) [$\xi_R^2 = -2$ dB inferred (Sewell *et al.*, 2012)] using $N \approx 5 \times 10^5$ atoms, mainly limited by the photon shot noise of the readout light. Polarization-based QND measurements find direct application in entanglement-assisted optical magnetometry, see Sec. VII.C.

2. Cavity-based quantum nondemolition measurements in the dispersive regime

The strength of the interaction between light and individual atoms is usually weak, but can be enhanced by placing the atoms inside an optical cavity. This method is very promising as the squeezing factor is proportional to the cavity finesse, which can be pushed to large values. Moreover, a small single-pass optical depth is advantageous for applications in atomic clocks, since it allows to reduce atomic-density-dependent atom losses, dephasing, and systematic errors.

The essential features of dispersive atom-light interaction in a cavity (Bouchoule and Mølmer, 2002a; Madsen and Mølmer, 2004; Nielsen and Mølmer, 2008; Vernac *et al.*, 2000) are captured by a simplified model comprising N three-level atoms, see Fig. 27(a). Each atom has two hyperfine levels $|a\rangle$ and $|b\rangle$ of energy difference $\hbar\omega$, and an excited state $|e\rangle$ with linewidth Γ (spontaneous decay rate into free space). The atoms are placed in




FIG. 27 **Spin squeezing via cavity-based quantum nondemolition measurements in the dispersive regime.** (a) Laser-cooled atoms are optically trapped in a standing wave (red line) inside an optical resonator. The cavity resonance is shifted in proportion to the relative population of two clock levels. The shift is measured from the transmission of a probe beam (blue line). (b) Measured spin-noise reduction $4(\Delta J_z)^2/N$ (dots) normalized to the coherent spin state projection noise. The solid line is a model fit. (c) Allan deviation of a clock that uses the generated squeezed states (black dots; the solid line indicates $9.7 \times 10^{-11} \text{ s}^{1/2}/\sqrt{\tau}$) or coherent spin states (blue circles; the dashed line is the standard quantum limit). Adapted from Hosten *et al.* (2016a).

an optical cavity with resonance frequency of ω_c and a linewidth κ_c , driven resonantly with a single-atom-single-photon effective intra-cavity Rabi frequency $2g$. The detuning of the cavity from the $|a\rangle \leftrightarrow |e\rangle$ and $|b\rangle \leftrightarrow |e\rangle$ transitions is chosen of equal magnitude $\Delta = \pm\omega/2$. Assuming homogeneous interaction (see below), low intra-cavity photon number ($n \ll \Delta^2/g^2$), and large detuning ($\Delta \gg \kappa_c, \Gamma, \sqrt{N}g$), the coupling Hamiltonian is (Schleier-Smith *et al.*, 2010a)

$$\hat{H} = \hbar\omega_c \hat{c}^\dagger \hat{c} + \hbar \frac{2g^2}{\Delta} \hat{c}^\dagger \hat{c} \hat{J}_z + \hbar\omega \hat{J}_z, \quad (83)$$

where $\hat{c}$ and $\hat{c}^\dagger$ are cavity mode operators. The effect of the light on the atoms is an ac Stark shift of the transition frequency $\delta\omega = \frac{2g^2}{\Delta} n$ between $|a\rangle$ and $|b\rangle$. Atoms in $|a\rangle$ ($|b\rangle$) increase (decrease) the index of refraction seen by the probe light, so that the net effect is a shift of the cavity resonance by $\delta\omega_c = \frac{2g^2}{\Delta} \frac{N_a - N_b}{2} = \frac{2g^2}{\Delta} J_z$, where N_a and N_b are the numbers of atoms in $|a\rangle$ and $|b\rangle$, respectively. This shift can be probed by injecting a laser into the cavity, providing a QND measurement of J_z . A detailed analysis including decoherence associated with free-space scattering of the probe light shows the possibility to achieve (Chen *et al.*, 2014; Hosten *et al.*, 2016a)

$$\xi_R^2 \approx \frac{1 + NC(\Gamma/\omega)^2}{\sqrt{NC}}, \quad (84)$$

where $C = \frac{(2g)^2}{\kappa_c \Gamma}$ is the single-atom cavity cooperativity. C is the ratio between the number of photons scattered into the cavity mode and those scattered into free space, and quantifies the optical depth of an atom with respect to the cavity mode (Tanji-Suzuki *et al.*, 2011). Note that C depends on the cavity geometry and is proportional to the cavity finesse (Tanji-Suzuki *et al.*, 2011).

The above assumption of homogeneous interaction between the trapped atoms and the cavity light field is not always valid. Inhomogeneous coupling holds when the atoms are trapped in a standing wave whose wavelength is incommensurate with that of the probe field (Leroux *et al.*, 2010b; McConnell *et al.*, 2015; Tanji-Suzuki *et al.*, 2011), when the atoms are trapped in a large volume that samples the spatial profile of the probe field (Appel *et al.*, 2009), or when the atoms move in space (Hammerer *et al.*, 2010). In these situations, Eq. (1) is modified so that each atom contributes to the collective spin with a weight given by its coupling g_i to the cavity mode: the internal-state dynamics of the effectively addressed atoms is well described by a spin operator $\hat{J}_{\text{eff}} = \sum_i (g_i/g_{\text{max}}) \hat{\sigma}_i/2$. Experimentally, this effective spin can be treated in the same way as a real spin of the same length; all the metrological methods of Sec. II remain valid. More care is required for conclusions about the correlations between the real (not effective) atoms, such as the entanglement depth (McConnell *et al.*, 2015).

Spin squeezing via cavity-based QND measurement has been first demonstrated by Schleier-Smith *et al.* (2010b), using magnetically-insensitive clock states of ^{87}Rb atoms. This experiment reached a spin squeezing $\xi_R^2 = -1.45$ dB with respect to $(\Delta\theta_{\text{SQL}})^2 = 1/N_0$, referring to $N_0 \approx 0.66N$ uncorrelated atoms (N_0 accounts for spatial variation in the atom-light coupling due to the trapping lattice being incommensurate with the cavity mode used for probing, and $N = 5 \times 10^4$). This result is mainly limited by inhomogeneous dephasing due to the cavity locking light. Hosten *et al.* (2016a) have used a cavity of higher cooperativity and exploited probing and trapping beams of commensurate frequencies, achieving a uniform atom-light coupling, see Fig. 27(a). This avoids the need for spin-echo techniques required for non-uniformly coupled systems. This experiment demonstrated $\xi_R^2 = 10.5$ dB of improved phase sensitivity with respect to $(\Delta\theta_{\text{SQL}})^2 = 1/N$, with $N = 1 \times 10^5$ ^{87}Rb atoms (and measured spin squeezing $\xi_R^2 = -18.5$ dB), see Fig. 27(b). This currently represents the highest value in expected metrological spin squeezing and measured phase sensitivity gain to date.

3. Heralded atomic entanglement created by single photon detection

In the preceding sections we have discussed the interaction of an atomic ensemble with an intense laser pulse (10^5 – 10^7 photons in typical experiments), which is subsequently detected by a standard linear photodetector that does not resolve individual photons. The light in these experiments is effectively a continuous-variable quantum system. A different approach uses weak laser pulses in combination with single-photon counting, in close analogy to the techniques developed for atomic ensemble-based quantum repeaters (Duan *et al.*, 2001). These techniques allow to create a range of highly non-classical states of a large number of atoms in a heralded way, *i.e.* conditioned on the detection of a single photon. The interaction of a linearly polarized photon with the atoms according to Eq. (80) produces a weak Faraday rotation of the photon polarization. Denoting $|\text{v}\rangle$ and $|\text{h}\rangle$ the vertical and horizontal polarizations, respectively, an atom-light system initialized in $|\text{v}\rangle \otimes |\vartheta, \varphi, N\rangle$ [where $|\vartheta, \varphi, N\rangle$ is a coherent spin state, see Sec. II.C] evolves by the QND interaction (80) to the state (McConnell *et al.*, 2013)

$$|\psi\rangle = \frac{1}{2}|\text{v}\rangle \otimes (|\vartheta, \varphi_+, N\rangle + |\vartheta, \varphi_-, N\rangle) + \frac{1}{2}|\text{h}\rangle \otimes (|\vartheta, \varphi_+, N\rangle - |\vartheta, \varphi_-, N\rangle), \quad (85)$$

where $\varphi_{\pm} = \varphi \pm \phi$ and ϕ is the phase accumulated during the interaction. The detection of a vertically polarized photon projects the atomic ensemble to $|\psi_{\text{v}}\rangle \propto |\vartheta, \varphi_+, N\rangle + |\vartheta, \varphi_-, N\rangle$, whereas the detection of a horizontally polarized photon prepares $|\psi_{\text{h}}\rangle \propto |\vartheta, \varphi_+, N\rangle - |\vartheta, \varphi_-, N\rangle$. The entanglement properties strongly depend on the phase ϕ , and the quantum Fisher information is (without loss of generality we assume here that $\vartheta = \pi/2$ and $\varphi = 0$)

$$\frac{F_Q[|\psi_{\text{h,v}}\rangle, \hat{J}_y]}{N} = \frac{1 + (N-1)\sin^2\phi \pm \cos^N\phi}{1 \pm \cos^N\phi}, \quad (86)$$

where the plus (minus) sign holds for the detection of a vertically (horizontally) polarized photon. To leading order in N , we have $F_Q[|\psi_{\text{v}}\rangle, \hat{J}_y]/N = 1 + N\phi^2/2 + \mathcal{O}(N^2\phi^4)$ and $F_Q[|\psi_{\text{h}}\rangle, \hat{J}_y]/N = 3 + N\phi^2/2 + \mathcal{O}(N^2\phi^4)$: for $|\phi| \lesssim 2/\sqrt{N}$, the state $|\psi_{\text{h}}\rangle$ is metrologically more useful than $|\psi_{\text{v}}\rangle$. In particular, for $|\phi| \ll 1/\sqrt{N}$, the (rare) detection of a horizontally polarized photon heralds the generation of $|\psi_{\text{h}}\rangle = |(N/2 - 1)_x\rangle$, which is the one-excitation Dicke state. For $|\phi| \gtrsim 2/\sqrt{N}$, we have $\cos^N\phi \approx 0$ in Eq. (86) and the two states give approximately the same quantum Fisher information. At $\phi = \pm\pi/2$, we get the NOON state $|\psi_{\text{h,v}}\rangle = (|(N/2)_y\rangle \pm |(-N/2)_y\rangle)/\sqrt{2}$. In the realistic scenario $|\phi| \ll 1/\sqrt{N}$, the detection of a sequence of photons conditionally prepares more and more entanglement. For instance, the detection




FIG. 28 **Entangled states generated by the detection of a single photon.** (a) An atomic ensemble confined in a cavity interacts with vertically polarized light. The detection of an outgoing horizontally polarized single photon projects the atoms into a state with a negative Wigner function. (b) Experimental Wigner function of the heralded atomic state with effective atom number $N = 2100$ (see Sec. V.A.2), obtained from the reconstructed atomic density matrix, and showing negative parts. The dashed line is the contour at which the Wigner function of an N -atom coherent spin state is equal to $1/\sqrt{e}$ of its maximum value. Adapted from McConnell *et al.* (2015).

of n_h h-polarized photons prepares the NOON state $(|(N/2)_z\rangle + |(-N/2)_z\rangle)/\sqrt{2}$ when $n_h \times |\phi| \gtrsim \sqrt{N}$. Finally, Chen *et al.* (2015) have generalized the above method showing the possibility to generate a broad class of entangled states of many atoms using single photon having a tailored frequency spectrum and time-resolved detection.

Experimentally, the use of single-photon counting for preparing entangled states of an atomic ensemble has been pioneered by Christensen *et al.* (2014, 2013). These experiments have shown the expected increase of spin fluctuations due to the single-photon detection. McConnell *et al.* (2015) have demonstrated the generation of entanglement in an atomic ensemble of $N_a = 3100$ laser-cooled ^{87}Rb atoms prepared in an optical cavity and probed with an off-resonant weak pulse (of about 200 photons) of vertically polarized light, see Fig. 28. The accumulated phase is $\phi = \frac{4g^2}{\Delta\kappa_c}$, where $2/\kappa_c$ is the characteristic atom-photon interaction time. For the experiment of McConnell *et al.* (2015) we have $\phi\sqrt{N} \approx 0.03$. This experiment has investigated the entangled state produced by the detection of a horizontally polarized photon, while the detection of vertically polarized photons gives, for these experimental parameters, only a slight spin squeezing. From the tomographic reconstructed density matrix it is possible to obtain a Wigner distribution with negative areas, see Fig. 28, and an entanglement depth (not related to metrological usefulness) of 2900 atoms. This work demonstrates how the information carried by a single (or a few) photon(s) can create entanglement in a large atomic ensemble. In Hu *et al.* (2016) the same authors have re-analyzed these data to show that the gener-

ated atomic state violates classical physics even if no assumptions related to quantum mechanics are made. This conclusion (Kot *et al.*, 2012), similarly to Bell inequalities (Brunner *et al.*, 2014), observes that marginal probability distributions measured for non-commuting observables do not always come from a joint probability distribution identified with a classical state. As in Schmied *et al.* (2016) (see Fig. 17), Hu *et al.* (2016) show that classical physics is insufficient for describing mesoscopic entangled states of atoms.

4. Cavity-based quantum nondemolition measurements in the normal mode splitting regime

The experiments discussed in the previous sections operate in the dispersive regime, where the cavity is far detuned from atomic resonance. In this section, we discuss quantum state preparation schemes where the cavity is tuned near an atomic resonance with respect to the atomic transition $|b\rangle \leftrightarrow |e\rangle$, see Fig. 29(a). Atoms in level $|a\rangle$ are assumed not to interact with the cavity mode. In the limit $N_b \gg 1$, and assuming that the system is driven weakly so that the mean number of atoms in $|e\rangle$ is small with respect to N_b , atoms in $|b\rangle$ cause a splitting $\omega_c \rightarrow \omega_c^\pm$ of the cavity resonance frequency (Chen *et al.*, 2014). We have

$$\omega_c^\pm = \omega_c - \frac{\delta_c \pm \sqrt{\delta_c^2 + 4g^2 N_b}}{2}, \quad (87)$$

where $\delta_c = \omega_c - \omega_{be}$ is the cavity detuning from the atomic transition ω_{be} , ω_c is the bare cavity frequency, N_b is the number of atoms in $|b\rangle$, and g is the single-atom-single-photon coupling rate (uniform coupling is assumed). At $\delta_c = 0$, the appearance of two well-resolved resonances separated by $2g\sqrt{N_b}$ is referred to as collective vacuum Rabi splitting. In the limit of collective strong coupling $g\sqrt{N_b} \gg \kappa_c, \Gamma$ the normal modes of the coupled system are well resolved: a measurement of $\omega_c^\pm$ thus allows to determine N_b .

In the experiments of Bohnet *et al.* (2014); Chen *et al.* (2011); and Cox *et al.* (2016), the atom number N_b is measured non-destructively by probing the cavity resonant frequency with an additional probe laser injected into the cavity. The number of particles in level $|a\rangle$ can be probed by first applying a resonant π pulse between $|a\rangle$ and $|b\rangle$ and then repeating the QND measurement. A detailed analysis of the impact of decoherence in this system (mainly due to scattering of cavity photons into free space) can be found in Chen *et al.* (2014). In (Chen *et al.*, 2011) the optical cavity is tuned to resonance, $\delta_c = 0$, with $NC \approx 1400$. Chen *et al.* (2011) reported a spin squeezing $\xi_R^2 = -1$ dB ($\xi_R^2 = -3.3$ dB inferred) relative to the standard quantum limit using $N_0 = 7 \times 10^5$ ^{87}Rb atoms (N_0 is an effective atom number taking into account inhomogeneous coupling in the cavity). In (Bohnet




FIG. 29 **Spin squeezing via quantum nondemolition measurements in the normal mode splitting regime.** (a) Energy-level diagram showing the cavity shift due to atoms in level $|b\rangle$. (b) ξ_R^2/N as a function of atom number N for uncorrelated particles ($\xi_R \geq 1$, black dots) and spin-squeezed states ($\xi_R < 1$, red dots). The red data follow a $1/N^2$ scaling (as shown also in the inset, the black line being a fit to the data). Adapted from [Bohnet et al. \(2014\)](#).

[et al., 2014](#); [Cox et al., 2016](#)) noise effects were considerably reduced taking advantage of a cycling transition ([Chen et al., 2014](#); [Saffman et al., 2009](#)), and higher collective cooperativity $NC \approx 6000$ was reached. Experimental results demonstrate $\xi_R^2 = -10.1$ dB ([Bohnet et al., 2014](#)), see Fig. 29, and $\xi_R^2 = -17.7$ dB ([Cox et al., 2016](#)), using $N_0 = 5 \times 10^5$ ^{87}Rb atoms. The achieved spin squeezing witnesses the presence of useful 170 ± 30 particle entanglement ([Cox et al., 2016](#)). This squeezing technique has been further combined with quantum feedback, reaching $\xi_R^2 = -7.4$ dB of deterministic spin squeezing ([Cox et al., 2016](#)).

Resonant atom-light coupling in an optical cavity also offers the possibility to create entangled states beyond spin squeezing. [Haas et al. \(2014\)](#) have used an atom chip integrated with a high-finesse fiber-optical cavity in the strong-coupling regime ([Volz et al., 2011](#)) to prepare a generalized W state ([Dür et al., 2000](#)) of $N \approx 40$ cold ^{87}Rb atoms. In this experiment the cavity and probe laser tuned on resonance with the $|b\rangle \leftrightarrow |e\rangle$ transition. The single-atom strong coupling $g \gg \kappa_c, \Gamma$ guarantees that the cavity only transmits light if all N atoms are in level $|a\rangle$: a single atom in $|b\rangle$ is sufficient to block the cavity transmission. The experiment of [Haas et al. \(2014\)](#) starts with all atoms in level $|a\rangle$, and a weak microwave pulse coupling $|a\rangle$ and $|b\rangle$ prepares the state $\sqrt{1-p}|N\rangle_a|0\rangle_b + \sqrt{p}|N-1\rangle_a|1\rangle_b$, where $p \ll 1$ to avoid multi-particle excitations. The cavity transmission is then measured with a probe-light beam. A low transmission heralds the QND preparation of the W state $|N-1\rangle_a|1\rangle_b$. The resulting state is characterized by state tomography, from which it is possible to extract a lower




FIG. 30 **State engineering via quantum Zeno dynamics.** (a) The $|a\rangle$ and $|b\rangle$ levels are coupled by a microwave field at Rabi frequency Ω , while the optical cavity transition $|b\rangle \leftrightarrow |e\rangle$ is probed with a coupling rate much faster than Ω . (b) Starting with all atoms in $|b\rangle$, microwave Rabi coupling and simultaneous measurement of the cavity transmission leads to a coherent evolution restricted to the subspace highlighted by the orange shaded area. The preparation of the state with no atoms in $|b\rangle$ (green shaded area) is forbidden by the quantum Zeno effect. The upper row shows the Husimi distributions of symmetric Dicke states. (c) Experimental Husimi distributions, measured directly (upper row) and obtained from the reconstructed density matrix (lower row), for different evolution times t/T , where $T = \pi/\Omega$ is the time for a Rabi π -pulse in the absence of cavity probing. Taken from [Barontini et al. \(2015\)](#).

bound on the entanglement depth of 13 atoms, at least.

5. Quantum state preparation via quantum Zeno dynamics

Quantum Zeno dynamics combines the coherent evolution of a system with the measurement of an initially unoccupied quantum state ([Facchi and Pascazio, 2008](#)): if the measurement is performed with high enough frequency, the measured state remains unoccupied. This profoundly modifies the system dynamics and allows engineering specific entangled states. In [Barontini et al. \(2015\)](#) a cavity-based measurement in the regime $C \gg 1$ ([Volz et al., 2011](#)) is used to prepare a generalized W state of 36 ^{87}Rb atoms via quantum Zeno dynamics, see Fig. 30. The N atoms are all initialized in level $|b\rangle$ and coupled to the $|a\rangle$ level by a microwave Rabi drive at Rabi frequency Ω . The cavity is in resonance with the $|b\rangle \leftrightarrow |e\rangle$ transition and transmits only if there are no atoms in level $|b\rangle$. The cavity transmission is probed at a rate much larger than Ω . When no measurement is performed, the microwave Rabi coupling prepares the state $|N\rangle_a|0\rangle_b$ after a time $T = \pi/\Omega$. In presence of the cavity measurement, the quantum Zeno effect forbids the preparation of $|N\rangle_a|0\rangle_b$ and deterministically prepares the W

state $|N-1\rangle_a|1\rangle_b$, see Fig 30. From the density matrix extracted via state tomography, it is possible to obtain the quantum Fisher information. Values of the quantum-Fisher information up to $F_Q/N = 1.51$ are reported by Barontini *et al.* (2015).

B. Light-mediated coherent interaction between distant atoms

In contrast to the measurement-based scheme of Sec. V.A, off-resonant atom-light coupling can also be used to realize a light-mediated coherent interaction between distant atoms. In situations where the light has a Stokes component $\hat{S}_z \propto \hat{J}_z$, Eq. (80) reduces to $\hat{H}_{\text{QND}} = \hbar k \hat{J}_z^2$ (Agarwal *et al.*, 1997; Zhang *et al.*, 2003), corresponding to an effective one-axis twisting that generates unconditional spin squeezing (Kitagawa and Ueda, 1993). As a main difference with respect to contact interaction discussed in Sec. III.B, this interaction is mediated by light and thus long-range, so that it can be applied to a dilute sample, which is preferred for minimizing systematic errors in precision measurements. Moreover, in contrast to spin squeezing obtained from QND measurements, it deterministically produces known entangled states, independently from the detector performance.

There are different proposals for the realization of an effective interaction between distant atoms. Takeuchi *et al.* (2005) have studied a double-pass Faraday interaction in free space where a polarized light beam passes twice through an atomic ensemble (after the first passage the polarization is properly rotated such that $\hat{S}_z \propto \hat{J}_z$ at the second passage). An improved version of this proposal has been discussed by Trail *et al.* (2010) using a quantum eraser to remove residual spin-probe entanglement. Schleier-Smith *et al.* (2010a) and Zhang *et al.* (2015) have proposed the realization of light-induced atom-atom interaction in an optical cavity (also called cavity squeezing). As shown by Eq. (83) the atoms modify the cavity resonance frequency by an amount proportional to $\hat{J}_z$. When the cavity is driven by a probe laser tuned to the slope of the cavity resonance, the intracavity photon number $\hat{c}^\dagger \hat{c}$ in Eq. (83), being detuning-dependent, changes linearly with the index of refraction of the atomic cloud (which, in turn, is proportional to $\hat{J}_z$). Notice that tracing out the light field (which carries information about the atomic spin) yields a dissipative dynamics that limits the attainable squeezing (Leroux *et al.*, 2012). Finally, an alternative proposal to realize an effective one-axis twisting atomic Hamiltonian uses a four-wave Raman scheme to directly drive the atomic ensemble, where atoms communicate with one another by exchanging photons via a weakly-occupied cavity mode (Sørensen, 2002).

One axis twisting by light-mediated coherent interaction has been experimentally implemented in an optical




FIG. 31 **Squeezing via light-mediated effective interaction between atoms.** Effective spin squeezing $\xi_{\text{eff}}^2 = \sigma^2(Q)C(0)/C(Q)^2$ giving the expected gain with respect to the sensitivity experimentally achieved in absence of entanglement, as a function of the shearing strength Q (proportional to the photon number in the cavity). Here $\sigma^2(Q) = (\Delta \hat{J}_z)^2/(N_0/4)$, and $C(Q)$ is the contrast of Rabi oscillations of the sheared state, shown in inset. The effective spin squeezing is related to the metrological spin squeezing Eq. (28) as $\xi_{\text{eff}}^2 = C(0)\xi_R^2$. An effective squeezing $\xi_{\text{eff}}^2 = -5.6$ dB is reached, corresponding to $\xi_R^2 = -4.6$ dB. Adapted from Leroux *et al.* (2010a).

cavity (Leroux *et al.*, 2010a,b), using an effective number $N_0 = 3 \times 10^4$ of ^{87}Rb atoms. The experiment implements the scheme of Schleier-Smith *et al.* (2010a). The squeezed state is characterized via spin-noise tomography and the contrast of Rabi oscillations. The results of Leroux *et al.* (2010a) are summarized in Fig. (31) and demonstrate a gain $\xi_R^2 = -4.6$ dB over $(\Delta\theta_{\text{SQL}})^2 = 1/N_0$ ($\xi_{\text{eff}}^2 = -5.6$ dB over the sensitivity experimentally expected in absence of entanglement). In Leroux *et al.* (2010b) the squeezed states generated with this method have been used to realize an atomic clock with sensitivity 3.5 dB below the standard quantum limit, see Sec. VII.

C. Quantum state transfer from non-classical light to atoms

Spin-squeezed states can be created via atom-light interaction by transferring quadrature squeezing of light to atomic spin squeezing (Agarwal and Puri, 1990; Kuzmich *et al.*, 1997). This effect can be understood from the Jaynes-Cummings model Hamiltonian

$$\hat{H}_{\text{JC}} = \hbar\Omega(\hat{c}\hat{J}_+ + \hat{c}^\dagger\hat{J}_-), \quad (88)$$

describing the interaction of nondecaying two-level atoms with a light mode. Here $\hat{c}^\dagger$ and $\hat{c}$ are the light creation and annihilation operators, respectively, $\hat{J}_\pm$ are spin ris-

ing and lowering operators, see Sec. II.A, and Ω is a coupling constant. The Heisenberg equations of motion for $\hat{c}$ and $\hat{J}_\pm$ can be found analytically within a frozen-spin approximation, assuming $\hat{J}_z$ constant and equal to $N/2$. We obtain (Wineland *et al.*, 1994, 1992)

$$\xi_R^2(t) = \xi_R^2(0) \cos^2(\Omega_N t) + \xi_p^2(0) \sin^2(\Omega_N t) \quad (89a)$$

$$\xi_p^2(t) = \xi_p^2(0) \cos^2(\Omega_N t) + \xi_R^2(0) \sin^2(\Omega_N t), \quad (89b)$$

where $\xi_p^2 = 2(\Delta\hat{P}_c)^2$, $\hat{P}_c = \frac{\hat{c}-\hat{c}^\dagger}{i\sqrt{2}}$, $\xi_R^2 = \frac{N(\Delta\hat{J}_x)^2}{\langle\hat{J}_z\rangle^2}$, and $\Omega_N = k\sqrt{N}$. Similar expressions can be obtained relating $\xi_R^2 = \frac{N(\Delta\hat{J}_y)^2}{\langle\hat{J}_z\rangle^2}$ with $\xi_x^2 = 2(\Delta\hat{X}_c)^2$, with $\hat{X}_c = \frac{\hat{c}+\hat{c}^\dagger}{\sqrt{2}}$. These equations predict that atomic spin squeezing $\xi_R < 1$ can be achieved by first squeezing the light field, $\xi_p < 1$, and then transferring this quadrature squeezing onto the spins. Optimal spin squeezing is reached at $t = \pi/(2\Omega_N)$. For strong squeezing the contrast $\langle\hat{J}_z\rangle$ decreases, and the frozen-spin approximation breaks down. Since the squeezed light is the source of spin squeezing in this method, the degree of squeezing is determined by the quality of the quadrature squeezed light.

Quantum state transfer from light to atoms has been first experimentally demonstrated by (Hald *et al.*, 1999) using a cloud of 10^7 cold Cs atoms, following the theoretical proposal of Kuzmich *et al.* (1997). This experiment used a V-level scheme consisting of three atomic hyperfine levels (Hald *et al.*, 1999, 2000). The reduction of atomic spin noise below the projection noise is generated by the absorption of polarized coherent and squeezed vacuum light with opposite circular polarizations. The atomic fluctuations are read out by the differential photocurrent of a probe beam at the output ports of a polarizing beam splitter.

An alternative approach, mapping a quantum state of light onto an atomic state via electromagnetically induced transparency has been proposed by Fleischhauer and Lukin (2000), see also (Liu *et al.*, 2001; Phillips *et al.*, 2001)

VI. ENTANGLED STATES OF TRAPPED IONS

Ensembles of trapped ions confined in electromagnetic traps and manipulated with laser beams are one of the most successful systems to generate and exploit entanglement (Blatt and Wineland, 2008; Wineland, 2013). Direct applications include quantum information and computation (Häffner *et al.*, 2008; Roos, 2014), quantum networks (Duan and Monroe, 2010), quantum simulations (Blatt and Roos, 2012; Martinez *et al.*, 2016; Schneider *et al.*, 2012) and, as reviewed here, quantum metrology. Trapped ions are well isolated from the environment, can be coherently manipulated, individually addressed and detected with almost unit efficiency (Leibfried *et al.*,

2003a). Turchette *et al.* (1998) reported the first deterministic generation of entanglement of two trapped ions. This experiment was readily followed by the demonstration of a phase sensitivity below the standard quantum limit (Sackett *et al.*, 2000) and the realization of an atomic clock exploiting entangled ions (Meyer *et al.*, 2001).

Typical experiments (Leibfried *et al.*, 2003a) confine ions in a linear Paul trap [Fig. 32(a)], surface-electrode Paul trap (Mielenz *et al.*, 2016; Sterling *et al.*, 2014), or Penning trap [Fig. 32(b)], followed by ground-state cooling. So far, up to 20 ions have been entangled in a Paul trap (Monz *et al.*, 2011) and up to 220 ions in a Penning trap (Bohnet *et al.*, 2016). One- and two-dimensional ion arrangements are well suited for individually controlling and reading out internal states of the ions. We note that individual particle manipulation and detection are crucial for many quantum-computation and quantum-information tasks, but they are not necessary for quantum-enhanced metrology. Interesting metrological experiments can thus be also performed in Penning traps (where local manipulation is currently not available), which are well suited to store, manipulate, and entangle large numbers of ions (Bohnet *et al.*, 2016; Britton *et al.*, 2012).

In trapped-ion experiments, the qubit is formed by two internal (usually hyperfine) states $|a\rangle$ and $|b\rangle$ of each ion. This internal qubit can be coupled by an electromagnetic field to external (motional, vibrational) degrees of freedom, see Fig. 32(c). A key aspect is the local detection of the internal ion's state with extremely high efficiency. This is usually achieved by resonant fluorescence. Detection errors arise if the qubit changes its state during the detection interval or if photons scattered off trap electrodes are mistaken for fluorescence photons. In Harty *et al.* (2014) and Myerson *et al.* (2008) qubit detection errors smaller than 10^{-4} are reported.

A. Generation of metrologically useful entangled states of many ions

The generation of entanglement between different qubits is one major challenge in quantum information processing with trapped ions (Wineland *et al.*, 1998). The Coulomb force pushes the ions apart to a distance much larger than the Bohr radius, and thus prevents state-dependent ion-ion interactions in absence of external electromagnetic fields. However, as a consequence of the Coulomb force, the normal modes of motion in the trap are shared among all ions and can provide a means for entangling the qubits. Most of the schemes for entangling trapped ion qubits are based on internal-state-dependent forces acting on the ions, generated by external electromagnetic fields. These forces can be arranged in time and frequency in such a way that an effective




FIG. 32 **Common ion-trapping setups.** a) Linear Paul trap using radio-frequency oscillating electric fields to trap ions, combined with static electric control fields. In tight radial confinement, laser-cooled ions form a linear string (see inset image for 8 ions) with a spacing determined by the balance between external confining fields and ion-ion Coulomb repulsion. Taken from Blatt and Wineland (2008). b) Penning trap. Ions are confined in the xy plane with a strong homogeneous magnetic field and axially by a quadrupole electric field. The green arrows indicate bi-chromatic light fields used to generate entanglement. The inset shows an image of a 2D crystal of 91 ions. Taken from Bohnet *et al.* (2016). c) Each trapped ion can be modeled as an effective internal two-level system— $|a\rangle$ and $|b\rangle$ interacting with a radiation characterized by Rabi frequency Ω and decay rate γ —and a shared external harmonic oscillator potential with equally spaced energy levels of mode frequency ω_m .

entangling operation is carried out between two or more qubits and no entanglement persists between the qubits and the vibrational degrees of freedom.

The entangling gates can be broadly classified in three categories, according to the way the electromagnetic fields interact with the ions: *i*) Quantum gates induced by a laser beam that interacts with a single ion at a time as originally proposed by Cirac and Zoller (1995). The ion is entangled with a vibrational mode of the ion string and the entanglement is subsequently transferred from the vibrational mode to the internal state of a second ion. *ii*) Quantum gates induced by an electromagnetic field that simultaneously addresses two or more ions, either through optical ac Stark shifts (García-Ripoll *et al.*, 2005; Kim *et al.*, 2009, 2010; Lee *et al.*, 2005; Leibfried *et al.*, 2003b; Milburn *et al.*, 2000; Mølmer and Sørensen, 1999; Solano *et al.*, 1999; Sørensen and Mølmer, 1999; Sørensen and Mølmer, 2000), through static magnetic-field gradients in combination with homogeneous radio-frequency fields (Johanning *et al.*, 2009a,b; Mintert and Wunderlich, 2001; Welzel *et al.*, 2011), or with radio-frequency field gradients (Ospelkaus *et al.*, 2008, 2011). As an important advantage, these gates do not require to cool all ion normal modes to their ground states, but only the ion motion to be well within the Lamb-Dicke

regime $\langle \hat{z}^2 \rangle \ll (\lambda/2\pi)^2$, where z is the displacement of an ion along the laser propagation direction and λ the laser wavelength. The basic idea is to drive a phonon (normal) mode of frequency ν by a periodic state-dependent force with frequency ω . In the rotating-wave approximation, the Hamiltonian describing this conditional interaction is (Roos, 2014)

$$\hat{H} = i\hbar k(\hat{c}^\dagger e^{i\delta t} - \hat{c} e^{-i\delta t})\hat{O}, \quad (90)$$

where $\hat{O}$ is an operator acting on the (internal) qubit states, $\hat{c}^\dagger$ and $\hat{c}$ are the harmonic oscillator creation and annihilation operators of the addressed normal mode, respectively, k is a coupling strength, and $\delta = \omega - \nu$. The propagator $\hat{U}$ corresponding to this (time-dependent) Hamiltonian is

$$\hat{U}(t) = \exp[\alpha(t)\hat{O}\hat{c}^\dagger - \alpha^*(t)\hat{O}^\dagger\hat{c}] \exp[i\phi(t)\hat{O}^2], \quad (91)$$

where $\alpha(t) = i(k/\delta)(1 - e^{i\delta t})$ and $\phi(t) = (k/\delta)^2(\delta t - \sin \delta t)$. If the interaction time $t = \tau$ is chosen such that $\alpha(\tau) = 0$, then the propagator reduces to $U(\tau) = \exp[i\phi(\tau)\hat{O}^2]$. A vibrational mode becomes transiently entangled with the qubits before getting disentangled at the end of the gate operation, resulting in an effective nonlocal operation capable of entangling the ions. The conditional phase gate (Leibfried *et al.*, 2003b) is obtained by setting $\hat{O} = \frac{1}{2} \sum_{i=1}^N \hat{\sigma}_z^{(i)} = \hat{J}_z$ in Eq. (91), and is realized by placing the ions in a qubit-state-dependent potential created by two counter-propagating laser beams. The Mølmer-Sørensen gate (Mølmer and Sørensen, 1999; Sørensen and Mølmer, 2000) is obtained setting $\hat{O} = \frac{1}{2} \sum_{i=1}^N \hat{\sigma}_y^{(i)} = \hat{J}_y$ (or $\hat{O} = \hat{J}_x$). This gate is realized by a bichromatic laser field tuned close to the upper and lower motional sideband of the qubit transition (Benhelm *et al.*, 2008; Sackett *et al.*, 2000). The propagator is thus of the kind $\hat{U} = \exp[i\phi(\tau)\hat{J}_n^2]$, formally equivalent to Kitagawa-Ueda one-axis twisting, see Sec. III.B. Spin-dependent optical dipole forces generated by a pair of off-resonance laser beams with different frequencies, addressing many normal modes simultaneously, have been recently used to engineer tunable Ising spin-spin coupling (Kim *et al.*, 2009; Lee *et al.*, 2005; Porras and Cirac, 2006). The resulting propagator is (Porras and Cirac, 2006)

$$\hat{U}(t) = \exp \left[i \sum_{i,j=1}^N V_{i,j} \hat{\sigma}_z^{(i)} \hat{\sigma}_z^{(j)} \right], \quad (92)$$

where $V_{ij} \propto d_{ij}^{-\alpha}$, d_{ij} is the distance between the i th and j th ions and α can be tuned by adjusting the laser frequencies in the range $0 \leq \alpha \leq 3$ (Britton *et al.*, 2012; Schmied *et al.*, 2011). Ising spin-spin couplings have been implemented in a linear Paul trap with up to ~ 20 ions (Islam *et al.*, 2013; Jurcevic *et al.*, 2014; Richerme *et al.*, 2014) and in a Penning trap forming a 2D Coulomb crystal of ~ 200 ions (Bohnet *et al.*, 2016; Britton *et al.*, 2012),

see Fig. 32(b). *iii*) A third way to generate ion entanglement is based on performing joint measurements on photons that are first entangled with ion qubits (Casabone *et al.*, 2013; Moehring *et al.*, 2007; Stute *et al.*, 2012). This scheme enables the generation of entanglement between ions separated by large distances and does not require the ions to be in the Lamb-Dicke regime.

B. Quantum metrology with trapped ions

1. Quantum metrology with two ions

Historically, the first atomic phase sensitivity below the standard quantum limit was demonstrated with two trapped ions (Meyer *et al.*, 2001; Sackett *et al.*, 2000). Meyer *et al.* (2001) reported the creation of the state

$$|\psi\rangle = \cos\beta|a\rangle^{\otimes 2} + i\sin\beta|b\rangle^{\otimes 2} \quad (93)$$

of two $^9\text{Be}^+$ ions via a Mølmer-Sørensen gate in a linear Paul trap. Here β is a tunable parameter proportional to the laser pulse duration. The state (93) is spin-squeezed, $\xi_R^2 = \frac{N(\Delta\hat{J}_\perp)^2}{\langle\hat{J}_z\rangle^2} = \frac{1-\sin 2\beta}{\cos^2 2\beta} < 1$ for $0 < \beta < \pi/2$. It reaches the Heisenberg limit $\xi_R^2 = 1/2$ at $\beta = \pi/4$, corresponding to the creation of a maximally entangled state. However, at $\beta = \pi/4$, $\langle\hat{J}_z\rangle = 0$ and any technical noise prevents detecting the maximally entangled state as spin squeezed. The experiment of Meyer *et al.* (2001) reached $\xi_R^2 = 0.85$ at the experimentally optimal working point $\beta = \pi/10$. A phase estimation with sensitivity below $\Delta\theta_{\text{SQL}}$ with the maximally entangled state ($\beta = \pi/4$) was demonstrated by applying a spin rotation $\exp(-i\theta\hat{J}_z)$ followed by $\exp(-i\frac{\pi}{2}\hat{J}_x)$. Measuring the parity $\hat{\Pi} = \hat{\sigma}_z^{(1)}\hat{\sigma}_z^{(2)}$ led to a sensitivity $(\Delta\theta)^2/(\Delta\theta_{\text{SQL}})^2 = 0.70$, where $\Delta\theta = \Delta\hat{\Pi}/|d\hat{\Pi}/d\theta|$. Other experiments with two $^9\text{Be}^+$ ions (Leibfried *et al.*, 2003b; Sackett *et al.*, 2000), two $^{111}\text{Cd}^+$ ions in magnetic-field-insensitive clock states (Haljan *et al.*, 2005), and two $^{40}\text{Ca}^+$ ions using an optical transition (Home *et al.*, 2006; Monz *et al.*, 2011; Noguchi *et al.*, 2012) have reported parity oscillations from which it is possible to extract the phase sensitivity. In particular, Leibfried *et al.* (2003b) demonstrated $(\Delta\theta)^2/(\Delta\theta_{\text{SQL}})^2 = 0.55$, very close to the Heisenberg limit $(\Delta\theta)^2/(\Delta\theta_{\text{SQL}})^2 = 0.5$, with a maximally entangled states obtained via a geometric phase gate. Finally, Meyer *et al.* (2001) also reported a full Ramsey sequence, where the maximally entangled state is generated between two Ramsey pulses and the phase θ is proportional to the interrogation time. In this case, the initial probe state is a symmetric Dicke (or twin-Fock) state. By measuring the parity, it was possible to reach a spectroscopic sensitivity $(\Delta\theta)^2/(\Delta\theta_{\text{SQL}})^2 = 0.77$. The creation of symmetric Dicke states of two and four $^{40}\text{Ca}^+$ ions is reported in Noguchi *et al.* (2012).




FIG. 33 **Phase sensitivity of ion Schrödinger cat states.** a) Typical parity oscillations obtained with cat states. These have a characteristic period $2\pi/N$ (here for $N = 8$). Adapted from Monz *et al.* (2011). b) Summary of the experimental achievements (symbols). Here we show the Fisher information as a function of the number of qubits N , $F = V^2 N^2$, obtained from the extracted experimental visibilities V . The upper thick line is the Heisenberg limit $F = N^2$, the lower thick line is the standard quantum limit, $F = N$. The different lines are bounds for useful k -particle entanglement, Eq. (25). In particular, the darker red region stands for useful genuine N -particle entanglement. The inset is a zoom for $N = 2$ ions. Adapted from (Pezzè *et al.*, 2015).

2. Quantum metrology with GHZ states of several ions

The maximally entangled (or “Schrödinger cat”) state

$$|\psi_{N,\text{cat}}\rangle = \frac{|a\rangle^{\otimes N} + |b\rangle^{\otimes N}}{\sqrt{2}} \quad (94)$$

has been created with up to $N = 6$ $^9\text{Be}^+$ ions (Leibfried *et al.*, 2005) and up to $N = 14$ $^{40}\text{Ca}^+$ ions (Monz *et al.*, 2011) in a linear Paul trap. This state is formally obtained from $\hat{U}_{\text{cat}}|b\rangle^{\otimes N}$, where $\hat{U}_{\text{cat}} = \exp(-i\frac{N\pi}{2}\hat{J}_x)\exp(i\frac{\pi}{2}\hat{J}_z^2)$, see Sec. III.B. When applying a spin rotation $\exp(-i\theta\hat{J}_z)$, the state (94) transforms as $|\psi_{N,\text{cat}}(\theta)\rangle = [\exp(i\theta N/2)|a\rangle^{\otimes N} + \exp(-i\theta N/2)|b\rangle^{\otimes N}]/\sqrt{2}$, such that the relative phase between $|a\rangle^{\otimes N}$ and $|b\rangle^{\otimes N}$ is $N\theta$. The state is de-




FIG. 34 **Spin squeezing of hundreds of trapped ions.** Metrological spin-squeezing parameter as a function of the number of ions N . The black points are results obtained for a coherent spin state. The solid purple squares show ξ_R^2 (optimized over probe preparation time) without subtractions of detection noise. The open squares show ξ_R^2 after subtracting photon shot noise. The dashed (solid) line is a theoretical prediction (including spontaneous emission). Taken from [Bohnet et al. \(2016\)](#).

tected after a $\pi/2$ Rabi rotation around the x -axis. The probability to obtain N_a atoms in level a and $N_b = N - N_a$ atoms in level b is $P(N_a|N, \theta) = 2^{-N} \binom{N}{N_a} \{1 + (-1)^{N_a} \cos[N(\theta + \pi/2)]\}$. These probabilities oscillate with a frequency N times faster than that of a single qubit, and therefore most observable properties of these states, such as the occurrence of N atoms in level a or b , or collective properties such as the parity of the relative number of particles in the output state ([Bollinger et al., 1996](#)), feature this rapid oscillation. Notably, applying the nonlinear transformation $\hat{U}_{\text{cat}}$ again after phase encoding, *i.e.* $\hat{U}_{\text{cat}} \exp(-i\theta \hat{J}_z) \hat{U}_{\text{cat}} |b\rangle^{\otimes N} = \cos(\frac{N\theta}{2} - \frac{\pi}{4}) |b\rangle^{\otimes N} + i \cos(\frac{N\theta}{2} + \frac{\pi}{4}) |a\rangle^{\otimes N}$, one obtains only two detection events (N atoms in level a or b) that can be experimentally distinguished with high efficiency ([Leibfried et al., 2004, 2005](#)). Experiments exploring quantum metrology with states (94) generally report dichotomic events—the parity of the relative atom number, in [Leibfried et al. \(2004\)](#) and [Monz et al. \(2011\)](#), or the detection of all atoms in the same level in [Leibfried et al. \(2004, 2005\)](#). From the reported probabilities, $P(\pm|\theta) = \frac{1 \pm V \cos N\theta}{2}$, where $0 \leq V \leq 1$ is a fitted visibility accounting for experimental noise and imperfect state preparation, see Fig. 33(a), the Fisher information $F(\theta) = \frac{1}{P(+|\theta)} \left(\frac{dP(+|\theta)}{d\theta} \right)^2 + \frac{1}{P(-|\theta)} \left(\frac{dP(-|\theta)}{d\theta} \right)^2$ is

$$F(\theta) = \frac{V^2 N^2 \sin^2 N\theta}{1 - V^2 \cos^2 N\theta}. \quad (95)$$

It reaches the maximum value $F_{\text{max}} = N^2 V^2$ when $\sin N\theta = \pm 1$. It is thus possible to detect useful entanglement and obtain sensitivities below the standard quantum limit when $V^2 > 1/N$ (*i.e.* $F > N$). Maximally entangled states are detected when $V^2 > (1 - \frac{1}{N})^2 + \frac{1}{N^2}$ [*i.e.* $F > (N - 1)^2 + 1$, obtained from Eq. (25) with $k = N - 1$]. Figure 33(b) gives an overview of the experimental results obtained with cat states of trapped ions. In [Monz et al. \(2011\)](#) the fringe visibility of parity oscillations was sufficiently high that the Fisher information witnesses metrologically useful genuine N -particle entanglement (and thus Heisenberg limit sensitivity) up to $N = 6$ ions ([Pezzè et al., 2015](#)).

We note that the output probabilities obtained with states (94) show fast oscillations as a function of θ , see Fig. 33. This seems to prevent the unambiguous estimate of a phase larger than $2\pi/N$ ([Berry et al., 2009](#); [Boto et al., 2000](#); [Mitchell, 2005](#); [Pezzè and Smerzi, 2007](#)). A phase estimation at the Heisenberg limit with no constraint on θ in $[-\pi, \pi]$ can be obtained using states (94) with different numbers of particles. The optimal sequence $N = 1, 2, 4, \dots, 2^p$ provides $\Delta\theta = 2.55/N_T$ (at $\theta = 0$), where $N_T = \sum_{n=0}^p 2^n = 2^{p+1} - 1$ is the total number of particles used ([Pezzè and Smerzi, 2007](#)).

It is finally worth pointing out that W states (that, ideally, are less useful than cat states for metrological applications) have been created experimentally with up to eight ions ([Häffner et al., 2005](#); [Roos et al., 2004](#)).

3. Spin squeezing and useful entanglement of many ions

Recently, entanglement and metrological spin-squeezing of hundreds of trapped $^9\text{Be}^+$ ions have been demonstrated in a Penning trap ([Bohnet et al., 2016](#)). The experiment realized a homogeneous Ising spin-spin interaction ([Kitagawa and Ueda, 1993](#)) (*i.e.* $\alpha \approx 0$ in Eq. 92), see Sec. III.B for a detailed analysis of the one-axis twisting model. Figure 34 shows ξ_R^2 , optimized over time, as a function of the number of ions ranging from $N = 20$ to $N = 220$. After longer interaction times, the state wraps around the Bloch sphere and spin squeezing is lost (see Sec. III.B). In this case, the experimental data of [Bohnet et al. \(2016\)](#) still show $F/N > 2.1$ signalling useful entanglement, where the Fisher information has been extracted with the Hellinger method ([Strobel et al., 2014](#)). For an analysis of decoherence in this system see [Foss-Feig et al. \(2013\)](#).

VII. WORKING ENTANGLEMENT-ENHANCED INTERFEROMETERS AND PROSPECTS FOR APPLICATIONS

Atomic ensembles are routinely used as precision sensors of inertial forces or external fields, or as atomic

clocks, which usually operate as Ramsey interferometers (Barrett *et al.*, 2016; Bord , 2002; Cronin *et al.*, 2009; Kitching *et al.*, 2011; Wynands, 2009; Wynands and Weyers, 2005). The accumulated phase is $\theta = \omega T_R$, where T_R is the interrogation time and ω is the parameter to be estimated. In an atomic sensor, $\omega = \Delta E/\hbar$, where ΔE is the energy shift between the two interferometer modes, induced by an external field, an acceleration, a rotation, or a force. In an atomic clock, $\omega = \omega_{LO} - \omega_0$, where ω_{LO} is the instantaneous frequency of a local oscillator to be locked to the transition frequency ω_0 between the two atomic levels that define the clock. The standard quantum limit for the estimation of the frequency ω is, from Eq. (21),

$$\Delta\omega_{\text{SQL}} = \frac{1}{T_R \sqrt{N\nu}}. \quad (96)$$

Using separable states, if quantum noise dominates over technical noise, the sensitivity can thus only be increased by increasing the interrogation time, the number N of particles, and/or the number ν of experimental runs. This is not always possible: every sensor has an optimal working range for T_R and N , and drifts always limit the length ν of the sequence in which identical state preparation can be assumed (Bak *et al.*, 1987). In general, increasing the atom number leads to systematic errors due to collisions that spoil the sensitivity. Interrogation times are limited by mechanical restrictions (*e.g.* the limited ballistic flight time in fountain clocks), by the desired temporal resolution (*e.g.* when measuring high-frequency magnetic fields in broadband magnetometers), or decoherence (*e.g.* the phase coherence time of the local oscillator in atomic clocks). Can entangled states improve the performance of atomic sensors and, if so, under what conditions?

Entangled states are typically more susceptible to decoherence than separable states during phase encoding. Thus their gain (*i.e.* their more favorable scaling of phase sensitivity with N) is often counterbalanced by the requirement of shorter interrogation times (Huelga *et al.*, 1997). Whether or not an improvement is possible can only be decided after a detailed analysis of a specific sensor taking its actual limitations into account, and ultimately hinges on an experimental demonstration. As an example we note that squeezing of light has recently been used to improve the performance of a laser-interferometer gravitational wave observatory (Aasi *et al.*, 2013; The LIGO Scientific Collaboration, 2011). The aim of this section is to review the state of the art of entanglement-enhanced measurements with atomic ensembles, and discuss prospects for applications.

A. The influence of noise and decoherence

A common and central aspect of quantum-enhanced metrology is the fragility of useful entangled states when coupled to the environment. Noise affects the preparation of the probe state, phase encoding, and detection. The literature has focused mainly on specific noise models, and only few general results are available. An important challenge is to devise protocols (*e.g.* differential schemes, or error correction), special phase sensing situations (*e.g.* non-Markovian, or transverse noise), and detection schemes that make metrology below the standard quantum limit robust against noise.

1. General results

The quantum Fisher information is convex in the state (Cohen, 1968; Fujiwara, 2001):

$$F_Q \left[\sum_k p_k \hat{\rho}_k(\theta) \right] \leq \sum_k p_k F_Q [\hat{\rho}_k(\theta)]. \quad (97)$$

This expresses the fact that mixing quantum states cannot increase the achievable estimation sensitivity. Furthermore, the quantum Fisher information never increases under any parameter-independent quantum channel (Petz, 2002):

$$F_Q [\Lambda(\hat{\rho}(\theta))] \leq F_Q [\hat{\rho}(\theta)], \quad (98)$$

where Λ is a θ -independent completely-positive trace-preserving map, which is a general model of the evolution of a system and its environment (Nielsen and Chuang, 2000; Rivas and Huelga, 2012). For example, the partial trace operation can only decrease the quantum Fisher information, consistent with the intuition that ignoring part of a system can only decrease the information about the estimated parameter. The equality sign in Eq. (98) is always obtained if Λ is unitary, because unitary θ -independent transformations can be absorbed into a redefinition of the optimal measurement.

2. Uncorrelated decoherence

A benchmark example of the fragility of frequency estimation with respect to noise has been pointed out by Huelga *et al.* (1997). Let us consider a single atom prepared in the superposition $(|a\rangle + |b\rangle)/\sqrt{2}$ that evolves freely for a time T_R , according to the Hamiltonian $\omega\hat{\sigma}_z$, and acquires a phase ωT_R in the presence of local Markovian dephasing. The probability to find the probe in its initial state is $P = [1 + e^{-\gamma T_R} \cos(\omega T_R)]/2$, where $\gamma \geq 0$ is the dephasing rate. If the phase estimation is repeated ν times (using a total of $N = \nu$ particles), a calculation of the Cram r-Rao bound gives

$\Delta\omega = \sqrt{2\gamma e/(NT_R)}$ at the optimal time $T_R = 1/(2\gamma)$. If we take a GHZ state $(|a\rangle^{\otimes N} + |b\rangle^{\otimes N})/\sqrt{2}$ in the presence of the same source of (single-particle) decoherence, the probability to find the probe in the initial state is $P = [1 + e^{-\gamma NT_R} \cos(\omega NT_R)]/2$: it oscillates N times faster than in the single-particle case (this is typical for GHZ states, see Sec. II.C.7), but the visibility of these oscillations decays N times faster. A calculation of the Cramér-Rao bound gives $\Delta\omega = \sqrt{2\gamma e/(NT_R)}$, the same as in the single-particle case, reached at the optimal time $T_R = 1/(2N\gamma)$. In this example, the ideal sensitivity enhancement offered by GHZ states for $\gamma = 0$ disappears in the presence of arbitrarily small uncorrelated dephasing ($\gamma > 0$). An optimization over probe states shows that at most a constant factor of $1/\sqrt{e}$ in the absolute frequency error can be gained (Escher *et al.*, 2011; Huelga *et al.*, 1997; Ulam-Orgikh and Kitagawa, 2001).

Frequency estimation in the presence of single-atom decoherence has been analyzed by Demkowicz-Dobrzański *et al.* (2012); Escher *et al.* (2011); Huelga *et al.* (1997); Kołodyński and Demkowicz-Dobrzański (2013); Shaji and Caves (2007); Smirne *et al.* (2016); and Ulam-Orgikh and Kitagawa (2001). When considering arbitrarily long interrogation times and taking the limit of large numbers of particles $N \rightarrow \infty$, an infinitesimally small amount of Markovian noise (independent of the particle number) prevents any sensible gain from the creation of entangled states (Demkowicz-Dobrzański *et al.*, 2012; Escher *et al.*, 2011). This result holds also for sequential feedback protocols when the channel associated with the system's evolution between two feedback operations is of full rank (Escher *et al.*, 2011). If the channel is of less than full rank, quantum error correction based on a sequential feedback protocol can be applied to counteract the effects of noise (Kessler *et al.*, 2014b) and quantum-enhanced phase estimation is possible.

For non-Markovian noise, on the other hand, quantum enhancement can be found in the limit $N \rightarrow \infty$. In particular, Chin *et al.* (2012) and Matsuzaki *et al.* (2011) have shown that a frequency variance $(\Delta\omega)^2 = \mathcal{O}(N^{-3/2})$ can be reached for an interrogation time $T_R = \mathcal{O}(N^{-1/2})$ under general models of non-Markovian phase noise; see also Berrada (2013); Macieszczak (2015); Smirne *et al.* (2016); and Szańkowski *et al.* (2014). Moreover, Chaves *et al.* (2013) have shown that if Markovian dephasing is acting along a spin direction perpendicular to the phase encoding, in the limit $N \rightarrow \infty$ a sensitivity $(\Delta\omega)^2 = \mathcal{O}(N^{-5/3})$ can be obtained for interrogation times $T_R = \mathcal{O}(N^{-1/3})$. The possibility to reach the Heisenberg limit $(\Delta\omega)^2 = \mathcal{O}(N^{-1})$ in this case has been discussed by Dür *et al.* (2014) and Kessler *et al.* (2014b) when making use of quantum error correction techniques (see below). We note that the orientation-dependent lifetime of spin-squeezed states has been investigated experimentally by Leroux *et al.* (2010b).

3. Correlated phase noise, differential interferometry, and decoherence-free subspaces

In many systems, a significant source of noise is correlated dephasing, where all atoms are subject to the same noise source. For instance, correlated dephasing is the major source of noise in experiments with ions stored in linear Paul traps (Monz *et al.*, 2011; Roos *et al.*, 2006). Collective phase fluctuations are caused by noisy stray fields inducing random energy shifts of the atomic levels. In the Markovian limit, the initial state $\hat{\rho}(0)$ evolves according to $\frac{d\hat{\rho}}{dt} = \frac{\gamma}{2}(\hat{J}_z\hat{\rho}\hat{J}_z - \frac{1}{2}\hat{\rho}\hat{J}_z^2 - \frac{1}{2}\hat{J}_z^2\hat{\rho})$. For states symmetric under particle exchange we have

$$\hat{\rho}_{\nu\mu}(t) = \hat{\rho}_{\nu\mu}(0)e^{-\frac{1}{4}\gamma t(\mu-\nu)^2}, \quad (99)$$

where $\hat{\rho}_{\nu\mu} = \langle \nu_z | \hat{\rho} | \mu_z \rangle$ and $|\nu_z\rangle$ are eigenstates of $\hat{J}_z$. In particular, Eq. (99) predicts that the coherence of an N -qubit GHZ state decays faster than that of a single qubit by a factor N^2 . This effect, also known as super-decoherence, has been investigated experimentally with maximally entangled states of $^{40}\text{Ca}^+$ ions by adding a variable delay between creation and coherence investigation (Monz *et al.*, 2011). The obtained data are consistent with a decay of off-diagonal terms of the density matrix with a $e^{-\frac{1}{4}\gamma t N^2}$ scaling law (Monz *et al.*, 2011), in full agreement with predictions for correlated Markovian noise at long evolution times.

Collective phase noise can be counteracted by encoding the qubits in a decoherence-free subspace (Lidar *et al.*, 1998; Zanardi and Rasetti, 1997), formally equivalent to a differential interferometer scheme (Landini *et al.*, 2014). If we consider the system to be made of two subsystems such that $\hat{J}_z = \hat{J}_z^{(1)} + \hat{J}_z^{(2)}$, and assume that the state of each subsystem is symmetric under particle exchange, then the components of the joint state evolve according to

$$\hat{\rho}_{\nu_1, \nu_2, \mu_1, \mu_2}(t) = \hat{\rho}_{\nu_1, \nu_2, \mu_1, \mu_2}(0)e^{-\frac{1}{4}\gamma t(\mu_1 + \mu_2 - \nu_1 - \nu_2)^2}. \quad (100)$$

Components with $\mu_1 + \mu_2 = \nu_1 + \nu_2$ are insensitive to phase noise. This condition is met by nontrivial states achieving the Heisenberg limit of phase sensitivity (Dorner, 2012; Jeske *et al.*, 2014; Landini *et al.*, 2014). Notice that under certain conditions, entanglement can be preserved in the presence of collective dephasing even if the state is not in a decoherence-free subspace (Carnio *et al.*, 2015). The experiment of Roos *et al.* (2006) demonstrated the preparation, and proved the robustness, of a state of two ions in a decoherence-free subspace. In Monz *et al.* (2011) decoherence-free states with up to 8 ions were constructed, achieving coherence times of ~ 100 ms.

4. Error correction

Quantum error correction techniques (Nielsen and Chuang, 2000; Shor, 1995; Steane, 1996) can be used to counteract the effect of noise in quantum metrology (Arrad *et al.*, 2014; Dür *et al.*, 2014; Herrera-Martí *et al.*, 2015; Kessler *et al.*, 2014b; Ozeri, 2013; Plenio and Huelga, 2016). These schemes protect the quantum states on which a signal parameter is imprinted by employing ancilla qubits that neither interact with the parameter nor are subject to noise. Noise operators map the state to ancilla Hilbert subspaces orthogonal to the phase encoding subspace. Error correction is accomplished by means of projective measurements into the orthogonal subspaces, and then applying a correction sequence. These techniques can be used to extend the coherence time and/or to achieve the Heisenberg limit for certain noise models (Arrad *et al.*, 2014; Dür *et al.*, 2014; Kessler *et al.*, 2014b; Ozeri, 2013). Lu *et al.* (2015) have provided conditions under which the quantum Fisher information, rather than the full quantum state, can be protected under a class of noisy channels.

Finally, it has been shown that entanglement between the probe and an ancillary system in the preparation and measurement stage can be useful in the presence of uncorrelated noise (Demkowicz-Dobrzański and Maccone, 2014; Haine and Szigeti, 2015; Haine *et al.*, 2015; Huang *et al.*, 2016).

5. Particle losses

The incoherent loss of particles generally has a strong impact on the usefulness of a state for quantum metrology. A paradigmatic example is the NOON state (Sec. II.C.7): the loss of a single particle transforms the NOON state into an incoherent mixture $\frac{1}{2}|N-1\rangle_a\langle N-1| \otimes |0\rangle_b\langle 0| + \frac{1}{2}|0\rangle_a\langle 0| \otimes |N-1\rangle_b\langle N-1|$ that is useless for phase sensing. The impact of losses in quantum metrology has been extensively studied in the context of optical interferometry (Dorner *et al.*, 2009; Huver *et al.*, 2008; Kacprowicz *et al.*, 2010; Knysh *et al.*, 2011; Lee *et al.*, 2009; Ono and Hofmann, 2010). For atoms, losses are especially relevant in experiments with Bose-Einstein condensates. There, one-body losses are due to collisions of condensate atoms with residual hot atoms due to imperfect vacuum, whereas two- and three-body losses are caused by inelastic collisions and are relevant in dense samples. The impacts of losses (Li *et al.*, 2008a; Sinatra *et al.*, 2013, 2012a, 2011; Spehner *et al.*, 2014), finite temperatures (Sinatra *et al.*, 2011) and collective dephasing (Ferrini *et al.*, 2008, 2011) during the preparation of spin-squeezed states via one-axis twisting (see Sec. III.B) have been studied (Sinatra *et al.*, 2012b). In particular, Li *et al.* (2008a) have shown that $\xi_R^2 = \mathcal{O}(N^{-4/15})$ for one-body losses, assuming $N \rightarrow \infty$ and an opti-

mal evolution time. For two-body losses, the optimal ξ_R^2 does not depend on N , while for three-body losses $\xi_R^2 = \mathcal{O}(N^{4/15})$ and there is a finite optimal number of particles for squeezing.

6. Finite detection efficiency

The sensitivity of parameter estimation is directly related to the distinguishability of probability distributions of some measurement observable when changing the parameter, see Sec. II.B.4. Finite detection efficiency blurs the probability distributions and limits their distinguishability. Low efficiency in the detection of large atom numbers is indeed one of the main limitations in many current quantum-enhanced metrology experiments.

A common detection method for neutral atoms is absorption imaging. The precision of this technique on mesoscopic ensembles is at the level of four atoms (Muessel *et al.*, 2015; Ockeloen *et al.*, 2010; Schmied *et al.*, 2016) (standard deviation on the detection of hundreds of atoms). Very high sensitivity in fluorescence detection of many atoms has been shown by spatially resolving each atom in an optical lattice (Bakr *et al.*, 2009; Nelson *et al.*, 2007; Sherson *et al.*, 2010). This technique can image and count individual atoms in large ensembles but do not determine the exact atom number as atom pairs are quickly lost due to light-assisted collisions. There are prospects to improve the detection efficiency using fluorescence imaging in Bose-Einstein condensates (Hume *et al.*, 2013; Stroescu *et al.*, 2015) and optical cavity-aided measurements with cold thermal ensembles (Zhang *et al.*, 2012).

Davis *et al.* (2016) have shown that a nonlinear readout (*i.e.* applying a nonlocal evolution prior to standard atom number measurement) can make phase estimation robust against detection noise removing the requirement of single-atom resolution to reach a precision approaching the Heisenberg limit. A nonlinear readout has been exploited by Leibfried *et al.* (2004, 2005) for quantum-enhanced metrology with GHZ states of $N \lesssim 6$ trapped ions (see Sec. VI.B), and, more recently, by Linemann *et al.* (2016) for the realization of a SU(1,1) interferometer with a spinor Bose-Einstein condensate (See Sec. IV.C.3). Hosten *et al.* (2016b) have exploited a nonlinear evolution after phase encoding to “magnify” the phase signal. In this experiment, the collective spin of $N = 5 \times 10^5$ ^{87}Rb atoms is first squeezed via a light-mediated interaction in an optical cavity (see Sec. V.B), and then rotated by an angle θ . After a second period of collective spin interactions in the cavity, the state is detected via fluorescence imaging. Hosten *et al.* (2016b) demonstrated a phase sensitivity 8 dB below the standard quantum limit, using a detection with a technical noise floor 10 dB above projection noise.

B. Atomic clocks

The performance of an atomic clock (Ludlow *et al.*, 2015; Wynands, 2009; Wynands and Weyers, 2005) can be characterized by its frequency accuracy and stability (Vanier and Audoin, 1992). Accuracy refers to the frequency offset from the ideal value, whereas stability describes the fluctuations of the instantaneous frequency $\omega(t)$ from the average frequency ω_0 . Improving the stability of time-keeping is one of the primary targets of quantum-enhanced phase estimation. It also allows for faster evaluation of systematic errors, which in turn can improve the accuracy (Nicholson *et al.*, 2015). Currently, the stability of atomic clocks is nearly (Bloom *et al.*, 2014; Hinkley *et al.*, 2013; Nicholson *et al.*, 2012) or already—as in the case of fountain clocks (Santarelli *et al.*, 1999)—limited by the standard quantum limit. In principle, the clock stability is optimized by extending the interrogation time until atomic decoherence dominates in the interferometer output signal. In practice, a main limitation of current clocks, in particular of optical clocks using trapped atoms, is frequency noise of the local oscillator, which limits the stability of frequency comparisons well short of the limits imposed by atomic decoherence (Chou *et al.*, 2011; Wineland *et al.*, 1998).

The standard figure of merit for quantifying the stability of a clock is the Allan standard deviation of the relative frequency fluctuations $y(t) = \omega(t)/\omega_0$ (Vanier and Audoin, 1992),

$$\sigma(\tau) = \sqrt{\frac{\sum_{k=1}^{\nu-1} (y_{k+1} - y_k)^2}{2(\nu-1)}}, \text{ with } y_k = \frac{1}{\tau} \int_{t_k}^{t_k+\tau} dt y(t), \quad (101)$$

where $y_{k+1} - y_k$ is the relative change of the clock frequency between two successive averaging bins of duration τ . The Allan deviation quantifies the discrepancy between two consecutive observations of the clock frequency averaged for a time τ , including dead times in preparation of the sample and data acquisition. For an atomic clock operating at the standard quantum limit with uncorrelated (shot-to-shot) measurements, we have

$$\sigma_{\text{SQL}}(\tau) = \frac{1}{\omega_0 T_R} \sqrt{\frac{T_{\text{cycle}}}{N\tau}}, \quad (102)$$

where $T_{\text{cycle}} \geq T_R$ is the clock cycle duration. Overcoming this limit is a concrete demonstration of the potential benefits of entanglement for improving clock stability.

Theoretical analyses of entanglement-enhanced atomic clocks have focused on the impact of fluctuations of the local oscillator frequency. In this case, entanglement may be beneficial. André *et al.* (2004) have shown, by optimizing over a family of squeezed states, the possibility of achieving an entanglement-enhanced stability $\sigma(\tau) \propto 1/(N^{2/3}\sqrt{\tau})$. Bužek *et al.* (1999) have considered

a different optimization strategy and shown the possibility of achieving $\sigma(\tau) \propto 1/(N\sqrt{\tau})$ for large N . Using a numerical optimization of probe states, measurements with up to 20 atoms, and realistic local oscillator noise, Rosenband (2011) has shown that the approach of André *et al.* (2004) is optimal up to 15 atoms and overtaken by the approach of Bužek *et al.* (1999) for larger N . The problem of devising efficient protocols for creating optimal states and measurements is still open. Furthermore, it has been shown that clock interrogation schemes exploiting adaptive protocols can realize sub-standard-quantum-limit instability (Borregaard and Sørensen, 2013; Kessler *et al.*, 2014a; Mullan and Knill, 2014; Rosenband and Leibbrandt, 2013; Rosenband, 2011). In these protocols, the phase noise of the laser is tracked and stabilized over long time scales.

1. Entanglement-assisted atomic clocks

Atomic clocks with a performance beating the standard quantum limit have been experimentally demonstrated with two trapped ions (Meyer *et al.*, 2001), ensembles of cold thermal atoms (Hosten *et al.*, 2016a; Leroux *et al.*, 2010b; Louchet-Chauvet *et al.*, 2010) and Bose-Einstein condensates (Gross *et al.*, 2010; Kruse *et al.*, 2016; Ockeloen *et al.*, 2013). The neutral atom experiments have realized microwave-frequency clocks on magnetic field-insensitive hyperfine transitions with spin-squeezed states as input.²³ Leroux *et al.* (2010b) measured an Allan deviation of $\sigma(\tau) = 1.1 \times 10^{-9} \text{ s}^{1/2}/\sqrt{\tau}$ for their squeezed clock, corresponding to an improvement of 4.5 dB in variance beyond the standard quantum limit, see Fig. 35. More recently, Hosten *et al.* (2016a) performed a similar measurement, reaching $\sigma(\tau) = 9.7 \times 10^{-11} \text{ s}^{1/2}/\sqrt{\tau}$, or 10.5 dB in variance beyond the standard quantum limit. While these experiments are impressive proof-of-principle demonstrations of entanglement-enhanced atomic clocks, they do not yet reach the frequency stability of state-of-the-art fountain clocks. This is mostly due to a much shorter Ramsey interrogation time of $T_R \sim 200 \mu\text{s}$, limited by the noise of the microwave local oscillator. For comparison, current fountain clocks using uncorrelated atoms operate with $T_R \sim 1 \text{ s}$ and reach $\sigma(\tau) \sim 10^{-14} \text{ s}^{1/2}/\sqrt{\tau}$ (Wynands, 2009).

2. Proposals for entanglement-assisted lattice clocks

Optical atomic clocks that interrogate narrow optical transitions in ensembles of atoms trapped in an optical

²³ Note that, to date, squeezing has not yet been realized on an optical transition. Monz *et al.* (2011) have created GHZ states on the optical clock transition of $^{40}\text{Ca}^+$ ions.




FIG. 35 **Stability of an atomic clock beating the standard quantum limit.** Allan deviation of an atomic clock with a spin-squeezed input state (solid red line with error bars), with $T_R = 200 \mu\text{s}$, $N = 3.5 \times 10^4$ and $T_{\text{cycle}} = 9 \text{ s}$. The red dotted line corresponds to $\sigma(\tau) = 1.1 \times 10^{-9} \text{ s}^{1/2}/\sqrt{\tau}$. Open black circles: reference measurements with uncorrelated atoms. The dashed line indicates the SQL. Taken from [Leroux et al. \(2010b\)](#).

lattice are currently the most precise and accurate measurement devices ([Ludlow et al., 2015](#); [Ye et al., 2008](#)), reaching stabilities of $\sigma(\tau) \sim 10^{-16} \text{ s}^{1/2}/\sqrt{\tau}$, integrating down to the lower 10^{-18} level after a few thousand seconds of averaging ([Nicholson et al., 2015](#)). Their stability is within a factor of two above the quantum projection noise limit ([Bloom et al., 2014](#); [Hinkley et al., 2013](#); [Ludlow et al., 2008](#)). Since, in these systems, there is a limit to the exploitable number of atoms, using entanglement to increase the sensitivity is of considerable interest. So far, only theoretical studies are available. [Meiser et al. \(2008\)](#) have studied the creation of spin-squeezed states in a neutral-atom optical lattice clock through a cavity-based QND measurement. [Weinstein et al. \(2010\)](#) considered the creation of GHZ states through the on-site interaction of an atom moving across the lattice. A further interesting possibility is to exploit long-range Rydberg-dressing interactions between atoms at different lattice sites induced by laser excitations ([Bouchoule and Mølmer, 2002b](#); [Henkel et al., 2010](#); [Pupillo et al., 2010](#)) [see [Browaeys et al. \(2016\)](#) and [Saffman et al. \(2010\)](#) for reviews] to generate spin-squeezed ([Gil et al., 2014](#)) and non-Gaussian entangled states ([Macrì et al., 2016](#)).

C. Optical magnetometers

Optical magnetometers ([Budker and Kimball, 2013](#); [Budker and Romalis, 2007](#)) exploit the interaction of light with a spin-polarized atomic ensemble (either thermal atoms in a vapor cell at room temperature or laser-

cooled atom in an optical trap²⁴) to measure the strength of a magnetic field ([Budker et al., 2002](#)). Generally, the device consists of N spin- F atoms initially optically pumped into a fully polarized state (pointing along the x -axis, for instance), such that the collective spin of the ensemble has length $\langle \hat{J}_x \rangle = FN$. A weak magnetic field along the y -axis causes the precession of the collective spin in the $x-z$ plane at a rate $g\mu_B B/\hbar$, where g is the gyromagnetic ratio, μ_B the Bohr magneton, and B the magnetic field strength. The atomic spin precession is measured by observing the polarization rotation of the probe light transmitted through the atomic sample. The interrogation time is limited by spin-relaxation due to collisions of the atoms with the cell walls that enclose the vapor or, for high density gases, due to spin-exchange collisions ([Budker and Romalis, 2007](#)). The sensitivity of current optical magnetometers is fundamentally limited by quantum noise in the form of atomic projection noise and photon (polarization) shot-noise. The back-action of light onto the atoms is a further limiting factor. Current optical magnetometers are approaching quantum noise limits ([Kominis et al., 2003](#); [Wasilewski et al., 2010](#)).

Polarization squeezing of the probe light or spin squeezing of the atomic ensemble ([Auzinsh et al., 2004](#); [Geremia et al., 2003](#); [Petersen et al., 2005](#)) can be used to reduce the quantum noise. Spin-squeezing induced by continuous quantum nondemolition measurements ([Auzinsh et al., 2004](#)) can enhance the sensitivity, and a scaling $N^{-3/4}$ is predicted when using shot-noise-limited light. This can be further pushed to the Heisenberg limit $1/N$ if squeezed light is used. However, single-atom spin-relaxation modifies this picture, and a reduction of quantum noise below the standard quantum limit is expected only for interrogation times shorter than the spin coherence time ([Auzinsh et al., 2004](#)). Spin-squeezed states can thus increase the measurement bandwidth (*i.e.* they are useful for short interrogation times) without loss of sensitivity, with important applications in biology and medicine. For long interrogation times, entanglement-enhanced sensitivity using spin squeezed states is expected for high-density ensembles due to the suppression of spin-relaxation noise ([Kominis, 2008](#); [Vasilakis et al., 2011](#)).

²⁴ Vapor cell magnetometers are currently the most sensitive measuring devices for low-frequency magnetic fields, reaching sensitivity levels below $1 \text{ fT}/\sqrt{\text{Hz}}$ with probe volumes of 10^6 - $10^{12} (\mu\text{m})^3$ and large atom numbers $N = 10^{11}$ - 10^{15} . Laser-cooled atoms in an optical trap contain a much smaller number of atoms $N = 10^5$ - 10^8 but are more compact and better suited for field measurements with high spatial resolution.




FIG. 36 **Entanglement-assisted optical magnetometry.** (a) Measurement sensitivity obtained in absence (squares) and in presence (open circles) of entanglement between atomic ensembles in a two-cell radio-frequency magnetometer. An improvement is demonstrated for short radio-frequency pulse durations τ , where $\delta_{\text{RF}} = 1/\tau$. Taken from Wasilewski *et al.* (2010). (b) Measurement sensitivity as a function of the number of atoms with a coherent spin state (blue circles) and spin-squeezed (red diamonds) probe. Solid lines are expected results while dashed lines are different contributions to the sensitivity. Taken from Sewell *et al.* (2012).

1. Entanglement-assisted optical magnetometers

An improvement in magnetic field sensing via the reduction of the optical polarization noise has been demonstrated by Horrom *et al.* (2012) and Wolfgramm *et al.* (2010) using squeezed probe light. Atomic projection noise limited and entanglement-assisted magnetometry has been demonstrated in the experiment of Wasilewski *et al.* (2010), who realized a pulsed RF magnetometer made of two atomic ensembles (two vapor cells with opposite atomic polarizations) with $N = 3.6 \times 10^{12}$ Cs atoms. A first light pulse is used to entangle the two ensembles via a quantum nondemolition measurement, as first demonstrated by Julsgaard *et al.* (2001). This is a useful resource to improve the sensitivity of the magnetometer for large bandwidths, as shown in Fig. 36(a). The improvement is observed for interrogation times shorter than the entanglement lifetime of 4 ms. The increase of measurement bandwidth in DC magnetometers has been shown by Shah *et al.* (2010) using continuous quantum nondemolition measurements. Sewell *et al.* (2012) generated spin-squeezed states in ensembles of cold spin-1 ^{87}Rb atoms via quantum nondemolition measurements (Koschorreck *et al.*, 2010a,b) and applied them to optical magnetometry. Results corresponding to interrogation pulses of $5 \mu\text{s}$ (giving a measurement bandwidth of 200 kHz) are shown in Fig. 36(b).

D. Scanning-probe magnetometers using Bose-Einstein condensates

Trapped Bose-Einstein condensates are particularly well suited for sensing applications requiring high spa-




FIG. 37 **Entanglement-enhanced Bose-Einstein condensate interferometer.** Performance of a spin-squeezed Bose-Einstein condensate interferometer expressed in terms of ξ_R^2 as a function of varying interrogation times T_R for squeezed (blue diamonds) and coherent (red circles) input states. Dashed lines model constant performance of $\xi_R^2 = -4.0$ dB (squeezed state) and $+0.2$ dB (coherent state) plus technical noise due to shot-to-shot frequency fluctuations. Inset: scanning probe measurements of microwave fields. The Bose-Einstein condensate (blue) is translated near the surface of an atom chip to measure the spatial dependence of microwave magnetic near-fields. Figure adapted from Ockeloen *et al.* (2013).

tial resolution, taking advantage of their small size, high degree of coherence, and the availability of sophisticated techniques for precise positioning of the atoms. Current Bose-Einstein condensate magnetometers reach sensitivities on the order of $10 \text{ pT}/\sqrt{\text{Hz}}$ with probe volumes of 10^2 - $10^4 (\mu\text{m})^3$ (Aigner *et al.*, 2008; Eto *et al.*, 2013; Vengalattore *et al.*, 2007; Wildermuth *et al.*, 2006). In these systems, the required small probe size gives rise to an upper bound on the atom number due to density-dependent collisional trap losses. It is thus crucial to use entangled states to increase the sensitivity without increasing the atom number.

Ockeloen *et al.* (2013) have demonstrated a scanning probe atom interferometer operating beyond the standard quantum limit. Bose-Einstein condensates of ^{87}Rb atoms were prepared on an atom chip and spin-squeezed by means of collisions in a spin-dependent trapping potential (see Sec. III.B). Using the spin-squeezed state as input, a Ramsey interferometer sequence was implemented, achieving a performance of 4 dB below the standard quantum limit with $N = 1400$ atoms for interrogation times up to $T_R = 10 \text{ ms}$, see Fig. 37. This interferometer is sensitive to microwave magnetic fields, with a sensitivity of $77 \text{ pT}/\sqrt{\text{Hz}}$ using a small probe volume of $20 (\mu\text{m})^3$. The interferometer was operated as a scanning probe, by translating the Bose-Einstein condensate on the atom chip to measure the spatial dependence of

an on-chip microwave near-field.

The magnetometer of [Muessel *et al.* \(2014\)](#) consists of ~ 30 independent Bose-Einstein condensates in a one-dimensional optical lattice each containing about $N = 400$ ^{87}Rb atoms. Spin squeezing is created via one-axis twisting (see Sec. III.B), with interactions tuned by a Feshbach resonance. Subsequently, a microwave π -pulse transfers the squeezing to a magnetic-field-sensitive hyperfine transition. This experiment demonstrated sub-SQL magnetometry (3.4 dB below the standard quantum limit) with interrogation times up to $340\ \mu\text{s}$, reaching a sensitivity of $1.9\ \text{nT}/\sqrt{\text{Hz}}$ for static magnetic fields. The array configuration is particularly well suited for differential measurements and magnetic field gradiometry.

Taking the small probe volume into account, the experiments of [Muessel *et al.* \(2014\)](#) and [Ockeloen *et al.* \(2013\)](#) already achieve state-of-the-art sensitivity ([Budker and Kimball, 2013](#)).

E. Nonlocal phase encoding

Usually, the coupling between an atomic ensemble and an external field is local in the particles. The phase-encoding transformation of standard (or linear) interferometers is thus modeled by $\exp(-i\theta \sum_{i=1}^N \hat{\sigma}_n^{(i)}/2)$, where $\hat{\sigma}_n^{(i)}$ is a single-particle Pauli operator (see Sec. II). Yet in some cases (see below) the coupling between the atoms and the external field to be measured may be associated with a nonlocal Hamiltonian that involves interactions between the particles. Nonlinear interferometry has been first proposed by [Luis \(2004\)](#) and extensively studied in the literature ([Boixo *et al.*, 2008, 2007](#); [Choi and Sundaram, 2008](#); [Luis, 2007](#); [Rivas and Luis, 2010](#); [Roy and Braunstein, 2008](#)). In this case, the notions of a standard quantum limit (*i.e.* the highest phase sensitivity achievable by separable states) and a Heisenberg limit (*i.e.* the ultimate allowed phase sensitivity) discussed in Sec. II.C still hold. However, the scalings of these bounds with the number of particles in the probe state depend on the specific phase-encoding Hamiltonian. For instance, [Boixo *et al.* \(2008, 2007\)](#); and [Choi and Sundaram \(2008\)](#) have considered phase encoding of the kind $\exp[-i\theta(\sum_{i=1}^N \hat{\sigma}_n^{(i)}/2)^k]$. For $N \gg 1$, the standard quantum limit and Heisenberg limit become

$$\Delta\theta_{\text{SQL}} = \frac{\alpha_k}{N^{k-1/2}\sqrt{\nu}} \text{ and } \Delta\theta_{\text{HL}} = \frac{\beta_k}{N^k\sqrt{\nu}}, \quad (103)$$

for $N \gg 1$, respectively, where α_k and β_k are constant prefactors. For $k > 1$ these bounds have faster scalings of phase sensitivity with the number of particles than $\Delta\theta_{\text{SQL}} = 1/\sqrt{N\nu}$ and $\Delta\theta_{\text{HL}} = 1/(N\sqrt{\nu})$, discussed in Sec. II for linear interferometers. Several systems have been proposed to observe these anomalous scalings of phase sensitivity, including Kerr nonlinearities ([Beltrán](#)

and [Luis, 2005](#)), collisions in Bose-Einstein condensates ([Boixo *et al.*, 2009, 2008, 2007](#); [Choi and Sundaram, 2008](#); [Rey *et al.*, 2007](#)), nonlinearities in nano-mechanical resonators ([Woolley *et al.*, 2008](#)), double-pass effective nonlinearities with a cold atomic ensemble ([Chase *et al.*, 2009](#)), topological excitations in nonlinear systems ([Negretti *et al.*, 2008](#)), and atom-photon interactions in cold atom systems ([Napolitano and Mitchell, 2010](#)). Nonlinear interferometers find applications in optical magnetometry ([Chase *et al.*, 2009](#); [Sewell *et al.*, 2014](#)) and in the measurement of atomic scattering properties ([Rey *et al.*, 2007](#)).

[Napolitano *et al.* \(2011\)](#) have engineered an effective atom-light Hamiltonian $\hat{H}_{\text{eff}} = \alpha \hat{J}_z \hat{S}_z + \beta \hat{J}_z \hat{S}_z N_{\text{ph}}/2$ using an optical pulse passing through a cold atom ensemble of ^{87}Rb atoms. Here, $\hat{J}$ is collective atomic spin, $\hat{S}$ is the Stokes vector of the light, and N_{ph} is the photon number. $\hat{H}_{\text{eff}}$ describes a paramagnetic (nonlinear) Faraday rotation of the light beam ([Napolitano and Mitchell, 2010](#)) with rotation angle proportional to $\langle \hat{J}_z \rangle$. The coefficients α and β strongly depend on the detuning of the light beam, and experimental conditions for which $\alpha = 0$ can be achieved ([Napolitano and Mitchell, 2010](#)). The second term in $\hat{H}_{\text{eff}}$ accounts for nonlinear photon-photon interactions. The measurement uncertainty achieved with unentangled photons is $\Delta J_z = 1/(AN_{\text{ph}}^{1/2} + BN_{\text{ph}}^{3/2})$ where $A \propto \alpha$ and $B \propto \beta$. A scaling $\Delta J_z \propto N_{\text{ph}}^{-3/2}$, obtained by increasing the photon number, has been demonstrated by [Napolitano *et al.* \(2011\)](#). [Sewell *et al.* \(2014\)](#), using a nonlinear Faraday rotation based on alignment-to-orientation conversion ([Budker *et al.*, 2000](#)), have demonstrated a sensitivity $\Delta J_z \propto N_{\text{ph}}^{-3/2}$, overcoming the sensitivity of a linear measurement with the same photon number.

VIII. OUTLOOK

The possibility to achieve phase sensitivities beyond the standard quantum limit using atomic ensembles is the fuel for a vivid and exciting research activity that focuses on the engineering, characterization and manipulation of entangled many-body quantum states. On the theoretical side, a comprehensive and convincing conceptual framework has been developed for quantum metrology with entangled states, centered around key concepts such as spin-squeezing and Fisher information. Connections between the concepts of quantum metrology and other fields of quantum information (such as Bell correlations, quantum Zeno dynamics, and Einstein-Podolsky-Rosen entanglement) can shed new light on quantum technologies. Current active research trends are centered on quantum-enhanced metrology, taking into account relevant experimental imperfections and searching for protocols where the fragile entanglement is protected against

noise sources.

On the experimental side, many of the key concepts that were proposed during the past decades, such as spin-squeezing created via particle-particle or atom-light interactions, have recently been implemented in proof-of-principle experiments. The progress—in terms of sensitivity gain with respect to the standard quantum limit—has been extremely fast, in particular when compared to squeezing of light. In atomic ensembles, less than ten years after the first spin-squeezing experiments, impressive gains up to 20 dB have been achieved. The field is now at the verge of moving from proof-of-principle experiments to applications in entanglement-enhanced precision measurements of time, external fields, and forces.

ACKNOWLEDGMENTS

We are indebted to our colleagues and collaborators with whom we have shared many useful discussions over the past years. In particular, we would like to thank M. Fattori, C. Klempt, W. Muessel, L. Santos, H. Strobel, and G. Tóth. R.S. and P.T. also acknowledge discussions with B. Allard, M. Fadel, and T. Zibold, as well as funding from the Swiss National Science Foundation. L.P. acknowledges financial support by MIUR through FIRB Project No. RBFR08H058. All authors acknowledge financial support by EU-STREP Project QIBEC, No. FP7-ICT-2011-C.

REFERENCES

- Aasi, J, J. Abadie, B. P. Abbott, R. Abbott, T. D. Abbott, M. R. Abernathy, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, et al. (2013), “Enhanced sensitivity of the ligo gravitational wave detector by using squeezed states of light,” *Nat. Photon.* **7**, 613–619.
- Agarwal, G S (1981), “Relation between atomic coherent-state representation, state multipoles, and generalized phase-space distributions,” *Phys. Rev. A* **24**, 2889–2896.
- Agarwal, G S, and R. R. Puri (1990), “Cooperative behavior of atoms irradiated by broadband squeezed light,” *Phys. Rev. A* **41**, 3782–3791.
- Agarwal, G S, and R. R. Puri (1994), “Atomic states with spectroscopic squeezing,” *Phys. Rev. A* **49**, 4968–4971.
- Agarwal, G S, R. R. Puri, and R. P. Singh (1997), “Atomic Schrödinger cat states,” *Phys. Rev. A* **56**, 2249–2254.
- Aigner, S, L. Della Pietra, Y. Japha, O. Entin-Wohlman, T. David, R. Salem, R. Folman, and J. Schmiedmayer (2008), “Long-range order in electronic transport through disordered metal films,” *Science* **319** (5867), 1226–1229.
- Albiez, M, R. Gati, J. Fölling, S. Hunsmann, M. Cristiani, and M. K. Oberthaler (2005), “Direct observation of tunneling and nonlinear self-trapping in a single bosonic Josephson junction,” *Phys. Rev. Lett.* **95**, 010402.
- Amico, L, R. Fazio, A. Osterloh, and V. Vedral (2008), “Entanglement in many-body systems,” *Rev. Mod. Phys.* **80**, 517–576.
- Ananikian, D, and T. Bergeman (2006), “Gross-Pitaevskii equation for Bose particles in a double-well potential: Two-mode models and beyond,” *Phys. Rev. A* **73**, 013604.
- André, A, A. S. Sørensen, and M. D. Lukin (2004), “Stability of atomic clocks based on entangled atoms,” *Phys. Rev. Lett.* **92**, 230801.
- Anglin, J R, P. Drummond, and A. Smerzi (2001), “Exact quantum phase model for mesoscopic Josephson junctions,” *Phys. Rev. A* **64**, 063605.
- Apellaniz, I, B. Lücke, J. Peise, C. Klempt, and G. Toth (2015a), “Detecting metrologically useful entanglement in the vicinity of Dicke states,” *New J. Phys.* **17**, 083027.
- Apellaniz, Iagoba, Matthias Kleinmann, Otfried Gühne, and Géza Tóth (2015b), “Optimal detection of metrologically useful entanglement,” *arXiv:1511.05203 [quant-ph]*.
- Appel, J, P. J. Windpassinger, D. Oblak, N. Hoff, U. B. Kjærgaard, and E. S. Polzik (2009), “Mesoscopic atomic entanglement for precision measurements beyond the standard quantum limit,” *PNAS* **106**, 10960.
- Aragone, C, G. Guerri, S. Salamo, and J. L. Tani (1974), “Intelligent spin states,” *J. Phys. A: Math., Nucl. Gen.* **7**, L149.
- Arecchi, F T, E. Courtens, R. Gilmore, and H. Thomas (1972), “Atomic coherent states in quantum optics,” *Phys. Rev. A* **6**, 2211–2237.
- Arrad, G, Y. Vinkler, D. Aharonov, and A. Retzker (2014), “Increasing sensing resolution with error correction,” *Phys. Rev. Lett.* **112**, 150801.
- Auzinsh, M, D. Budker, D. F. Kimball, S. M. Rochester, J. E. Stalnaker, A. O. Sushkov, and V. V. Yashchuk (2004), “Can a quantum nondemolition measurement improve the sensitivity of an atomic magnetometer?” *Phys. Rev. Lett.* **93**, 173002.
- Bak, P, C. Tang, and K. Wiesenfeld (1987), “Self-organized criticality: An explanation of the $1/f$ noise,” *Phys. Rev. Lett.* **59**, 381–384.
- Bakr, W S, J. I. Gillen, A. Peng, S. Fölling, and M. Greiner (2009), “A quantum gas microscope for detecting single atoms in a Hubbard-regime optical lattice,” *Nature* **462** (7269), 74–77.
- Bar-Gill, Nir, Christian Gross, Igor Mazets, Markus Oberthaler, and Gershon Kurizki (2011), “Einstein-Podolsky-Rosen correlations of ultracold atomic gases,” *Phys. Rev. Lett.* **106**, 120404.
- Barontini, G, L. Hohmann, F. Haas, J. Estève, and J. Reichel (2015), “Deterministic generation of multiparticle entanglement by quantum Zeno dynamics,” *Science* **349** (6254), 1317–1321.
- Barrett, B, A. Bertoldi, and P. Bouyer (2016), “Inertial quantum sensors using light and matter,” *Physica Scripta* **91** (5), 053006.
- Behbood, N, F. Martin Ciurana, G. Colangelo, M. Napolitano, G. Tóth, R. J. Sewell, and M. W. Mitchell (2014), “Generation of macroscopic singlet states in a cold atomic ensemble,” *Phys. Rev. Lett.* **113**, 093601.
- Bell, J S (1964), “On the Einstein Podolsky Rosen paradox,” *Physics* **1**, 195–200.
- Beltrán, J, and A. Luis (2005), “Breaking the Heisenberg limit with inefficient detectors,” *Phys. Rev. A* **72**, 045801.
- Benhelm, J, G. Kirchmair, C. F. Roos, and R. Blatt (2008), “Towards fault-tolerant quantum computing with trapped ions,” *Nat. Phys.* **4** (6), 463–466.
- Berrada, K (2013), “Non-Markovian effect on the precision of parameter estimation,” *Phys. Rev. A* **88**, 035806.

- Berrada, T, S. van Frank, R. Bücker, T. Schumm, J.-F. Schaff, and J. Schmiedmayer (2013), “Integrated Mach-Zehnder interferometer for Bose-Einstein condensates,” *Nat. Comm.* **4**, 2077.
- Berrada, T, S. van Frank, R. Bücker, T. Schumm, J.-F. Schaff, J. Schmiedmayer, B. Juliá-Díaz, and A. Polls (2016), “Matter-wave recombiners for trapped Bose-Einstein condensates,” *Phys. Rev. A* **93**, 063620.
- Berry, D W, B. L. Higgins, S. D. Bartlett, M. W. Mitchell, G. J. Pryde, and H. M. Wiseman (2009), “How to perform the most accurate possible phase measurements,” *Phys. Rev. A* **80**, 052114.
- Berry, D W, and B. C. Sanders (2002), “Near-optimal two-mode spin squeezing via feedback,” *Phys. Rev. A* **66**, 012313.
- Bhattacharyya, A (1943), “On a measure of divergence between two statistical populations defined by their probability distributions,” *Bull. Calcutta Math. Soc.* **35**, 99.
- Biedenharn, L G, and J. C. Louck (1981), *Angular Momentum in Quantum Physics: Theory and Applications* (Addison-Wesley, Reading, MA).
- Blatt, R, and C. F. Roos (2012), “Quantum simulations with trapped ions,” *Nat. Phys.* **4**, 277.
- Blatt, R, and D. J. Wineland (2008), “Entangled states of trapped atomic ions,” *Nature* **453**, 1008.
- Bloch, I, J. Dalibard, and W. Zwerger (2008), “Many-body physics with ultracold gases,” *Rev. Mod. Phys.* **80**, 885–964.
- Bloom, B J, T. L. Nicholson, J. R. Williams, S. L. Campbell, M. Bishof, X. Zhang, W. Zhang, S. L. Bromley, and J. Ye (2014), “An optical lattice clock with accuracy and stability at the 10^{-18} level,” *Nature* **506**, 71–75.
- Blume-Kohout, R (2010), “Optimal, reliable estimation of quantum states,” *New J. Phys.* **12** (4), 043034.
- Bodet, C, J. Estève, M. K. Oberthaler, and T. Gasenzer (2010), “Two-mode Bose gas: Beyond classical squeezing,” *Phys. Rev. A* **81**, 063605.
- Böhi, P, M. F. Riedel, J. Hoffrogge, J. Reichel, T. W. Hänsch, and P. Treutlein (2009), “Coherent manipulation of Bose-Einstein condensates with state-dependent microwave potentials on an atom chip,” *Nat. Phys.* **5** (8), 592–597.
- Bohnet, J G, K. C. Cox, M. A. Norcia, J. M. Weiner, Z. Chen, and J. K. Thompson (2014), “Reduced spin measurement back-action for a phase sensitivity ten times beyond the standard quantum limit,” *Nat. Photon.* **8** (9), 731–736.
- Bohnet, J G, B. C. Sawyer, J. W. Britton, M. L. Wall, A. M. Rey, M. Foss-Feig, and J. J. Bollinger (2016), “Quantum spin dynamics and entanglement generation with hundreds of trapped ions,” *Science* **352** (6291), 1297–1301.
- Boixo, S, A. Datta, M. J. Davis, A. Shaji, A. B. Tacla, and C. M. Caves (2009), “Quantum-limited metrology and Bose-Einstein condensates,” *Phys. Rev. A* **80**, 032103.
- Boixo, S, A. Datta, S. T. Flammia, A. Shaji, E. Bagan, and C. M. Caves (2008), “Quantum-limited metrology with product states,” *Phys. Rev. A* **77**, 012317.
- Boixo, S, S. T. Flammia, C. M. Caves, and J. M. Geremia (2007), “Generalized limits for single-parameter quantum estimation,” *Phys. Rev. Lett.* **98**, 090401.
- Bollinger, J J, Wayne M. Itano, D. J. Wineland, and D. J. Heinzen (1996), “Optimal frequency measurements with maximally correlated states,” *Phys. Rev. A* **54**, R4649–R4652.
- Bonneau, M, J. Ruauadel, R. Lopes, J.-C. Jaskula, A. Aspect, D. Boiron, and C. I. Westbrook (2013), “Tunable source of correlated atom beams,” *Phys. Rev. A* **87**, 061603.
- Bookjans, E M, C. D. Hamley, and M. S. Chapman (2011), “Strong quantum spin correlations observed in atomic spin mixing,” *Phys. Rev. Lett.* **107**, 210406.
- Bordè, Ch J (2002), “Atomic clocks and inertial sensors,” *Metrologia* **39** (5), 435–463.
- Borregaard, J, and A. S. Sørensen (2013), “Near-Heisenberg-limited atomic clocks in the presence of decoherence,” *Phys. Rev. Lett.* **111**, 090801.
- Botet, R, R. Jullien, and P. Pfeuty (1982), “Size scaling for infinitely coordinated systems,” *Phys. Rev. Lett.* **49**, 478–481.
- Boto, A N, P. Kok, D. S. Abrams, S. L. Braunstein, C. P. Williams, and J. P. Dowling (2000), “Quantum interferometric optical lithography: Exploiting entanglement to beat the diffraction limit,” *Phys. Rev. Lett.* **85**, 2733–2736.
- Bouchoule, I, and K. Mølmer (2002a), “Preparation of spin-squeezed atomic states by optical-phase-shift measurement,” *Phys. Rev. A* **66**, 043811.
- Bouchoule, I, and K. Mølmer (2002b), “Spin squeezing of atoms by the dipole interaction in virtually excited Rydberg states,” *Phys. Rev. A* **65**, 041803.
- Braginsky, V B, and F. Y. Khalili (1992), *Quantum Measurement* (Cambridge University Press, Cambridge).
- Braginsky, V B, Y. I. Vorontsov, and K. S. Thorne (1980), “Quantum nondemolition measurements,” *Science* **209** (4456), 547–557.
- Braunstein, S L, and C. M. Caves (1994), “Statistical distance and the geometry of quantum states,” *Phys. Rev. Lett.* **72**, 3439–3443.
- Braunstein, S L, C. M. Caves, and G. J. Milburn (1996), “Generalized uncertainty relations: Theory, examples, and Lorentz invariance,” *Ann. Phys.* **247** (1), 135–173.
- Braunstein, S L, and P. van Loock (2005), “Quantum information with continuous variables,” *Rev. Mod. Phys.* **77**, 513–577.
- Breitenbach, G, S. Schiller, and J. Mlynek (1997), “Measurement of the quantum states of squeezed light,” *Nature* **387** (6632), 471–475.
- Brif, C, and A. Mann (1996), “Nonclassical interferometry with intelligent light,” *Phys. Rev. A* **54**, 4505–4518.
- Britton, J W, B. C. Sawyer, A. C. Keith, C. C. J. Wang, J. K. Freericks, H. Uys, M. J. Biercuk, and J. J. Bollinger (2012), “Engineered two-dimensional Ising interactions in a trapped-ion quantum simulator with hundreds of spins,” *Nature* **484**, 489–492.
- Browaeys, A, D. Barredo, and T. Lahaye (2016), “Experimental investigations of dipole-dipole interactions between a few Rydberg atoms,” *J. Phys. B: At. Mol. Opt. Phys.* **49** (15), 152001.
- Brunner, N, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner (2014), “Bell nonlocality,” *Rev. Mod. Phys.* **86**, 419–478.
- Bücker, R, J. Grond, S. Manz, T. Berrada, T. Betz, C. Koller, U. Hohenester, T. Schumm, A. Perrin, and J. Schmiedmayer (2011), “Twin-atom beams,” *Nat. Phys.* **7**, 1608–611.
- Budker, D, W. Gawlik, D. F. Kimball, S. M. Rochester, V. V. Yashchuk, and A. Weis (2002), “Resonant nonlinear magneto-optical effects in atoms,” *Rev. Mod. Phys.* **74**, 1153–1201.
- Budker, D, D. F. Kimball, S. M. Rochester, and V. V. Yashchuk (2000), “Nonlinear magneto-optical rotation via alignment-to-orientation conversion,” *Phys. Rev. Lett.* **85**,

- 2088–2091.
- Budker, D., and D. F. J. (eds.) Kimball (2013), *Optical Magnetometry* (Cambridge Univ. Press).
- Budker, D., and M. V. Romalis (2007), “Optical magnetometry,” *Nat. Phys.* **3**, 227.
- Buonsante, P., R. Burioni, E. Vescovi, and A. Vezzani (2012), “Quantum criticality in a bosonic Josephson junction,” *Phys. Rev. A* **85**, 043625.
- Bures, D (1969), “An extension of Kakutani’s theorem on infinite product measures to the tensor product of semifinite w^* -algebras,” *Trans. Amer. Math. Soc.* **135**, 199.
- Bužek, V., R. Derka, and S. Massar (1999), “Optimal quantum clocks,” *Phys. Rev. Lett.* **82**, 2207–2210.
- Byrnes, T (2013), “Fractality and macroscopic entanglement in two-component Bose-Einstein condensates,” *Phys. Rev. A* **88**, 023609.
- Byrnes, T, K. Wen, and Y. Yamamoto (2012), “Macroscopic quantum computation using Bose-Einstein condensates,” *Phys. Rev. A* **85**, 040306.
- Campos, R A, Christopher C. Gerry, and A. Benmoussa (2003), “Optical interferometry at the Heisenberg limit with twin Fock states and parity measurements,” *Phys. Rev. A* **68**, 023810.
- Carnio, E G, A. Buchleitner, and M. Gessner (2015), “Robust asymptotic entanglement under multipartite collective dephasing,” *Phys. Rev. Lett.* **115**, 010404.
- Carruthers, P., and M. M. Nieto (1968), “Phase and angle variables in quantum mechanics,” *Rev. Mod. Phys.* **40**, 411–440.
- Casabone, B., A. Stute, K. Friebe, B. Brandstätter, K. Schüppert, R. Blatt, and T. E. Northup (2013), “Heralded entanglement of two ions in an optical cavity,” *Phys. Rev. Lett.* **111**, 100505.
- Castin, Y., and J. Dalibard (1997), “Relative phase of two Bose-Einstein condensates,” *Phys. Rev. A* **55**, 4330–4337.
- Caves, C M (1981), “Quantum-mechanical noise in an interferometer,” *Phys. Rev. D* **23**, 1693–1708.
- Chang, M-S, C. D. Hamley, M. D. Barrett, J. A. Sauer, K. M. Fortier, W. Zhang, L. You, and M. S. Chapman (2004), “Observation of spinor dynamics in optically trapped ^{87}Rb Bose-Einstein condensates,” *Phys. Rev. Lett.* **92**, 140403.
- Chase, B A, B. Q. Baragiola, H. L. Partner, B. D. Black, and J. M. Geremia (2009), “Magnetometry via a double-pass continuous quantum measurement of atomic spin,” *Phys. Rev. A* **79**, 062107.
- Chaudhury, S, G. A. Smith, K. Schulz, and P. S. Jessen (2006), “Continuous nondemolition measurement of the Cs clock transition pseudospin,” *Phys. Rev. Lett.* **96**, 043001.
- Chaves, R, J. B. Brask, M. Markiewicz, J. Kołodyński, and A. Acín (2013), “Noisy metrology beyond the standard quantum limit,” *Phys. Rev. Lett.* **111**, 120401.
- Chen, W, J. Hu, Y. Duan, B. Braverman, H. Zhang, and V. Vuletić (2015), “Carving complex many-atom entangled states by single-photon detection,” *Phys. Rev. Lett.* **115**, 250502.
- Chen, Z, J. G. Bohnet, S. R. Sankar, J. Dai, and J. K. Thompson (2011), “Conditional spin squeezing of a large ensemble via the vacuum Rabi splitting,” *Phys. Rev. Lett.* **106**, 133601.
- Chen, Z, J. G. Bohnet, J. M. Weiner, K. C. Cox, and J. K. Thompson (2014), “Cavity-aided nondemolition measurements for atom counting and spin squeezing,” *Phys. Rev. A* **89**, 043837.
- Chin, A W, S. F. Huelga, and M. B. Plenio (2012), “Quantum metrology in non-Markovian environments,” *Phys. Rev. Lett.* **109**, 233601.
- Chin, C, R. Grimm, P. Julienne, and E. Tiesinga (2010), “Feshbach resonances in ultracold gases,” *Rev. Mod. Phys.* **82**, 1225–1286.
- Choi, S., and N. P. Bigelow (2005), “Quantum squeezing and entanglement in a two-mode Bose-Einstein condensate with time-dependent Josephson-like coupling,” *Phys. Rev. A* **72**, 033612.
- Choi, S., and B. Sundaram (2008), “Bose-Einstein condensate as a nonlinear Ramsey interferometer operating beyond the Heisenberg limit,” *Phys. Rev. A* **77**, 053613.
- Chou, C W, D. B. Hume, M. J. Thorpe, D. J. Wineland, and T. Rosenband (2011), “Quantum coherence between two atoms beyond $Q = 10^{15}$,” *Phys. Rev. Lett.* **106**, 160801.
- Christensen, S L, J.-B. Béguin, E. Bookjans, H. L. Sørensen, J. H. Müller, J. Appel, and E. S. Polzik (2014), “Quantum interference of a single spin excitation with a macroscopic atomic ensemble,” *Phys. Rev. A* **89**, 033801.
- Christensen, S L, J.-B. Béguin, H. L. Sørensen, E. Bookjans, D. Oblak, J. H. Müller, J. Appel, and E. S. Polzik (2013), “Toward quantum state tomography of a single polariton state of an atomic ensemble,” *New J. Phys.* **15**, 015002.
- Cirac, J I, M. Lewenstein, K. Mølmer, and P. Zoller (1998), “Quantum superposition states of Bose-Einstein condensates,” *Phys. Rev. A* **57**, 1208–1218.
- Cirac, J I, and P. Zoller (1995), “Quantum computations with cold trapped ions,” *Phys. Rev. Lett.* **74**, 4091–4094.
- Cohen, M (1968), “The Fisher information and convexity,” *IEEE Transactions on Information Theory* **14**, 591–592.
- Cox, K C, G. P. Greve, J. M. Weiner, and J. K. Thompson (2016), “Deterministic squeezed states with collective measurements and feedback,” *Phys. Rev. Lett.* **116**, 093602.
- Cramér, H (1946), *Mathematical Methods of Statistics* (Princeton Univ. Press, Princeton, NJ).
- Cronin, A D, J. Schmiedmayer, and D. E. Pritchard (2009), “Optics and interferometry with atoms and molecules,” *Rev. Mod. Phys.* **81**, 1051–1129.
- Davis, E, G. Bentsen, and M. Schleier-Smith (2016), “Approaching the Heisenberg limit without single-particle detection,” *Phys. Rev. Lett.* **116**, 053601.
- Demkowicz-Dobrzański, R, J. Kołodyński, and M. Guţă (2012), “The elusive Heisenberg limit in quantum-enhanced metrology,” *Nat. Comm.* **3**, 1063.
- Demkowicz-Dobrzański, R., and L. Maccone (2014), “Using entanglement against noise in quantum metrology,” *Phys. Rev. Lett.* **113**, 250801.
- Dicke, R H (1954), “Coherence in spontaneous radiation processes,” *Phys. Rev.* **93**, 99–110.
- Dorner, U (2012), “Quantum frequency estimation with trapped ions and atoms,” *New J. Phys.* **14** (4), 043011.
- Dorner, U, R. Demkowicz-Dobrzański, B. J. Smith, J. S. Lundeen, W. Wasilewski, K. Banaszek, and I. A. Walmsley (2009), “Optimal quantum phase estimation,” *Phys. Rev. Lett.* **102**, 040403.
- Dowling, J P (1998), “Correlated input-port, matter-wave interferometer: Quantum-noise limits to the atom-laser gyroscope,” *Phys. Rev. A* **57**, 4736–4746.
- Dowling, J P, G. S. Agarwal, and W. P. Schleich (1994), “Wigner distribution of a general angular-momentum state: Applications to a collection of two-level atoms,” *Phys. Rev. A* **49**, 4101–4109.

- Duan, L-M (2011), “Entanglement detection in the vicinity of arbitrary Dicke states,” *Phys. Rev. Lett.* **107**, 180502.
- Duan, L-M, J. I. Cirac, and P. Zoller (2002), “Quantum entanglement in spinor Bose-Einstein condensates,” *Phys. Rev. A* **65**, 033619.
- Duan, L-M, G. Giedke, J. I. Cirac, and P. Zoller (2000a), “Inseparability criterion for continuous variable systems,” *Phys. Rev. Lett.* **84**, 2722–2725.
- Duan, L-M, M. D. Lukin, J. I. Cirac, and P. Zoller (2001), “Long-distance quantum communication with atomic ensembles and linear optics,” *Nature* **414**, 413.
- Duan, L-M, and C. Monroe (2010), “*Colloquium*: Quantum networks with trapped ions,” *Rev. Mod. Phys.* **82**, 1209–1224.
- Duan, L-M, A. Sørensen, J. I. Cirac, and P. Zoller (2000b), “Squeezing and entanglement of atomic beams,” *Phys. Rev. Lett.* **85**, 3991–3994.
- Dunningham, J A, K. Burnett, and Stephen M. Barnett (2002), “Interferometry below the standard quantum limit with Bose-Einstein condensates,” *Phys. Rev. Lett.* **89**, 150401.
- Dür, W, M. Skotiniotis, F. Fröwis, and B. Kraus (2014), “Improved quantum metrology using quantum error correction,” *Phys. Rev. Lett.* **112**, 080801.
- Dür, W, G. Vidal, and J. I. Cirac (2000), “Three qubits can be entangled in two inequivalent ways,” *Phys. Rev. A* **62**, 062314.
- Dusuel, S, and J. Vidal (2005), “Continuous unitary transformations and finite-size scaling exponents in the Lipkin-Meshkov-Glick model,” *Phys. Rev. B* **71**, 224420.
- de Echaniz, S R, M. W. Mitchell, M. Kubasik, M. Koschorreck, H. Crepaz, J. Eschner, and E. S. Polzik (2005), “Conditions for spin squeezing in a cold ^{87}Rb ensemble,” *J. Opt. B: Quantum and Semiclassical Optics* **7** (12), S548.
- Eibl, M, S. Gaertner, M. Bourennane, C. Kurtsiefer, M. Żukowski, and H. Weinfurter (2003), “Experimental observation of four-photon entanglement from parametric down-conversion,” *Phys. Rev. Lett.* **90**, 200403.
- Einstein, A, B. Podolsky, and N. Rosen (1935), “Can quantum-mechanical description of physical reality be considered complete?” *Phys. Rev.* **47**, 777–780.
- Escher, B M, R. L. de Matos Filho, and L. Davidovich (2011), “General framework for estimating the ultimate precision limit in noisy quantum-enhanced metrology,” *Nat. Phys.* **7**, 406.
- Estève, J, C. Gross, A. Weller, S. Giovanazzi, and M. K. Oberthaler (2008), “Squeezing and entanglement in a Bose-Einstein condensate,” *Nature* **455**, 1216–1219.
- Eto, Y, H. Ikeda, H. Suzuki, S. Hasegawa, Y. Tomiyama, S. Sekine, M. Sadgrove, and T. Hirano (2013), “Spin-echo-based magnetometry with spinor Bose-Einstein condensates,” *Phys. Rev. A* **88**, 031602.
- Facchi, P, and S. Pascazio (2008), “Quantum Zeno dynamics: mathematical and physical aspects,” *J. Phys. A: Math. Theor.* **41** (49), 493001.
- Ferrini, G, A. Minguzzi, and F. W. J. Hekking (2008), “Number squeezing, quantum fluctuations, and oscillations in mesoscopic Bose Josephson junctions,” *Phys. Rev. A* **78**, 023606.
- Ferrini, G, D. Spehner, A. Minguzzi, and F. W. J. Hekking (2011), “Effect of phase noise on quantum correlations in Bose-Josephson junctions,” *Phys. Rev. A* **84**, 043628.
- Ferris, A J, M. K. Olsen, E. G. Cavalcanti, and M. J. Davis (2008), “Detection of continuous variable entanglement without coherent local oscillators,” *Phys. Rev. A* **78**, 060104.
- Feynman, R A, F. L. Jr. Vernon, and R. W. Hellwarth (1957), “Geometrical representation of the Schrödinger equation for solving maser problems,” *J. App. Phys.* **28**, 69.
- Fisher, R A (1922), “On the mathematical foundations of theoretical statistics,” *Phil. Trans. R. Soc. Lond. Ser.* **222** (602), 309–368.
- Fisher, R A (1925), “Theory of statistical estimation,” *Math. Proc. Camb. Phil. Soc.* **22**, 700.
- Fleischhauer, M, and M. D. Lukin (2000), “Dark-state polaritons in electromagnetically induced transparency,” *Phys. Rev. Lett.* **84**, 5094–5097.
- Foss-Feig, M, K. R. A. Hazzard, J. J. Bollinger, and A. M. Rey (2013), “Nonequilibrium dynamics of arbitrary-range Ising models with decoherence: An exact analytic solution,” *Phys. Rev. A* **87**, 042101.
- Fréchet, M (1943), “Sur l’extension de certaines évaluations statistiques au cas de petits échantillons,” *Revue de l’Institut International de Statistique* **11** (3/4), 182–205.
- Fujiwara, A (2001), “Quantum channel identification problem,” *Phys. Rev. A* **63**, 042304.
- Gabbriellini, M, L. Pezzè, and A. Smerzi (2015), “Spin-mixing interferometry with Bose-Einstein condensates,” *Phys. Rev. Lett.* **115**, 163002.
- García-Ripoll, J J, P. Zoller, and J. I. Cirac (2005), “Coherent control of trapped ions using off-resonant lasers,” *Phys. Rev. A* **71**, 062309.
- Gati, R, B. Hemmerling, J. Fölling, M. Albiez, and M. K. Oberthaler (2006), “Noise thermometry with two weakly coupled Bose-Einstein condensates,” *Phys. Rev. Lett.* **96**, 130404.
- Gerbier, F, A. Widera, S. Fölling, O. Mandel, and I. Bloch (2006), “Resonant control of spin dynamics in ultracold quantum gases by microwave dressing,” *Phys. Rev. A* **73**, 041602.
- Geremia, J M, J. K. Stockton, A. C. Doherty, and H. Mabuchi (2003), “Quantum Kalman filtering and the Heisenberg limit in atomic magnetometry,” *Phys. Rev. Lett.* **91**, 250801.
- Gerry, C C, R. A. Campos, and A. Benmoussa (2004), “Comment on “interferometric detection of optical phase shifts at the Heisenberg limit”,” *Phys. Rev. Lett.* **92**, 209301.
- Gil, L I R, R. Mukherjee, E. M. Bridge, M. P. A. Jones, and T. Pohl (2014), “Spin squeezing in a Rydberg lattice clock,” *Phys. Rev. Lett.* **112**, 103601.
- Gill, R D (2008), “Conciliation of Bayes and pointwise quantum state estimation,” in *Quantum Stochastics and Information: Statistics, Filtering and Control*, edited by V.P. Belavkin & M. Guță (World Scientific) p. 239.
- Gilmore, R, and D. H. Feng (1978), “Phase transitions in nuclear matter described by pseudospin Hamiltonians,” *Nucl. Phys. A* **301**, 189.
- Giovannetti, V, S. Lloyd, and L. Maccone (2004), “Quantum-enhanced measurements: Beating the standard quantum limit,” *Science* **306**, 1330.
- Giovannetti, V, S. Lloyd, and L. Maccone (2006), “Quantum metrology,” *Phys. Rev. Lett.* **96**, 010401.
- Giovannetti, V, S. Lloyd, and L. Maccone (2011), “Advances in quantum metrology,” *Nat. Photon.* **5**, 222.
- Giovannetti, V, S. Mancini, D. Vitali, and P. Tombesi (2003), “Characterizing the entanglement of bipartite quantum systems,” *Phys. Rev. A* **67**, 022320.

- Glauber, R J (1963), “Coherent and incoherent states of the radiation field,” *Phys. Rev.* **131**, 2766–2788.
- Glick, A J, H. J. Lipkin, and N. Meshkov (1965), “Validity of many-body approximation methods for a solvable model: (iii). diagram summations,” *Nucl. Phys.* **62**, 211.
- Goldstein, E V, and P. Meystre (1999), “Quantum theory of atomic four-wave mixing in Bose-Einstein condensates,” *Phys. Rev. A* **59**, 3896–3901.
- Gordon, D, and C. M. Savage (1999), “Creating macroscopic quantum superpositions with Bose-Einstein condensates,” *Phys. Rev. A* **59**, 4623–4629.
- Grangier, P, J. A. Levenson, and J.-P. Poizat (1998), “Quantum non-demolition measurements in optics,” *Nature* **396**, 537.
- Greenberger, D M, M. A. Horne, A. Shimony, and A. Zeilinger (1990), “Bell’s theorem without inequalities,” *Am. J. Phys.* **58**, 1131.
- Grond, J, J. Schmiedmayer, and U. Hohenester (2009a), “Optimizing number squeezing when splitting a mesoscopic condensate,” *Phys. Rev. A* **79**, 021603.
- Grond, J, G. von Winckel, J. Schmiedmayer, and U. Hohenester (2009b), “Optimal control of number squeezing in trapped Bose-Einstein condensates,” *Phys. Rev. A* **80**, 053625.
- Gross, C, H. Strobel, E. Nicklas, T. Zibold, N. Bar-Gill, G. Kurizki, and M. K. Oberthaler (2011), “Atomic homodyne detection of continuous-variable entangled twin-atom states,” *Nature* **480**, 219–223.
- Gross, C, T Zibold, E Nicklas, J Estève, and M.K. Oberthaler (2010), “Nonlinear atom interferometer surpasses classical precision limit,” *Nature* **464**, 1165–1169.
- Gühne, O, and G. Tóth (2009), “Entanglement detection,” *Phys. Rep.* **474**, 1.
- Gühne, O, G. Tóth, and H.J. Briegel (2005), “Multipartite entanglement in spin chains,” *New J. Phys.* **7**, 1.
- Haas, F, J. Volz, R. Gehr, J. Reichel, and J. Estève (2014), “Entangled states of more than 40 atoms in an optical fiber cavity,” *Science* **344**, 180.
- Hadzibabic, Z, P. Krüger, M. Cheneau, B. Battelier, and J. Dalibard (2006), “Berezinskii-Kosterlitz-Thouless crossover in a trapped atomic gas,” *Nature* **441** (7097), 1118–1121.
- Häffner, H, W. Hänsel, C. F. Roos, J. Benhelm, D. Chek-al kar, M. Chwalla, T. Körber, U. D. Rapol, M. Riebe, P. O. Schmidt, C. Becher, O. Gühne, W. Dür, and R. Blatt (2005), “Scalable multiparticle entanglement of trapped ions,” *Nature* **438** (7068), 643–646.
- Häffner, H, C. F. Roos, and R. Blatt (2008), “Quantum computing with trapped ions,” *Phys. Rep.* **469** (4), 155–203.
- Haine, S A, and S. S. Szigeti (2015), “Quantum metrology with mixed states: When recovering lost information is better than never losing it,” *Phys. Rev. A* **92**, 032317.
- Haine, S A, S. S. Szigeti, M. D. Lang, and C. M. Caves (2015), “Heisenberg-limited metrology with information recycling,” *Phys. Rev. A* **91**, 041802.
- Hald, J, J. L. Sørensen, C. Schori, and E. S. Polzik (1999), “Spin squeezed atoms: A macroscopic entangled ensemble created by light,” *Phys. Rev. Lett.* **83**, 1319–1322.
- Hald, J, J. L. Sørensen, C. Schori, and E. S. Polzik (2000), “Entanglement transfer from light to atoms,” *J. Mod. Opt.* **47**, 2599.
- Haljan, P C, P. J. Lee, K.-A. Brickman, M. Acton, L. Deslauriers, and C. Monroe (2005), “Entanglement of trapped-ion clock states,” *Phys. Rev. A* **72**, 062316.
- Hall, B V, S. Whitlock, R. Anderson, P. Hannaford, and A. I. Sidorov (2007), “Condensate splitting in an asymmetric double well for atom chip based sensors,” *Phys. Rev. Lett.* **98**, 030402.
- Hall, D S, M. R. Matthews, J. R. Ensher, C. E. Wieman, and E. A. Cornell (1998a), “Dynamics of component separation in a binary mixture of Bose-Einstein condensates,” *Phys. Rev. Lett.* **81**, 1539–1542.
- Hall, D S, M. R. Matthews, C. E. Wieman, and E. A. Cornell (1998b), “Measurements of relative phase in two-component Bose-Einstein condensates,” *Phys. Rev. Lett.* **81**, 1543–1546.
- Hamley, C D, C. S. Gerving, T. M. Hoang, E. M. Bookjans, and M. S. Chapman (2012), “Spin-nematic squeezed vacuum in a quantum gas,” *Nat. Phys.* **8**, 305.
- Hammerer, K, K. Mølmer, E. S. Polzik, and J. I. Cirac (2004), “Light-matter quantum interface,” *Phys. Rev. A* **70**, 044304.
- Hammerer, K, A. S. Sørensen, and E. S. Polzik (2010), “Quantum interface between light and atomic ensembles,” *Rev. Mod. Phys.* **82**, 1041–1093.
- Harty, T P, D. T. C. Allcock, C. J. Ballance, L. Guidoni, H. A. Janacek, N. M. Linke, D. N. Stacey, and D. M. Lucas (2014), “High-fidelity preparation, gates, memory, and readout of a trapped-ion quantum bit,” *Phys. Rev. Lett.* **113**, 220501.
- Hauke, Philipp, Markus Heyl, Luca Tagliacozzo, and Peter Zoller (2016), “Measuring multipartite entanglement through dynamic susceptibilities,” *Nat. Phys.* **12** (8), 778–782.
- Hayashi, M (Ed) (2005), *Asymptotic Theory of Quantum Statistical Inference* (World Scientific Singapore).
- He, Q Y, M. D. Reid, T. G. Vaughan, C. Gross, M. Oberthaler, and P. D. Drummond (2011), “Einstein-Podolsky-Rosen entanglement strategies in two-well Bose-Einstein condensates,” *Phys. Rev. Lett.* **106**, 120405.
- Helstrom, C W (1967), “Minimum mean-squared error of estimates in quantum statistics,” *Phys. Lett. A* **25** (2), 101.
- Helstrom, C W (1976), *Quantum Detection and Estimation Theory* (Academic Press, New York).
- Henkel, N, R. Nath, and T. Pohl (2010), “Three-dimensional roton excitations and supersolid formation in rydberg-excited bose-einstein condensates,” *Phys. Rev. Lett.* **104**, 195302.
- Hensen, B, H. Bernien, A. E. Dreau, A. Reiserer, N. Kalb, M. S. Blok, J. Ruitenbergh, R. F. L. Vermeulen, R. N. Schouten, C. Abellan, W. Amaya, V. Pruneri, M. W. Mitchell, M. Markham, D. J. Twitchen, D. Elkouss, S. Wehner, T. H. Taminiau, and R. Hanson (2015), “Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres,” *Nature* **526** (7575), 682–686.
- Herrera-Martí, D A, T. Gefen, D. Aharonov, N. Katz, and A. Retzker (2015), “Quantum error-correction-enhanced magnetometer overcoming the limit imposed by relaxation,” *Phys. Rev. Lett.* **115**, 200501.
- Hill, S, and W. K. Wootters (1997), “Entanglement of a pair of quantum bits,” *Phys. Rev. Lett.* **78**, 5022–5025.
- Hillery, M, and L. Mlodinow (1993), “Interferometers and minimum-uncertainty states,” *Phys. Rev. A* **48**, 1548–1558.
- Hinkley, N, J. A. Sherman, N. B. Phillips, M. Schioppo, N. D. Lemke, K. Beloy, M. Pizzocaro, C. W. Oates, and A. D. Ludlow (2013), “An atomic clock with 10^{-18} instability,” *Science* **341**, 1215.

- Ho, T-L (1998), “Spinor Bose condensates in optical traps,” *Phys. Rev. Lett.* **81**, 742–745.
- Ho, T L, and C. V. Ciobanu (2004), “The Schrödinger cat family in attractive Bose gases,” *J. Low Temp. Phys.* **125**, 257.
- Hofferberth, S, I. Lesanovsky, B. Fischer, J. Verdu, and J. Schmiedmayer (2006), “Radiofrequency-dressed-state potentials for neutral atoms,” *Nat. Phys.* **2**, 710.
- Hofmann, J, M. Krug, N. Ortegel, L. Gérard, M. Weber, W. Rosenfeld, and H. Weinfurter (2012), “Heralded entanglement between widely separated atoms,” *Science* **337** (6090), 72–75.
- Holevo, A S (1982), *Probabilistic and statistical aspects of quantum theory* (North-Holland Publishing Company, Amsterdam).
- Holland, M J, and K. Burnett (1993), “Interferometric detection of optical phase shifts at the Heisenberg limit,” *Phys. Rev. Lett.* **71**, 1355–1358.
- Home, J P, M. J. McDonnell, D. L. Lucas, G. Imreh, B. C. Keitch, D. J. Szwer, N. R. Thomas, S. C. Webster, N. D. Stacey, and A. M. Steane (2006), “Deterministic entanglement and tomography of ion-spin qubits,” *New J. Phys.* **8** (9), 188.
- Horodecki, R, P. Horodecki, M. Horodecki, and K. Horodecki (2009), “Quantum entanglement,” *Rev. Mod. Phys.* **81**, 865–942.
- Horrom, T, R. Singh, J. P. Dowling, and E. E. Mikhailov (2012), “Quantum-enhanced magnetometer with low-frequency squeezing,” *Phys. Rev. A* **86**, 023803.
- Hosten, O, N. J. Engelsens, R. Krishnakumar, and M. A. Kasevich (2016a), “Measurement noise 100 times lower than the quantum-projection limit using entangled atoms,” *Nature* **529** (7587), 505–508.
- Hosten, O, R. Krishnakumar, N. J. Engelsens, and M. A. Kasevich (2016b), “Quantum phase magnification,” *Science* **352** (6293), 1552–1555.
- Hu, Jiazhong, Zachary Vendeiro, Wenlan Chen, Hao Zhang, Robert McConnell, Anders Sørensen, and Vladan Vuletić (2016), “Violation of classical physics by a mesoscopic system,” [arXiv:1606.05392 \[quant-ph\]](https://arxiv.org/abs/1606.05392).
- Huang, Y P, and M. G. Moore (2006), “Creation, detection, and decoherence of macroscopic quantum superposition states in double-well Bose-Einstein condensates,” *Phys. Rev. A* **73**, 023606.
- Huang, Y P, and M. G. Moore (2008), “Optimized double-well quantum interferometry with Gaussian squeezed states,” *Phys. Rev. Lett.* **100**, 250406.
- Huang, Z, C. Macchiavello, and L. Maccone (2016), “Usefulness of entanglement-assisted quantum metrology,” *Phys. Rev. A* **94**, 012101.
- Hübner, M (1992), “Explicit computation of the Bures distance for density matrices,” *Phys. Lett. A* **163** (4), 239.
- Hudelist, F, J. Kong, C. Liu, J. Jing, Z. Y. Ou, and W. Zhang (2014), “Quantum metrology with parametric amplifier-based photon correlation interferometers,” *Nat. Comm.* **5**, 3049.
- Huelga, S F, C. Macchiavello, T. Pellizzari, A. K. Ekert, M. B. Plenio, and J. I. Cirac (1997), “Improvement of frequency standards with quantum entanglement,” *Phys. Rev. Lett.* **79**, 3865–3868.
- Hume, D B, I. Stroescu, M. Joos, W. Muessel, H. Strobel, and M. K. Oberthaler (2013), “Accurate atom counting in mesoscopic ensembles,” *Phys. Rev. Lett.* **111**, 253001.
- Husimi, K (1940), “Some formal properties of the density matrix,” *Proc. Phys.-Math. Soc. Jap.* **22** (4), 264–314.
- Huuer, S D, C. F. Wildfeuer, and J. P. Dowling (2008), “Entangled Fock states for robust quantum optical metrology, imaging, and sensing,” *Phys. Rev. A* **78**, 063828.
- Hyllus, P, O. Gühne, and A. Smerzi (2010), “Not all pure entangled states are useful for sub-shot-noise interferometry,” *Phys. Rev. A* **82**, 012337.
- Hyllus, P, W. Laskowski, R. Krischek, C. Schwemmer, W. Wiczeorek, H. Weinfurter, L. Pezzè, and A. Smerzi (2012a), “Fisher information and multiparticle entanglement,” *Phys. Rev. A* **85**, 022321.
- Hyllus, P, L. Pezzè, A. Smerzi, and G. Tóth (2012b), “Entanglement and extreme spin squeezing for a fluctuating number of indistinguishable particles,” *Phys. Rev. A* **86**, 012337.
- Inguscio, M, and L. Fallani (2013), *Atomic Physics: Precise Measurements and Ultracold Matter* (Oxford University Press).
- Inoue, R, S.-I.-R. Tanaka, R. Namiki, T. Sagawa, and Y. Takahashi (2013), “Unconditional quantum-noise suppression via measurement-based quantum feedback,” *Phys. Rev. Lett.* **110**, 163602.
- Isella, L, and J. Ruostekoski (2005), “Nonadiabatic dynamics of a Bose-Einstein condensate in an optical lattice,” *Phys. Rev. A* **72**, 011601.
- Islam, R, C. Senko, W. C. Campbell, S. Korenblit, J. Smith, A. Lee, E. E. Edwards, C.-C. J. Wang, J. K. Freericks, and C. Monroe (2013), “Emergence and frustration of magnetism with variable-range interactions in a quantum simulator,” *Science* **340** (6132), 583–587.
- Itano, W M, J. C. Bergquist, J. J. Bollinger, J. M. Gilligan, D. J. Heinzen, F. L. Moore, M. G. Raizen, and D. J. Wineland (1993), “Quantum projection noise: Population fluctuations in two-level systems,” *Phys. Rev. A* **47**, 3554–3570.
- Jääskeläinen, M, W. Zhang, and P. Meystre (2004), “Limits to phase resolution in matter-wave interferometry,” *Phys. Rev. A* **70**, 063612.
- Jaskula, J-C, M. Bonneau, G. B. Partridge, V. Krachmalnicoff, P. Deuar, K. V. Kheruntsyan, A. Aspect, D. Boiron, and C. I. Westbrook (2010), “Sub-Poissonian number differences in four-wave mixing of matter waves,” *Phys. Rev. Lett.* **105**, 190402.
- Javanainen, J (1986), “Oscillatory exchange of atoms between traps containing Bose condensates,” *Phys. Rev. Lett.* **57**, 3164–3166.
- Javanainen, J, and M. Yu. Ivanov (1999), “Splitting a trap containing a Bose-Einstein condensate: Atom number fluctuations,” *Phys. Rev. A* **60**, 2351–2359.
- Javanainen, J, and M. Wilkens (1997), “Phase and phase diffusion of a split Bose-Einstein condensate,” *Phys. Rev. Lett.* **78**, 4675–4678.
- Jeske, J, J. H. Cole, and S. F. Huelga (2014), “Quantum metrology subject to spatially correlated Markovian noise: restoring the Heisenberg limit,” *New J. Phys.* **16** (7), 073039.
- Jo, G-B, J.-H. Choi, C. A. Christensen, T. A. Pasquini, Y.-R. Lee, W. Ketterle, and D. E. Pritchard (2007a), “Phase-sensitive recombination of two Bose-Einstein condensates on an atom chip,” *Phys. Rev. Lett.* **98**, 180401.
- Jo, G-B, Y. Shin, S. Will, T. A. Pasquini, M. Saba, W. Ketterle, D. E. Pritchard, M. Vengalattore, and M. Prentiss (2007b), “Long phase coherence time and number squeez-

- ing of two Bose-Einstein condensates on an atom chip," *Phys. Rev. Lett.* **98**, 030407.
- Johanning, M, A. Braun, N. Timoney, V. Elman, W. Neuhauser, and C. Wunderlich (2009a), "Individual addressing of trapped ions and coupling of motional and spin states using rf radiation," *Phys. Rev. Lett.* **102**, 073004.
- Johanning, M, A. F. Varón, and C. Wunderlich (2009b), "Quantum simulations with cold trapped ions," *J. Phys. B: At. Mol. Opt. Phys.* **42** (15), 154009.
- Jozsa, R (1994), "Fidelity for mixed quantum states," *J. Mod. Opt.* **41**, 2315–2323.
- Juliá-Díaz, B, E. Torrontegui, J. Martorell, J. G. Muga, and A. Polls (2012), "Fast generation of spin-squeezed states in bosonic Josephson junctions," *Phys. Rev. A* **86**, 063623.
- Juliá-Díaz, B, T. Zibold, M. K. Oberthaler, M. Melé-Messeguer, J. Martorell, and A. Polls (2012), "Dynamic generation of spin-squeezed states in bosonic Josephson junctions," *Phys. Rev. A* **86**, 023615.
- Julsgaard, B, A. Kozhekin, and E. S. Polzik (2001), "Experimental long-lived entanglement of two macroscopic objects," *Nature* **413**, 400.
- Jurcevic, P, B. P. Lanyon, P. Hauke, C. Hempel, P. Zoller, R. Blatt, and C. F. Roos (2014), "Quasiparticle engineering and entanglement propagation in a quantum many-body system," *Nature* **511**, 202–205.
- Kacprowicz, M, R. Demkowicz-Dobrzański, W. Wasilewski, K. Banaszek, and I. A. Walmsley (2010), "Experimental quantum-enhanced estimation of a lossy phase shift," *Nat. Photon.* **4** (6), 357–360.
- Kawaguchi, Y, and M. Ueda (2012), "Spinor Bose-Einstein condensates," *Phys. Rep.* **520**, 253–381.
- Kay, S M (1993), *Fundamentals of Statistical Signal Processing, Volume I: Estimation Theory* (Prentice Hall, Upper Saddle River, NJ).
- Kessler, E M, P. Kómár, M. Bishof, L. Jiang, A. S. Sørensen, J. Ye, and M. D. Lukin (2014a), "Heisenberg-limited atom clocks based on entangled qubits," *Phys. Rev. Lett.* **112**, 190403.
- Kessler, E M, I. Lovchinsky, A. O. Sushkov, and M. D. Lukin (2014b), "Quantum error correction for metrology," *Phys. Rev. Lett.* **112**, 150802.
- Kiesel, T, W. Vogel, V. Parigi, A. Zavatta, and M. Bellini (2008), "Experimental determination of a nonclassical Glauber-Sudarshan P function," *Phys. Rev. A* **78**, 021804.
- Kim, K, M.-S. Chang, R. Islam, S. Korenblit, L.-M. Duan, and C. Monroe (2009), "Entanglement and tunable spin-spin couplings between trapped ions using multiple transverse modes," *Phys. Rev. Lett.* **103**, 120502.
- Kim, K, M. S. Chang, S. Korenblit, R. Islam, E. E. Edwards, J. K. Freericks, G. D. Lin, L. M. Duan, and C. Monroe (2010), "Quantum simulation of frustrated Ising spins with trapped ions," *Nature* **465** (7298), 590–593.
- Kim, T, O. Pfister, M. J. Holland, J. Noh, and J. L. Hall (1998), "Influence of decorrelation on Heisenberg-limited interferometry with quantum correlated photons," *Phys. Rev. A* **57**, 4004–4013.
- Kimble, H J (2008), "The quantum internet," *Nature* **453**, 1023.
- Kitagawa, M, and M. Ueda (1993), "Squeezed spin states," *Phys. Rev. A* **47**, 5138–5143.
- Kitching, J, S. Knappe, and E. A. Donley (2011), "Atomic sensors – a review," *IEEE Sensors Journal* **11** (9), 1749–1758.
- Klempt, C, O. Topic, G. Gebreyesus, M. Scherer, T. Henninger, P. Hyllus, W. Ertmer, L. Santos, and J. J. Arlt (2010), "Parametric amplification of vacuum fluctuations in a spinor condensate," *Phys. Rev. Lett.* **104**, 195303.
- Knysh, S, V. N. Smelyanskiy, and G. A. Durkin (2011), "Scaling laws for precision in quantum interferometry and the bifurcation landscape of the optimal state," *Phys. Rev. A* **83**, 021804.
- Kolodyński, J, and R. Demkowicz-Dobrzański (2013), "Efficient tools for quantum metrology with uncorrelated noise," *New J. Phys.* **15** (7), 073043.
- Kominis, I K (2008), "Sub-shot-noise magnetometry with a correlated spin-relaxation dominated alkali-metal vapor," *Phys. Rev. Lett.* **100**, 073002.
- Kominis, I K, T. W. Kornack, J. C. Allred, and M. V. Romalis (2003), "A subfemtotesla multichannel atomic magnetometer," *Nature* **422**, 596–599.
- Korbicz, J K, J. I. Cirac, and M. Lewenstein (2005), "Spin squeezing inequalities and entanglement of N qubit states," *Phys. Rev. Lett.* **95**, 120502.
- Korbicz, J K, O. Gühne, M. Lewenstein, H. Häffner, C. F. Roos, and R. Blatt (2006), "Generalized spin-squeezing inequalities in N -qubit systems: Theory and experiment," *Phys. Rev. A* **74**, 052319.
- Koschorreck, M, M. Napolitano, B. Dubost, and M. W. Mitchell (2010a), "Quantum nondemolition measurement of large-spin ensembles by dynamical decoupling," *Phys. Rev. Lett.* **105**, 093602.
- Koschorreck, M, M. Napolitano, B. Dubost, and M. W. Mitchell (2010b), "Sub-projection-noise sensitivity in broadband atomic magnetometry," *Phys. Rev. Lett.* **104**, 093602.
- Kot, E, N. Grønbech-Jensen, B. M. Nielsen, J. S. Neergaard-Nielsen, E. S. Polzik, and A. S. Sørensen (2012), "Breakdown of the classical description of a local system," *Phys. Rev. Lett.* **108**, 233601.
- Krischek, R, C. Schwemmer, W. Wieczorek, H. Weinfurter, P. Hyllus, L. Pezzè, and A. Smerzi (2011), "Useful multiparticle entanglement and sub-shot-noise sensitivity in experimental phase estimation," *Phys. Rev. Lett.* **107**, 080504.
- Kruse, I, K. Lange, J. Peise, B. Lücke, L. Pezzè, J. Arlt, W. Ertmer, C. Lisdat, L. Santos, A. Smerzi, and C. Klempt (2016), "0.75 atoms improve the clock signal of 10,000 atoms," [arXiv:1605.07754 \[quant-ph\]](https://arxiv.org/abs/1605.07754).
- Kurkjian, H, K. Pawłowski, A. Sinatra, and P. Treutlein (2013), "Spin squeezing and Einstein-Podolsky-Rosen entanglement of two bimodal condensates in state-dependent potentials," *Phys. Rev. A* **88**, 043605.
- Kuzmich, A, N. P. Bigelow, and L. Mandel (1998), "Atomic quantum non-demolition measurements and squeezing," *Europhys. Lett.* **42** (5), 481.
- Kuzmich, A, L. Mandel, and N. P. Bigelow (2000), "Generation of spin squeezing via continuous quantum nondemolition measurement," *Phys. Rev. Lett.* **85**, 1594–1597.
- Kuzmich, A, L. Mandel, J. Janis, Y. E. Young, R. Egnis-man, and N. P. Bigelow (1999), "Quantum nondemolition measurements of collective atomic spin," *Phys. Rev. A* **60**, 2346–2350.
- Kuzmich, A, K. Mølmer, and E. S. Polzik (1997), "Spin squeezing in an ensemble of atoms illuminated with squeezed light," *Phys. Rev. Lett.* **79**, 4782–4785.
- Kuzmich, A, and E. S. Polzik (2003), "Atomic continuous variable processing and light-atoms quantum interface," in

- Quantum Information with Continuous Variables*, edited by S. L. Braunstein and A. K. Pati, Chap. 18 (Kluwer Academic Publishers) p. 231.
- Kwiat, P. G., K. Mattle, H. Weinfurter, A. Zeilinger, A. V. Sergienko, and Y. Shih (1995), “New high-intensity source of polarization-entangled photon pairs,” *Phys. Rev. Lett.* **75**, 4337–4341.
- Laloë, F., and W. J. Mullin (2009), “Interferometry with independent Bose-Einstein condensates: parity as an EPR/Bell quantum variable,” *Eur. Phys. J. B* **70** (3), 377–396.
- Landau, L., and E. Lifchitz (1975), *Physique théorique*, 3rd ed., Vol. 3: Mécanique quantique (Éditions Mir).
- Landini, M., M. Fattori, L. Pezzè, and A. Smerzi (2014), “Phase-noise protection in quantum-enhanced differential interferometry,” *New J. Phys.* **16** (11), 113074.
- Lanyon, B. P., M. Zwerger, P. Jurcevic, C. Hempel, W. Dür, H. J. Briegel, R. Blatt, and C. F. Roos (2014), “Experimental violation of multipartite Bell inequalities with trapped ions,” *Phys. Rev. Lett.* **112**, 100403.
- Lapert, M., G. Ferrini, and D. Sugny (2012), “Optimal control of quantum superpositions in a bosonic Josephson junction,” *Phys. Rev. A* **85**, 023611.
- Law, C. K., H. T. Ng, and P. T. Leung (2001), “Coherent control of spin squeezing,” *Phys. Rev. A* **63**, 055601.
- Law, C. K., H. Pu, and N. P. Bigelow (1998), “Quantum spins mixing in spinor Bose-Einstein condensates,” *Phys. Rev. Lett.* **81**, 5257–5261.
- LeBlanc, L. J., A. B. Bardou, J. McKeever, M. H. T. Extavour, D. Jervis, J. H. Thywissen, F. Piazza, and A. Smerzi (2011), “Dynamics of a tunable superfluid junction,” *Phys. Rev. Lett.* **106**, 025302.
- Lee, H., P. Kok, and J. P. Dowling (2002), “A quantum Rosetta stone for interferometry,” *J. Mod. Opt.* **49**, 2325.
- Lee, P. J., K.-A. Brickman, L. Deslauriers, P. C. Haljan, L.-M. Duan, and C. Monroe (2005), “Phase control of trapped ion quantum gates,” *J. Opt. B: Quantum and Semiclassical Optics* **7** (10), S371.
- Lee, T.-W., S. D. Huver, H. Lee, L. Kaplan, S. B. McCracken, C. Min, D. B. Uskov, C. F. Wildfeuer, G. Veronis, and J. P. Dowling (2009), “Optimization of quantum interferometric metrological sensors in the presence of photon loss,” *Phys. Rev. A* **80**, 063803.
- Leggett, A. J. (2001), “Bose-Einstein condensation in the alkali gases: Some fundamental concepts,” *Rev. Mod. Phys.* **73**, 307–356.
- Leggett, A. J., and F. Sols (1991), “On the concept of spontaneously broken gauge symmetry in condensed matter physics,” *Found. Phys.* **21**, 353.
- Leggett, A. J., and F. Sols (1998), “Comment on ‘phase and phase diffusion of a split Bose-Einstein condensate,’” *Phys. Rev. Lett.* **81**, 1344–1344.
- Lehmann, E. L., and G. Casella (2003), *Theory of Point Estimation* (Springer Texts in Statistics).
- Leibfried, D., M. D. Barrett, T. Schaetz, J. Britton, J. Chiaverini, W. M. Itano, J. D. Jost, C. Langer, and D. J. Wineland (2004), “Toward Heisenberg-limited spectroscopy with multiparticle entangled states,” *Science* **304**, 1476.
- Leibfried, D., R. Blatt, C. Monroe, and D. Wineland (2003a), “Quantum dynamics of single trapped ions,” *Rev. Mod. Phys.* **75**, 281–324.
- Leibfried, D., B. DeMarco, V. Meyer, D. Lucas, M. Barrett, J. Britton, W. M. Itano, B. Jelenković, C. Langer, T. Rosenband, and D. J. Wineland (2003b), “Experimental demonstration of a robust, high-fidelity geometric two ion-qubit phase gate,” *Nature* **422**, 412.
- Leibfried, D., E. Knill, S. Seidelin, J. Britton, R. B. Blakestad, J. Chiaverini, D. B. Hume, W. M. Itano, J. D. Jost, C. Langer, R. Ozeri, R. Reichle, and D. J. Wineland (2005), “Creation of a six-atom ‘Schrödinger cat’ state,” *Nature* **438**, 639.
- Leibfried, D., D. M. Meekhof, B. E. King, C. Monroe, W. M. Itano, and D. J. Wineland (1996), “Experimental determination of the motional quantum state of a trapped atom,” *Phys. Rev. Lett.* **77**, 4281–4285.
- Leroux, I. D., M. H. Schleier-Smith, and V. Vuletić (2010a), “Implementation of cavity squeezing of a collective atomic spin,” *Phys. Rev. Lett.* **104**, 073602.
- Leroux, I. D., M. H. Schleier-Smith, and V. Vuletić (2010b), “Orientation-dependent entanglement lifetime in a squeezed atomic clock,” *Phys. Rev. Lett.* **104**, 250801.
- Leroux, I. D., M. H. Schleier-Smith, H. Zhang, and V. Vuletić (2012), “Unitary cavity spin squeezing by quantum erasure,” *Phys. Rev. A* **85**, 013803.
- Leslie, S. R., J. Guzman, M. Vengalattore, Jay D. Sau, Marvin L. Cohen, and D. M. Stamper-Kurn (2009), “Amplification of fluctuations in a spinor Bose-Einstein condensate,” *Phys. Rev. A* **79**, 043631.
- Levy, S., E. Lahoud, I. Shomroni, and J. Steinhauer (2007), “The a.c. and d.c. Josephson effects in a Bose-Einstein condensate,” *Nature* **449**, 579.
- Lewenstein, M., and L. You (1996), “Quantum phase diffusion of a Bose-Einstein condensate,” *Phys. Rev. Lett.* **77**, 3489–3493.
- Lewis-Swan, R. J., and K. V. Kheruntsyan (2014), “Proposal for demonstrating the Hong–Ou–Mandel effect with matter waves,” *Nat. Comm.* **5**, 3752 EP –.
- Li, Y., Y. Castin, and A. Sinatra (2008a), “Optimum spin squeezing in Bose-Einstein condensates with particle losses,” *Phys. Rev. Lett.* **100**, 210401.
- Li, Y., P. Treutlein, J. Reichel, and A. Sinatra (2008b), “Spin squeezing in a bimodal condensate: spatial dynamics and particle losses,” *Eur. Phys. J. B* **68**, 365.
- Lidar, D. A., I. L. Chuang, and K. B. Whaley (1998), “Decoherence-free subspaces for quantum computation,” *Phys. Rev. Lett.* **81**, 2594–2597.
- Linnemann, D., H. Strobel, W. Muessel, J. Schulz, R. J. Lewis-Swan, K. V. Kheruntsyan, and M. K. Oberthaler (2016), “Quantum-enhanced sensing based on time reversal of nonlinear dynamics,” *Phys. Rev. Lett.* **117**, 013001.
- Lipkin, H. J., N. Meshkov, and A. J. Glick (1965), “Validity of many-body approximation methods for a solvable model: (i). exact solutions and perturbation theory,” *Nucl. Phys.* **62**, 188.
- Liu, C., Z. Dutton, C. H. Behroozi, and L. V. Hau (2001), “Observation of coherent optical information storage in an atomic medium using halted light pulses,” *Nature* **409** (6819), 490–493.
- Liu, Y., E. Gomez, S. E. Maxwell, L. D. Turner, E. Tiesinga, and P. D. Lett (2009), “Number fluctuations and energy dissipation in sodium spinor condensates,” *Phys. Rev. Lett.* **102**, 225301.
- Lopes, R., A. Imanaliev, A. Aspect, M. Cheneau, D. Boiron, and C. I. Westbrook (2015), “Atomic Hong-Ou-Mandel experiment,” *Nature* **520**, 66.
- Louchet-Chauvet, A., J. Appel, J. J. Renema, D. Oblak, N. Kjørgaard, and E. S. Polzik (2010), “Entanglement-

- assisted atomic clock beyond the projection noise limit,” *New J. Phys.* **12** (6), 065032.
- Lu, X-M, S. Yu, and C. H. Oh (2015), “Robust quantum metrological schemes based on protection of quantum Fisher information,” *Nat. Comm.* **6** (6), 065032.
- Lücke, B, J. Peise, G. Vitagliano, J. Arlt, L. Santos, G. Tóth, and C. Klempt (2014), “Detecting multiparticle entanglement of Dicke states,” *Phys. Rev. Lett.* **112**, 155304.
- Lücke, B, M. Scherer, J. Kruse, L. Pezzè, F. Deuretzbacher, P. Hyllus, O. Topic, J. Peise, W. Ertmer, J. Arlt, L. Santos, A. Smerzi, and C. Klempt (2011), “Twin matter waves for interferometry beyond the classical limit,” *Science* **334**, 773.
- Ludlow, A D, M. M. Boyd, J. Ye, E. Peik, and P. O. Schmidt (2015), “Optical atomic clocks,” *Rev. Mod. Phys.* **87**, 637–701.
- Ludlow, A D, T. Zelevinsky, G. K. Campbell, S. Blatt, M. M. Boyd, M. H. G. de Miranda, M. J. Martin, J. W. Thomsen, S. M. Foreman, J. Ye, T. M. Fortier, J. E. Stalnaker, S. A. Diddams, Y. Le Coq, Z. W. Barber, N. Poli, N. D. Lemke, K. M. Beck, and C. W. Oates (2008), “Sr lattice clock at 1×10^{-16} fractional uncertainty by remote optical evaluation with a Ca clock,” *Science* **319**, 1805.
- Luis, A (2004), “Nonlinear transformations and the Heisenberg limit,” *Phys. Lett. A* **329**, 8.
- Luis, A (2007), “Quantum limits, nonseparable transformations, and nonlinear optics,” *Phys. Rev. A* **76**, 035801.
- Lvovsky, A I, H. Hansen, T. Aichele, O. Benson, J. Mlynek, and S. Schiller (2001), “Quantum state reconstruction of the single-photon Fock state,” *Phys. Rev. Lett.* **87**, 050402.
- Lynch, R (1995), “The quantum phase problem: a critical review,” *Phys. Rep.* **256**, 367.
- Ma, J, and X. Wang (2009), “Fisher information and spin squeezing in the Lipkin-Meshkov-Glick model,” *Phys. Rev. A* **80**, 012318.
- Ma, J, X. Wang, C. P. Sun, and F. Nori (2011), “Quantum spin squeezing,” *Phys. Rep.* **509**, 89.
- Mach, L (1892), “Ueber einen Interferenzrefraktor,” *Z. Instrumentenkunde* **12** (3), 89–93.
- Macieszczak, K (2015), “Zeno limit in frequency estimation with non-Markovian environments,” *Phys. Rev. A* **92**, 010102.
- Macrì, T, A. Smerzi, and L. Pezzè (2016), “Loschmidt echo for quantum metrology,” *Phys. Rev. A* **94**, 010102.
- Madsen, L B, and K. Mølmer (2004), “Spin squeezing and precision probing with light and samples of atoms in the Gaussian description,” *Phys. Rev. A* **70**, 052324.
- Marino, A M, N. V. Corzo Trejo, and P. D. Lett (2012), “Effect of losses on the performance of an SU(1,1) interferometer,” *Phys. Rev. A* **86**, 023844.
- Martinez, E A, C. A. Muschik, P. Schindler, D. Nigg, A. Erhard, M. Heyl, P. Hauke, M. Dalmonte, T. Monz, P. Zoller, and R. Blatt (2016), “Real-time dynamics of lattice gauge theories with a few-qubit quantum computer,” *Nature* **534** (7608), 516–519.
- Matsuzaki, Y, S. C. Benjamin, and J. Fitzsimons (2011), “Magnetic field sensing beyond the standard quantum limit under the effect of decoherence,” *Phys. Rev. A* **84**, 012103.
- Maussang, K, G. E. Marti, T. Schneider, P. Treutlein, Y. Li, A. Sinatra, R. Long, J. Estève, and J. Reichel (2010), “Enhanced and reduced atom number fluctuations in a BEC splitter,” *Phys. Rev. Lett.* **105**, 080403.
- McConnell, R, H. Zhang, S. Čuk, J. Hu, M. H. Schleier-Smith, and V. Vuletić (2013), “Generating entangled spin states for quantum metrology by single-photon detection,” *Phys. Rev. A* **88**, 063802.
- McConnell, Robert, Hao Zhang, Jiazhong Hu, Senka Cuk, and Vladan Vuletić (2015), “Entanglement with negative Wigner function of almost 3,000 atoms heralded by one photon,” *Nature* **519** (7544), 439–442.
- Meiser, D, J. Ye, and M. J. Holland (2008), “Spin squeezing in optical lattice clocks via lattice-based QND measurements,” *New J. Phys.* **10**, 073014.
- Menotti, C, J. R. Anglin, J. I. Cirac, and P. Zoller (2001), “Dynamic splitting of a Bose-Einstein condensate,” *Phys. Rev. A* **63**, 023601.
- Meshkov, N, A. J. Glick, and H. J. Lipkin (1965), “Validity of many-body approximation methods for a solvable model: (ii). linearization procedures,” *Nucl. Phys.* **62**, 199.
- Meyer, V, M. A. Rowe, D. Kielpinski, C. A. Sackett, W. M. Itano, C. Monroe, and D. J. Wineland (2001), “Experimental demonstration of entanglement-enhanced rotation angle estimation using trapped ions,” *Phys. Rev. Lett.* **86**, 5870–5873.
- Mias, G I, N. R. Cooper, and S. M. Girvin (2008), “Quantum noise, scaling, and domain formation in a spinor Bose-Einstein condensate,” *Phys. Rev. A* **77**, 023616.
- Micheli, A, D. Jaksch, J. I. Cirac, and P. Zoller (2003), “Many-particle entanglement in two-component Bose-Einstein condensates,” *Phys. Rev. A* **67**, 013607.
- Mielenz, M, H. Kalis, M. Wittemer, F. Hakelberg, U. Warring, R. Schmied, M. Blain, P. Maunz, D. L. Moehring, D. Leibfried, and T. Schaetz (2016), “Arrays of individually controlled ions suitable for two-dimensional quantum simulations,” *Nat. Comm.* **7**, ncomms11839.
- Milburn, G J, J. Corney, E. M. Wright, and D. F. Walls (1997), “Quantum dynamics of an atomic Bose-Einstein condensate in a double-well potential,” *Phys. Rev. A* **55**, 4318–4324.
- Milburn, G J, S. Schneider, and D. F. V. James (2000), “Ion trap quantum computing with warm ions,” *Fortschr. Phys.* **48**, 801–810.
- Miller, R, T. E. Northup, K. M. Birnbaum, A. Boca, A. D. Boozer, and H. J. Kimble (2005), “Trapped atoms in cavity QED: coupling quantized light and matter,” *J. Phys. B: At. Mol. Opt. Phys.* **38** (9), S551.
- Mintert, F, and C. Wunderlich (2001), “Ion-trap quantum logic using long-wavelength radiation,” *Phys. Rev. Lett.* **87**, 257904.
- Mitchell, M W (2005), “Metrology with entangled states,” *Proc. SPIE* **5893**, 589310.
- Moehring, D L, P. Maunz, S. Olmschenk, K. C. Younge, D. N. Matsukevich, L. M. Duan, and C. Monroe (2007), “Entanglement of single-atom quantum bits at a distance,” *Nature* **449** (7158), 68–71.
- Mølmer, K, and A. Sørensen (1999), “Multiparticle entanglement of hot trapped ions,” *Phys. Rev. Lett.* **82**, 1835–1838.
- Monz, T, P. Schindler, J. T. Barreiro, M. Chwalla, D. Nigg, W. A. Coish, M. Harlander, W. Hänsel, M. Hennrich, and R. Blatt (2011), “14-qubit entanglement: Creation and coherence,” *Phys. Rev. Lett.* **106**, 130506.
- Morsch, O, and M. K. Oberthaler (2006), “Dynamics of Bose-Einstein condensates in optical lattices,” *Rev. Mod. Phys.* **78**, 179–215.
- Muessel, W, H. Strobel, D. Linnemann, D. B. Hume, and M. K. Oberthaler (2014), “Scalable spin squeezing for quantum-enhanced magnetometry with Bose-Einstein condensates,” *Phys. Rev. Lett.* **113**, 103004.

- Muessel, W, H. Strobel, D. Linnemann, T. Zibold, B. Juliá-Díaz, and M. K. Oberthaler (2015), “Twist-and-turn spin squeezing in Bose-Einstein condensates,” *Phys. Rev. A* **92**, 023603.
- Mullan, M, and E. Knill (2014), “Optimizing passive quantum clocks,” *Phys. Rev. A* **90**, 042310.
- Mullin, W J, and F. Laloë (2008), “Interference of Bose-Einstein condensates: Quantum nonlocal effects,” *Phys. Rev. A* **78**, 061605.
- Müstecaplıoğlu, Ö E, M. Zhang, and L. You (2002), “Spin squeezing and entanglement in spinor condensates,” *Phys. Rev. A* **66**, 033611.
- Myerson, A H, D. J. Szwer, S. C. Webster, D. T. C. Allcock, M. J. Curtis, G. Imreh, J. A. Sherman, D. N. Stacey, A. M. Steane, and D. M. Lucas (2008), “High-fidelity readout of trapped-ion qubits,” *Phys. Rev. Lett.* **100**, 200502.
- Nagata, T, R. Okamoto, J. L. O’Brien, K. Sasaki, and S. Takeuchi (2007), “Beating the standard quantum limit with four-entangled photons,” *Science* **316** (5825), 726–729.
- Napolitano, M, M. Koschorreck, B. Dubost, N. Behbood, R. J. Sewell, and M. W. Mitchell (2011), “Interaction-based quantum metrology showing scaling beyond the Heisenberg limit,” *Nature* **471**, 486.
- Napolitano, M, and M. W. Mitchell (2010), “Nonlinear metrology with a quantum interface,” *New J. Phys.* **12** (9), 093016.
- Negretti, A, and C. Henkel (2004), “Enhanced phase sensitivity and soliton formation in an integrated BEC interferometer,” *J. Phys. B: At. Mol. Opt. Phys.* **37**, L385.
- Negretti, A, C. Henkel, and K. Mølmer (2008), “Quantum-limited position measurements of a dark matter-wave soliton,” *Phys. Rev. A* **77**, 043606.
- Nelson, Karl D, Xiao Li, and David S. Weiss (2007), “Imaging single atoms in a three-dimensional array,” *Nat Phys* **3** (8), 556–560.
- Nicholson, T L, S. L. Campbell, R. B. Hutson, G. E. Marti, B. J. Bloom, R. L. McNally, W. Zhang, M. D. Barrett, M. S. Safronova, G. F. Strouse, W. L. Tew, and J. Ye (2015), “Systematic evaluation of an atomic clock at 2×10^{-18} total uncertainty,” *Nat. Comm.* **6**, 6896 EP –.
- Nicholson, T L, M. J. Martin, J. R. Williams, B. J. Bloom, M. Bishof, M. D. Swallows, S. L. Campbell, and J. Ye (2012), “Comparison of two independent Sr optical clocks with 1×10^{-17} stability at 10^3 s,” *Phys. Rev. Lett.* **109**, 230801.
- Nicklas, E, M. Karl, M. Höfer, A. Johnson, W. Muesel, H. Strobel, J. Tomkovič, T. Gasenzer, and M. K. Oberthaler (2015), “Observation of scaling in the dynamics of a strongly quenched quantum gas,” *Phys. Rev. Lett.* **115**, 245301.
- Nielsen, A E B, and K. Mølmer (2008), “Atomic spin squeezing in an optical cavity,” *Phys. Rev. A* **77**, 063811.
- Nielsen, M A, and I. L. Chuang (2000), *Quantum Computation and Quantum Information* (Cambridge Univ. Press, Cambridge, England).
- Noguchi, A, K. Toyoda, and S. Urabe (2012), “Generation of Dicke states with phonon-mediated multilevel stimulated Raman adiabatic passage,” *Phys. Rev. Lett.* **109**, 260502.
- Oblak, D, P. G. Petrov, C. L. Garrido Alzar, W. Tittel, A. K. Vershovski, J. K. Mikkelsen, J. L. Sørensen, and E. S. Polzik (2005), “Quantum-noise-limited interferometric measurement of atomic noise: Towards spin squeezing on the Cs clock transition,” *Phys. Rev. A* **71**, 043807.
- O’Brien, J L, A. Furusawa, and J. Vučković (2009), “Photonic quantum technologies,” *Nat. Photon.* **3** (12), 687–695.
- Ockeloen, C F, R. Schmied, M. F. Riedel, and P. Treutlein (2013), “Quantum metrology with a scanning probe atom interferometer,” *Phys. Rev. Lett.* **111**, 143001.
- Ockeloen, C F, A. F. Tauschinsky, R. J. C. Spreeuw, and S. Whitlock (2010), “Detection of small atom numbers through image processing,” *Phys. Rev. A* **82**, 061606.
- Ohmi, T, and K. Machida (1998), “Bose-Einstein condensation with internal degrees of freedom in alkali atom gases,” *J. Phys. Soc. Jpn.* **67** (6), 1822.
- Ono, T, and H. F. Hofmann (2010), “Effects of photon losses on phase estimation near the Heisenberg limit using coherent light and squeezed vacuum,” *Phys. Rev. A* **81**, 033819.
- Orús, R, S. Dusuel, and J. Vidal (2008), “Equivalence of critical scaling laws for many-body entanglement in the Lipkin-Meshkov-Glick model,” *Phys. Rev. Lett.* **101**, 025701.
- Ospelkaus, C, C. E. Langer, J. M. Amini, K. R. Brown, D. Leibfried, and D. J. Wineland (2008), “Trapped-ion quantum logic gates based on oscillating magnetic fields,” *Phys. Rev. Lett.* **101**, 090502.
- Ospelkaus, C, U. Warring, Y. Colombe, K. R. Brown, J. M. Amini, D. Leibfried, and D. J. Wineland (2011), “Microwave quantum logic gates for trapped ions,” *Nature* **476** (7359), 181–184.
- Ou, Z Y, S. F. Pereira, H. J. Kimble, and K. C. Peng (1992), “Realization of the Einstein-Podolsky-Rosen paradox for continuous variables,” *Phys. Rev. Lett.* **68**, 3663–3666.
- Ozeri, R (2013), “Heisenberg limited metrology using quantum error-correction codes,” [arXiv:1310.3432 \[quant-ph\]](https://arxiv.org/abs/1310.3432).
- Paraoanu, G S, S. Kohler, F. Sols, and A. J. Leggett (2001), “The Josephson plasmon as a Bogoliubov quasiparticle,” *J. Phys. B: At. Mol. Opt. Phys.* **34**, 4689.
- Paris, M, and J. Řeháček, Eds. (2004), *Quantum State Estimation*, Lect. Notes Phys., Vol. 649 (Springer, Berlin Heidelberg).
- Paris, M G A (2009), “Quantum estimation for quantum technology,” *Int. J. Quant. Inf.* **07**, 125–137.
- Peise, J, I. Kruse, K. Lange, B. Lücke, L. Pezzè, J. Arlt, W. Ertmer, K. Hammerer, L. Santos, A. Smerzi, and C. Klempt (2015a), “Satisfying the Einstein-Podolsky-Rosen criterion with massive particles,” *Nat. Comm.* **6**, 1038.
- Peise, J, B. Lücke, L. Pezzè, F. Deuretzbacher, W. Ertmer, J. Arlt, A. Smerzi, L. Santos, and C. Klempt (2015b), “Interaction-free measurements by quantum Zeno stabilization of ultracold atoms,” *Nat. Comm.* **6**, 6811.
- Pelisson, Sophie, Luca Pezzè, and Augusto Smerzi (2016), “Nonlocality with ultracold atoms in a lattice,” *Phys. Rev. A* **93**, 022115.
- Peres, A (1995), *Quantum Theory: Concepts and Methods* (Springer Netherlands).
- Petersen, V, L. B. Madsen, and K. Mølmer (2005), “Magnetometry with entangled atomic samples,” *Phys. Rev. A* **71**, 012312.
- Petz, D (2002), “Covariance and Fisher information in quantum mechanics,” *J. Phys. A: Math. Gen.* **35** (4), 929–939.
- Pezzè, L, L. A. Collins, A. Smerzi, G. P. Berman, and A. R. Bishop (2005), “Sub-shot-noise phase sensitivity with a Bose-Einstein condensate Mach-Zehnder interferometer,” *Phys. Rev. A* **72**, 043612.
- Pezzè, L, Y. Li, W. Li, and A. Smerzi (2015), “Witnessing entanglement without entanglement witness operators,” [arXiv:1512.06213 \[quant-ph\]](https://arxiv.org/abs/1512.06213).

- Pezzè, L., and A. Smerzi (2006), “Phase sensitivity of a Mach-Zehnder interferometer,” *Phys. Rev. A* **73**, 011801.
- Pezzè, L., and A. Smerzi (2007), “Sub shot-noise interferometric phase sensitivity with beryllium ions Schrödinger cat states,” *Europhys. Lett.* **78**, 30004.
- Pezzè, L., and A. Smerzi (2008), “Mach-Zehnder interferometry at the Heisenberg limit with coherent and squeezed-vacuum light,” *Phys. Rev. Lett.* **100**, 073601.
- Pezzè, L., and A. Smerzi (2009), “Entanglement, nonlinear dynamics, and the Heisenberg limit,” *Phys. Rev. Lett.* **102**, 100401.
- Pezzè, L., and A. Smerzi (2013), “Ultrasensitive two-mode interferometry with single-mode number squeezing,” *Phys. Rev. Lett.* **110**, 163604.
- Pezzè, L., and A. Smerzi (2014), “Quantum theory of phase estimation,” in *Atom Interferometry, Proceedings of the International School of Physics “Enrico Fermi”, Course 188, Varenna*, edited by G. M. Tino and M. A. Kasevich (IOS Press, Amsterdam) p. 691.
- Phillips, D F, A. Fleischhauer, A. Mair, R. L. Walsworth, and M. D. Lukin (2001), “Storage of light in atomic vapor,” *Phys. Rev. Lett.* **86**, 783–786.
- Pichler, T, T. Caneva, S. Montangero, M. D. Lukin, and T. Calarco (2016), “Noise-resistant optimal spin squeezing via quantum control,” *Phys. Rev. A* **93**, 013851.
- Plenio, M B, and S. F. Huelga (2016), “Sensing in the presence of an observed environment,” *Phys. Rev. A* **93**, 032123.
- Porras, D, and J. I. Cirac (2006), “Quantum manipulation of trapped ions in two dimensional coulomb crystals,” *Phys. Rev. Lett.* **96**, 250501.
- Pu, H, C. K. Law, S. Raghavan, J. H. Eberly, and N. P. Bigelow (1999), “Spin-mixing dynamics of a spinor Bose-Einstein condensate,” *Phys. Rev. A* **60**, 1463–1470.
- Pu, H, and P. Meystre (2000), “Creating macroscopic atomic Einstein-Podolsky-Rosen states from Bose-Einstein condensates,” *Phys. Rev. Lett.* **85**, 3987–3990.
- Pupillo, G, A. Micheli, M. Boninsegni, I. Lesanovsky, and P. Zoller (2010), “Strongly correlated gases of rydberg-dressed atoms: Quantum and classical dynamics,” *Phys. Rev. Lett.* **104**, 223002.
- Radcliffe, J M (1971), “Some properties of coherent spin states,” *J. Phys. A: Gen. Phys.* **4** (3), 313.
- Raghavan, S, H. Pu, P. Meystre, and N. P. Bigelow (2001), “Generation of arbitrary Dicke states in spinor Bose-Einstein condensates,” *Opt. Comm.* **188**, 149.
- Raghavan, S, A. Smerzi, S. Fantoni, and S. R. Shenoy (1999), “Coherent oscillations between two weakly coupled Bose-Einstein condensates: Josephson effects, π oscillations, and macroscopic quantum self-trapping,” *Phys. Rev. A* **59**, 620–633.
- Ramsey, N F (1963), *Molecular Beams* (International Series of Monographs on Physics, Oxford University Press, London).
- Rao, C R (1945), “Information and the accuracy attainable in the estimation of statistical parameters,” *Bull. Calcutta Math. Soc.* **37**, 81–91.
- Rashid, M A (1978), “The intelligent states. i. group-theoretic study and the computation of matrix elements,” *J. Math. Phys.* **19**, 1391.
- Raymer, M G, A. C. Funk, B. C. Sanders, and H. de Guise (2003), “Separability criterion for separate quantum systems,” *Phys. Rev. A* **67**, 052104.
- Reid, M D (1989), “Demonstration of the Einstein-Podolsky-Rosen paradox using nondegenerate parametric amplification,” *Phys. Rev. A* **40**, 913–923.
- Reid, M D, P. D. Drummond, W. P. Bowen, E. G. Cavalcanti, P. K. Lam, H. A. Bachor, U. L. Andersen, and G. Leuchs (2009), “*Colloquium*: The Einstein-Podolsky-Rosen paradox: From concepts to applications,” *Rev. Mod. Phys.* **81**, 1727–1751.
- Rey, A M, L. Jiang, and M. D. Lukin (2007), “Quantum-limited measurements of atomic scattering properties,” *Phys. Rev. A* **76**, 053617.
- Richerme, P, Z.-X. Gong, A. Lee, C. Senko, J. Smith, M. Foss-Feig, S. Michalakis, A. V. Gorshkov, and C. Monroe (2014), “Non-local propagation of correlations in quantum systems with long-range interactions,” *Nature* **511**, 198–201.
- Riedel, M F, P. Böhi, Y. Li, T. W. Hänsch, A. Sinatra, and P. Treutlein (2010), “Atom-chip-based generation of entanglement for quantum metrology,” *Nature* **464**, 1170–1173.
- Ritsch, H, P. Domokos, F. Brennecke, and T. Esslinger (2013), “Cold atoms in cavity-generated dynamical optical potentials,” *Rev. Mod. Phys.* **85**, 553–601.
- Rivas, Á, and S. F. Huelga (2012), *Open Quantum Systems* (Springer, New York).
- Rivas, Á, and A. Luis (2010), “Precision quantum metrology and nonclassicality in linear and nonlinear detection schemes,” *Phys. Rev. Lett.* **105**, 010403.
- Roos, C (2014), “Quantum information processing with trapped ions,” in *Fundamental Physics in Particle Traps, Springer Tracts in Modern Physics, Vol. 256*, edited by Wolfgang Quint and Manuel Vogel (Springer-Verlag) pp. 253–292.
- Roos, C F, M. Chwalla, K. Kim, M. Riebe, and R. Blatt (2006), “‘Designer atoms’ for quantum metrology,” *Nature* **443**, 316.
- Roos, C F, M. Riebe, H. Häffner, W. Hänsel, J. Benhelm, G. P. T. Lancaster, C. Becher, F. Schmidt-Kaler, and R. Blatt (2004), “Control and measurement of three-qubit entangled states,” *Science* **304** (5676), 1478–1480.
- Rosenband, T, and D. R. Leibbrandt (2013), “Exponential scaling of clock stability with atom number,” [arXiv:1303.6357 \[quant-ph\]](https://arxiv.org/abs/1303.6357).
- Rosenband, Till, Ed. (2011), *Numerical test of few-qubit clock protocols* (Proceedings of the 20th International Conference on Laser Spectroscopy).
- Roy, S M, and S. L. Braunstein (2008), “Exponentially enhanced quantum metrology,” *Phys. Rev. Lett.* **100**, 220501.
- Sackett, C A, D. Kielpinski, B. E. King, C. Langer, V. Meyer, C. J. Myatt, M. Rowe, Q. A. Turchette, W. M. Itano, D. J. Wineland, and C. Monroe (2000), “Experimental entanglement of four particles,” *Nature* **404**, 256.
- Sadler, L E, J. M. Higbie, S. R. Leslie, M. Vengalattore, and D. M. Stamper-Kurn (2006), “Spontaneous symmetry breaking in a quenched ferromagnetic spinor Bose-Einstein condensate,” *Nature* **443** (7109), 312–315.
- Saffman, M, D. Oblak, J. Appel, and E. S. Polzik (2009), “Spin squeezing of atomic ensembles by multicolor quantum nondemolition measurements,” *Phys. Rev. A* **79**, 023831.
- Saffman, M, T. G. Walker, and K. Mølmer (2010), “Quantum information with rydberg atoms,” *Rev. Mod. Phys.* **82**, 2313–2363.
- Sanders, B C, and G. J. Milburn (1995), “Optimal quantum measurements for phase estimation,” *Phys. Rev. Lett.* **75**, 2944–2947.

- Santarelli, G, Ph. Laurent, P. Lemonde, A. Clairon, A. G. Mann, S. Chang, A. N. Luiten, and C. Salomon (1999), “Quantum projection noise in an atomic fountain: A high stability cesium frequency standard,” *Phys. Rev. Lett.* **82**, 4619–4622.
- Sau, J D, S. R. Leslie, M. L. Cohen, and D. M. Stamper-Kurn (2010), “Spin squeezing of high-spin, spatially extended quantum fields,” *New J. Phys.* **12**, 085011.
- Schaff, J-F, T. Langen, and J. Schmiedmayer (2014), “Interferometry with atoms,” in *Atom Interferometry, Proceedings of the International School of Physics “Enrico Fermi”, Course 188, Varenna*, edited by G. M. Tino and M. A. Kasevich (IOS Press, Amsterdam) p. 1.
- Scherer, M, B. Lücke, G. Gebreyesus, O. Topic, F. Deuretzbacher, W. Ertmer, L. Santos, J. J. Arlt, and C. Klempt (2010), “Spontaneous breaking of spatial and spin symmetry in spinor condensates,” *Phys. Rev. Lett.* **105**, 135302.
- Schleich, W P (2001), *Quantum Optics in Phase Space* (Wiley-VCH, Berlin).
- Schleier-Smith, M H, I. D. Leroux, and V. Vuletić (2010a), “Squeezing the collective spin of a dilute atomic ensemble by cavity feedback,” *Phys. Rev. A* **81**, 021804.
- Schleier-Smith, M H, I. D. Leroux, and V. Vuletić (2010b), “States of an ensemble of two-level atoms with reduced quantum uncertainty,” *Phys. Rev. Lett.* **104**, 073604.
- Schmaljohann, H, M. Erhard, J. Kronjäger, M. Kottke, S. van Staa, L. Cacciapuoti, J. J. Arlt, K. Bongs, and K. Sengstock (2004), “Dynamics of $F = 2$ spinor Bose-Einstein condensates,” *Phys. Rev. Lett.* **92**, 040402.
- Schmied, R, J.-D. Bancal, B. Allard, M. Fadel, V. Scarani, P. Treutlein, and N. Sangouard (2016), “Bell correlations in a Bose-Einstein condensate,” *Science* **352** (6284), 441–444.
- Schmied, R, and P. Treutlein (2011), “Tomographic reconstruction of the Wigner function on the Bloch sphere,” *New J. Phys.* **13** (6), 065019.
- Schmied, R, J. H. Wesenberg, and D. Leibfried (2011), “Quantum simulation of the hexagonal Kitaev model with trapped ions,” *New J. Phys.* **13** (11), 115011.
- Schmied, Roman (2016), “Quantum state tomography of a single qubit: comparison of methods,” *J. Mod. Opt.* **63** (18), 1744–1758.
- Schnabel, R, N. Mavalvala, D. E. McClelland, and P. K. Lam (2010), “Quantum metrology for gravitational wave astronomy,” *Nat. Comm.* **1**, 121.
- Schneider, C, D. Porras, and T. Schaetz (2012), “Experimental quantum simulations of many-body physics with trapped ions,” *Rep. Prog. Phys.* **75** (2), 024401.
- Schumm, T, S. Hofferberth, L. M. Andersson, S. Wildermuth, S. Groth, I. Bar-Joseph, J. Schmiedmayer, and P. Krüger (2005), “Matter-wave interferometry in a double well on an atom chip,” *Nat. Phys.* **1**, 57–62.
- Scott, R G, T. E. Judd, and T. M. Fromhold (2008), “Exploiting soliton decay and phase fluctuations in atom chip interferometry of Bose-Einstein condensates,” *Phys. Rev. Lett.* **100**, 100402.
- Scully, M O, and M. S. Zubairy (1997), *Quantum Optics* (Cambridge Univ. Press).
- Sewell, R J, M. Koschorreck, M. Napolitano, B. Dubost, N. Behbood, and M. W. Mitchell (2012), “Magnetic sensitivity beyond the projection noise limit by spin squeezing,” *Phys. Rev. Lett.* **109**, 253605.
- Sewell, R J, M. Napolitano, N. Behbood, G. Colangelo, F. Martin Ciurana, and M. W. Mitchell (2014), “Ultra-sensitive atomic spin measurements with a nonlinear interferometer,” *Phys. Rev. X* **4**, 021045.
- Sewell, R J, M. Napolitano, N. Behbood, G. Colangelo, and M. W. Mitchell (2013), “Certified quantum non-demolition measurement of a macroscopic material system,” *Nat. Photon.* **7**, 517.
- Shah, V, G. Vasilakis, and M. V. Romalis (2010), “High bandwidth atomic magnetometry with continuous quantum nondemolition measurements,” *Phys. Rev. Lett.* **104**, 013601.
- Shaji, A, and C. M. Caves (2007), “Qubit metrology and decoherence,” *Phys. Rev. A* **76**, 032111.
- Shchesnovich, V S, and M. Trippenbach (2008), “Fock-space WKB method for the boson Josephson model describing a Bose-Einstein condensate trapped in a double-well potential,” *Phys. Rev. A* **78**, 023611.
- Shchukin, E, and W. Vogel (2005), “Inseparability criteria for continuous bipartite quantum states,” *Phys. Rev. Lett.* **95**, 230502.
- Sherson, J F, C. Weitenberg, M. Endres, M. Cheneau, I. Bloch, and S. Kuhr (2010), “Single-atom-resolved fluorescence imaging of an atomic Mott insulator,” *Nature* **467** (7311), 68–72.
- Shin, Y, M. Saba, T. A. Pasquini, W. Ketterle, D. E. Pritchard, and A. E. Leanhardt (2004), “Atom interferometry with Bose-Einstein condensates in a double-well potential,” *Phys. Rev. Lett.* **92**, 050405.
- Shor, P W (1995), “Scheme for reducing decoherence in quantum computer memory,” *Phys. Rev. A* **52**, R2493–R2496.
- Simon, R (2000), “Peres-Horodecki separability criterion for continuous variable systems,” *Phys. Rev. Lett.* **84**, 2726–2729.
- Sinatra, A, Y. Castin, and E. Witkowska (2013), “Limit of spin squeezing in trapped Bose-Einstein condensates,” *Europhys. Lett.* **102** (4), 40001.
- Sinatra, A, E. Witkowska, and Y. Castin (2012a), “Spin squeezing in finite temperature Bose-Einstein condensates: Scaling with the system size,” *Eur. Phys. J. Special Topics* **203**, 87.
- Sinatra, A, E. Witkowska, J.-C. Dornstetter, Yun Li, and Y. Castin (2011), “Limit of spin squeezing in finite-temperature Bose-Einstein condensates,” *Phys. Rev. Lett.* **107**, 060404.
- Sinatra, Alice, Jean-Christophe Dornstetter, and Yvan Castin (2012b), “Spin squeezing in Bose-Einstein condensates: Limits imposed by decoherence and non-zero temperature,” *Frontiers of Physics* **7** (1), 86–97.
- Slusher, R E, L. W. Hollberg, B. Yurke, J. C. Mertz, and J. F. Valley (1985), “Observation of squeezed states generated by four-wave mixing in an optical cavity,” *Phys. Rev. Lett.* **55**, 2409–2412.
- Smerzi, A, S. Fantoni, S. Giovanazzi, and S. R. Shenoy (1997), “Quantum coherent atomic tunneling between two trapped Bose-Einstein condensates,” *Phys. Rev. Lett.* **79**, 4950–4953.
- Smerzi, A, and S. Raghavan (2000), “Macroscopic quantum fluctuations in the Josephson dynamics of two weakly linked Bose-Einstein condensates,” *Phys. Rev. A* **61**, 063601.
- Smirne, A, J. Kołodyński, S. F. Huelga, and R. Demkowicz-Dobrzański (2016), “Ultimate precision limits for noisy frequency estimation,” *Phys. Rev. Lett.* **116**, 120801.
- Solano, E, R. L. de Matos Filho, and N. Zagury (1999), “Deterministic Bell states and measurement of the motional

- state of two trapped ions,” *Phys. Rev. A* **59**, R2539–R2543.
- Sorelli, G, L. Pezzè, and A. Smerzi (2015), “Fast generation of entanglement in bosonic Josephson junctions,” [arXiv:1511.01835 \[quant-ph\]](#).
- Sørensen, A S (2002), “Bogoliubov theory of entanglement in a Bose-Einstein condensate,” *Phys. Rev. A* **65**, 043610.
- Sørensen, A S, L.-M. Duan, J. I. Cirac, and P. Zoller (2001), “Many-particle entanglement with Bose-Einstein condensates,” *Nature* **409**, 63–66.
- Sørensen, A S, and K. Mølmer (1999), “Quantum computation with ions in thermal motion,” *Phys. Rev. Lett.* **82**, 1971–1974.
- Sørensen, A S, and K. Mølmer (2000), “Entanglement and quantum computation with ions in thermal motion,” *Phys. Rev. A* **62**, 022311.
- Sørensen, AS, and K. Mølmer (2001), “Entanglement and extreme spin squeezing,” *Phys. Rev. Lett.* **86**, 4431–4434.
- Spehner, D, K. Pawłowski, G. Ferrini, and A. Minguzzi (2014), “Effect of one-, two-, and three-body atom loss processes on superpositions of phase states in Bose-Josephson junctions,” *Eur. Phys. J. B* **87** (7), 1–22.
- Spehner, Dominique (2014), “Quantum correlations and distinguishability of quantum states,” *J. Math. Phys.* **55** (7), 075211.
- Spekkens, R W, and J. E. Sipe (1999), “Spatial fragmentation of a Bose-Einstein condensate in a double-well potential,” *Phys. Rev. A* **59**, 3868–3877.
- Stamper-Kurn, D M, and M. Ueda (2013), “Spinor Bose gases: Symmetries, magnetism, and quantum dynamics,” *Rev. Mod. Phys.* **85**, 1191–1244.
- Steane, A M (1996), “Error correcting codes in quantum theory,” *Phys. Rev. Lett.* **77**, 793–797.
- Steel, M J, and M. J. Collett (1998), “Quantum state of two trapped Bose-Einstein condensates with a Josephson coupling,” *Phys. Rev. A* **57**, 2920–2930.
- Sterling, R C, H. Rattanasonti, S. Weidt, K. Lake, P. Srinivasan, S. C. Webster, M. Kraft, and W. K. Hensinger (2014), “Fabrication and operation of a two-dimensional ion-trap lattice on a high-voltage microchip,” *Nat. Comm.* **5**, 3637.
- Streltsov, A I, O. E. Alon, and L. S. Cederbaum (2007), “Role of excited states in the splitting of a trapped interacting Bose-Einstein condensate by a time-dependent barrier,” *Phys. Rev. Lett.* **99**, 030402.
- Strobel, H, W. Muessel, D. Linnemann, T. Zibold, D. B. Hume, L. Pezzè, A. Smerzi, and M. K. Oberthaler (2014), “Fisher information and entanglement of non-Gaussian spin states,” *Science* **345**, 424.
- Stroescu, I, D. B. Hume, and M. K. Oberthaler (2015), “Double-well atom trap for fluorescence detection at the Heisenberg limit,” *Phys. Rev. A* **91**, 013412.
- Stute, A, B. Casabone, P. Schindler, T. Monz, P. O. Schmidt, B. Brandstatter, T. E. Northup, and R. Blatt (2012), “Tunable ion-photon entanglement in an optical cavity,” *Nature* **485** (7399), 482–485.
- Sudarshan, E C G (1963), “Equivalence of semiclassical and quantum mechanical descriptions of statistical light beams,” *Phys. Rev. Lett.* **10**, 277–279.
- Szańkowski, P, M. Trippenbach, and J. Chwedeńczuk (2014), “Parameter estimation in memory-assisted noisy quantum interferometry,” *Phys. Rev. A* **90**, 063619.
- Takahashi, Y, K. Honda, N. Tanaka, K. Toyoda, K. Ishikawa, and T. Yabuzaki (1999), “Quantum nondemolition measurement of spin via the paramagnetic Faraday rotation,” *Phys. Rev. A* **60**, 4974–4979.
- Takano, T, M. Fuyama, R. Namiki, and Y. Takahashi (2009), “Spin squeezing of a cold atomic ensemble with the nuclear spin of one-half,” *Phys. Rev. Lett.* **102**, 033601.
- Takano, T, S.-I.-R. Tanaka, R. Namiki, and Y. Takahashi (2010), “Manipulation of nonclassical atomic spin states,” *Phys. Rev. Lett.* **104**, 013602.
- Takeuchi, M, S. Ichihara, T. Takano, M. Kumakura, T. Yabuzaki, and Y. Takahashi (2005), “Spin squeezing via one-axis twisting with coherent light,” *Phys. Rev. Lett.* **94**, 023003.
- Tanji-Suzuki, H, I. D. Leroux, M. H. Schleier-Smith, M. Cetina, A. Grier, J. Simon, and V. Vuletić (2011), “Interaction between atomic ensembles and optical resonators: Classical description,” in *Advances in Atomic, Molecular, and Optical Physics*, Vol. 60, edited by E. Arimondo, P.R. Berman, and C.C. Lin, Chap. 4 (Academic Press) pp. 201–237.
- The LIGO Scientific Collaboration, (2011), “A gravitational wave observatory operating beyond the quantum shot-noise limit,” *Nat. Phys.* **7**, 962–965.
- Thomsen, L K, S. Mancini, and H. M. Wiseman (2002), “Spin squeezing via quantum feedback,” *Phys. Rev. A* **65**, 061801.
- Tobias, M, P. Hyllus, G. Tóth, C. Schwemmer, A. Niggelbaum, S. Gaile, O. Gühne, and H. Weinfurter (2012), “Permutationally invariant state reconstruction,” *New J. Phys.* **14** (10), 105001.
- Tóth, G (2012), “Multipartite entanglement and high-precision metrology,” *Phys. Rev. A* **85**, 022322.
- Tóth, G, and I. Apellaniz (2014), “Quantum metrology from a quantum information science perspective,” *J. Phys. A: Math. Theor.* **47**, 424006.
- Tóth, G, C. Knapp, O. Gühne, and H. J. Briegel (2007), “Optimal spin squeezing inequalities detect bound entanglement in spin models,” *Phys. Rev. Lett.* **99**, 250405.
- Tóth, G, C. Knapp, O. Gühne, and H. J. Briegel (2009), “Spin squeezing and entanglement,” *Phys. Rev. A* **79**, 042334.
- Tóth, G, and M. W. Mitchell (2010), “Generation of macroscopic singlet states in atomic ensembles,” *New J. Phys.* **12** (5), 053007.
- Tóth, G, W. Wieczorek, D. Gross, R. Krischek, C. Schwemmer, and H. Weinfurter (2010), “Permutationally invariant quantum tomography,” *Phys. Rev. Lett.* **105**, 250403.
- Trail, C M, P. S. Jessen, and I. H. Deutsch (2010), “Strongly enhanced spin squeezing via quantum control,” *Phys. Rev. Lett.* **105**, 193602.
- Trenkwalder, A, G. Spagnoli, G. Semeghini, S. Coop, M. Landini, P. Castilho, L. Pezzè, G. Modugno, M. Inguscio, A. Smerzi, and M. Fattori (2016), “Quantum phase transitions with parity-symmetry breaking and hysteresis,” *Nat. Phys.* **12** (9), 826–829.
- Treutlein, P, T. W. Hänsch, J. Reichel, A. Negretti, M. A. Cirone, and T. Calarco (2006), “Microwave potentials and optimal control for robust quantum gates on an atom chip,” *Phys. Rev. A* **74**, 022312.
- Truax, D R (1985), “Baker-Campbell-Hausdorff relations and unitarity of SU(2) and SU(1,1) squeeze operators,” *Phys. Rev. D* **31**, 1988–1991.
- Tura, J, R. Augusiak, A. B. Sainz, T. Vértesi, M. Lewenstein, and A. Acín (2014), “Detecting nonlocality in many-body quantum states,” *Science* **344** (6189), 1256–1258.

- Turchette, Q. A., C. S. Wood, B. E. King, C. J. Myatt, D. Leibfried, W. M. Itano, C. Monroe, and D. J. Wineland (1998), “Deterministic entanglement of two trapped ions,” *Phys. Rev. Lett.* **81**, 3631–3634.
- Uhlmann, A. (1976), “The ‘transition probability’ in the state space of a $\ast$ -algebra,” *Rep. Math. Phys.* **9**, 273.
- Ulam-Orgikh, D., and M. Kitagawa (2001), “Spin squeezing and decoherence limit in Ramsey spectroscopy,” *Phys. Rev. A* **64**, 052106.
- Ulyanov, V. V., and O. B. Zaslavskii (1992), “New methods in the theory of quantum spin systems,” *Phys. Rep.* **216**, 179.
- Vahlbruch, H., S. Chelkowski, B. Hage, A. Franzen, K. Danzmann, and R. Schnabel (2005), “Demonstration of a squeezed-light-enhanced power- and signal-recycled Michelson interferometer,” *Phys. Rev. Lett.* **95**, 211102.
- Van Trees, H. L., and K. L. Bell (2007), *Bayesian Bounds for Parameter Estimation and Nonlinear Filtering/Tracking* (Wiley-IEEE Press).
- Vanderbruggen, T., R. Kohlhaas, A. Bertoldi, S. Bernon, A. Aspect, A. Landragin, and P. Bouyer (2013), “Feedback control of trapped coherent atomic ensembles,” *Phys. Rev. Lett.* **110**, 210503.
- Vanier, J., and C. Audoin (1992), *The Quantum Physics of Atomic Frequency Standards* (Adam Higler, New York).
- Vasilakis, G., V. Shah, and M. V. Romalis (2011), “Stroboscopic backaction evasion in a dense alkali-metal vapor,” *Phys. Rev. Lett.* **106**, 143601.
- Vengalattore, M., J. M. Higbie, S. R. Leslie, J. Guzman, L. E. Sadler, and D. M. Stamper-Kurn (2007), “High-resolution magnetometry with a spinor Bose-Einstein condensate,” *Phys. Rev. Lett.* **98**, 200801.
- Vernac, L., M. Pinard, and E. Giacobino (2000), “Spin squeezing in two-level systems,” *Phys. Rev. A* **62**, 063812.
- Vitagliano, G., I. Apellaniz, I. L. Egusquiza, and G. Tóth (2014), “Spin squeezing and entanglement for an arbitrary spin,” *Phys. Rev. A* **89**, 032307.
- Vitagliano, G., I. Apellaniz, M. Kleinmann, B. Lücke, C. Klempt, and G. Tóth (2016), “Entanglement and extreme spin squeezing of unpolarized states,” [arXiv:1605.07202](https://arxiv.org/abs/1605.07202).
- Vitagliano, G., P. Hyllus, I. L. Egusquiza, and G. Tóth (2011), “Spin squeezing inequalities for arbitrary spin,” *Phys. Rev. Lett.* **107**, 240502.
- Volz, J., R. Gehr, G. Dubois, J. Estève, and J. Reichel (2011), “Measurement of the internal state of a single atom without energy exchange,” *Nature* **475**, 210–213.
- Walborn, S. P., B. G. Taketani, A. Salles, F. Toscano, and R. L. de Matos Filho (2009), “Entropic entanglement criteria for continuous variables,” *Phys. Rev. Lett.* **103**, 160505.
- Walls, D., and G. Milburn (1994), *Quantum Optics* (Springer, New York).
- Walls, D. F. (1983), “Squeezed states of light,” *Nature* **306** (5939), 141–146.
- Wang, X., and K. Mølmer (2002), “Pairwise entanglement in symmetric multi-qubit systems,” *Eur. Phys. J. D* **18**, 385.
- Wang, X., and B. C. Sanders (2003a), “Relations between bosonic quadrature squeezing and atomic spin squeezing,” *Phys. Rev. A* **68**, 033821.
- Wang, X., and B. C. Sanders (2003b), “Spin squeezing and pairwise entanglement for symmetric multiqubit states,” *Phys. Rev. A* **68**, 012101.
- Wasilewski, W., K. Jensen, H. Krauter, J. J. Renema, M. V. Balabas, and E. S. Polzik (2010), “Quantum noise limited and entanglement-assisted magnetometry,” *Phys. Rev. Lett.* **104**, 133601.
- Weinstein, J. D., K. Beloy, and A. Derevianko (2010), “Entangling the lattice clock: Towards Heisenberg-limited time-keeping,” *Phys. Rev. A* **81**, 030302.
- Welzel, J., A. Bautista-Salvador, C. Abarbanel, V. Wineman-Fisher, C. Wunderlich, R. Folman, and F. Schmidt-Kaler (2011), “Designing spin-spin interactions with one and two dimensional ion crystals in planar micro traps,” *Eur. Phys. J. D* **65** (1), 285–297.
- Werner, R. F. (1989), “Quantum states with Einstein-Podolsky-Rosen correlations admitting a hidden-variable model,” *Phys. Rev. A* **40**, 4277–4281.
- Wigner, E. (1932), “On the quantum correction for thermodynamic equilibrium,” *Phys. Rev.* **40**, 749–759.
- Wildermuth, S., S. Hofferberth, I. Lesanovsky, S. Groth, P. Krüger, J. Schmiedmayer, and I. Bar-Joseph (2006), “Sensing electric and magnetic fields with Bose-Einstein condensates,” *Appl. Phys. Lett.* **88** (26), 264103.
- Windpassinger, P. J., D. Oblak, P. G. Petrov, M. Kubasik, M. Saffman, C. L. Garrido Alzar, J. Appel, J. H. Müller, N. Kjærgaard, and E. S. Polzik (2008), “Nondestructive probing of Rabi oscillations on the cesium clock transition near the standard quantum limit,” *Phys. Rev. Lett.* **100**, 103601.
- Wineland, D. J. (2013), “Nobel lecture: Superposition, entanglement, and raising Schrödinger’s cat,” *Rev. Mod. Phys.* **85**, 1103–1114.
- Wineland, D. J., J. J. Bollinger, W. M. Itano, and D. J. Heinzen (1994), “Squeezed atomic states and projection noise in spectroscopy,” *Phys. Rev. A* **50**, 67–88.
- Wineland, D. J., J. J. Bollinger, W. M. Itano, F. L. Moore, and D. J. Heinzen (1992), “Spin squeezing and reduced quantum noise in spectroscopy,” *Phys. Rev. A* **46**, R6797–R6800.
- Wineland, D. J., C. Monroe, W. M. Itano, D. Leibfried, B. E. King, and D. M. Meekhof (1998), “Experimental issues in coherent quantum-state manipulation of trapped atomic ions,” *J. Res. Natl. Inst. Stand. Technol.* **103**, 259.
- Wódkiewicz, K., and J. Eberly (1985), “Coherent states, squeezed fluctuations, and the SU(2) and SU(1,1) groups in quantum-optics applications,” *J. Opt. Soc. Am. B* **2**, 458.
- Wolfgramm, F., A. Cerè, F. A. Beduini, A. Predojević, M. Koschorreck, and M. W. Mitchell (2010), “Squeezed-light optical magnetometry,” *Phys. Rev. Lett.* **105**, 053601.
- Woolley, M. J., G. J. Milburn, and C. M. Caves (2008), “Nonlinear quantum metrology using coupled nanomechanical resonators,” *New J. Phys.* **10** (12), 125018.
- Wootters, W. K. (1981), “Statistical distance and Hilbert space,” *Phys. Rev. D* **23**, 357–362.
- Wootters, W. K. (1998), “Entanglement of formation of an arbitrary state of two qubits,” *Phys. Rev. Lett.* **80**, 2245–2248.
- Wu, L.-A., H. J. Kimble, J. L. Hall, and H. Wu (1986), “Generation of squeezed states by parametric down conversion,” *Phys. Rev. Lett.* **57**, 2520–2523.
- Wynands, R. (2009), *Atomic Clocks*, edited by G. Muga, A. Ruschhaupt, and A. Campo, Lecture Notes in Physics, Vol. 789 (Springer Berlin Heidelberg, Berlin, Heidelberg).
- Wynands, R., and S. Weyers (2005), “Atomic fountain clocks,” *Metrologia* **42** (3), S64.
- Xiang, G. Y., B. L. Higgins, D. W. Berry, H. M. Wiseman, and G. J. Pryde (2011), “Entanglement-enhanced measurement of a completely unknown optical phase,” *Nat. Photon.*

- 5 (1), 43–47.
- Ye, J, H. J. Kimble, and H. Katori (2008), “Quantum state engineering and precision metrology using state-insensitive light traps,” *Science* **320** (5884), 1734–1738.
- Yurke, B, S. L. McCall, and J. R. Klauder (1986), “SU(2) and SU(1,1) interferometers,” *Phys. Rev. A* **33**, 4033–4054.
- Yuste, A, B. Juliá-Díaz, E. Torrontegui, J. Martorell, J. G. Muga, and A. Polls (2013), “Shortcut to adiabaticity in internal bosonic Josephson junctions,” *Phys. Rev. A* **88**, 043647.
- Zanardi, P, and M. Rasetti (1997), “Noiseless quantum codes,” *Phys. Rev. Lett.* **79**, 3306–3309.
- Zapata, I, F. Sols, and A. J. Leggett (1998), “Josephson effect between trapped Bose-Einstein condensates,” *Phys. Rev. A* **57**, R28–R31.
- Zare, R N (1988), *Angular Momentum* (Wiley-Interscience).
- Zehnder, L (1891), “Ein neuer Interferenzrefraktor,” *Z. Instrumentenkunde* **11** (8), 275–285.
- Zhang, H, R. McConnell, S. Ćuk, Q. Lin, M. H. Schleier-Smith, I. D. Leroux, and V. Vuletić (2012), “Collective state measurement of mesoscopic ensembles with single-atom resolution,” *Phys. Rev. Lett.* **109**, 133603.
- Zhang, J, K. Peng, and S. L. Braunstein (2003), “Backaction-induced spin-squeezed states in a detuned quantum-nondemolition measurement,” *Phys. Rev. A* **68**, 035802.
- Zhang, W-M, Da H. Feng, and R. Gilmore (1990), “Coherent states: Theory and some applications,” *Rev. Mod. Phys.* **62**, 867–927.
- Zhang, Y-L, C.-L. Zou, X.-B. Zou, L. Jiang, and G.-C. Guo (2015), “Detuning-enhanced cavity spin squeezing,” *Phys. Rev. A* **91**, 033625.
- Zhao, Z, T. Yang, Y.-A. Chen, A.-N. Zhang, M. Żukowski, and J.-W. Pan (2003), “Experimental violation of local realism by four-photon Greenberger-Horne-Zeilinger entanglement,” *Phys. Rev. Lett.* **91**, 180401.
- Ziń, P, J. Chwedeńczuk, B. Oleś, K. Sacha, and M. Trippenbach (2008), “Critical fluctuations of an attractive Bose gas in a double-well potential,” *Europhys. Lett.* **83**, 64007.